

The nitrogen book



Principles of soil nitrogen fertility management in southern Queensland and northern New South Wales farming systems

Includes easy-to-use electronic nitrogen fertiliser calculator

Compiled by: Howard Cox and Wayne Strong

© State of Queensland, 2015.

The Queensland Government supports and encourages the dissemination and exchange of its information. The copyright in this publication is licensed under a Creative Commons Attribution 3.0 Australia (CC BY) licence.



Under this licence you are free, without having to seek our permission, to use this publication in accordance with the licence terms.

You must keep intact the copyright notice and attribute the State of Queensland as the source of the publication.

Note: Some content in this publication may have different licence terms as indicated.

For more information on this licence, visit <http://creativecommons.org/licenses/by/3.0/au/deed.en>.

The nitrogen book

Principles of soil nitrogen fertility
management in southern Queensland and northern New South
Wales farming systems

Includes easy-to-use electronic nitrogen fertiliser calculator

Contents

Preface	iv
Acknowledgements	iv
Executive summary	1
1. Introduction	2
2. Understanding and managing soil biology	3
3. Gains and losses of soil nitrogen	9
4. Determining crop demand for nitrogen	20
5. The soil's capacity to supply nitrogen	31
6. Determining nitrogen fertiliser requirement	39
7. Alternative methods to calculate nitrogen fertiliser requirement	40
8. Evaluating the financial risk of applying nitrogen fertiliser in south Queensland	42
9. Applying nitrogen fertiliser in south Queensland	47
10. Nitrogen contribution of ley legumes and pulses	52
Glossary of terms	61
More information	61

Preface

Declining fertility of arable soils remains a problem of national and international significance. In eastern Australia's northern grain belt, soil erosion and nutrient removal are the main factors causing soil fertility decline.

In most cropping regions of northern Australia (Darling Downs, Western Downs and northern NSW) soil fertility rundown has occurred and needs to be corrected in order to obtain satisfactory yields in good seasons because in most areas the period of continuous grain cropping has exceeded after 40-50 years. Hence the frequency response to applied nitrogen is likely to be quite high, drought years not withstanding.

This manual details:

- identifying causes of variability
- methods to obtain a 'target' yield
- discussion of soil sampling
- alternative strategies to improve soil fertility.

In this manual, outputs from crop simulation models are used quite extensively. The reader is encouraged to obtain the program *WhopperCropper* for their own use. *WhopperCropper* is an easy way to visualise the full range of yield (and gross margin) outcomes that are possible. This allows the user to choose a strategy to match a targeted seasonal outcome in keeping with the grower's knowledge of paddock performance over a range seasons and attitude to managing financial risk.

A web-based version is in preparation; search for CropARM or contact 13 25 23 or visit www.daf.qld.gov.au

The grower may choose to manage for the lower end of the yield range, minimising costs but also limiting the potential for high yields. Alternatively the grower may apply fertiliser rates targeting a seasonal outcome with high returns in good seasons but with higher financial risk in poorer seasons.

Nitrogen management strategies and aids to guide the use of nitrogen fertilisers, designed to cope with declining soil fertility in southern Queensland and northern New South Wales, are described in this manual. The role of other nutrients is mentioned briefly but will be more extensively described in a subsequent publication.

Acknowledgements

This book arose as an extension of the 'The nitrogen book' produced as an initiative of the Central Queensland Sustainable Farming Systems Project. As such this product has been jointly funded by the Grains Research and Development Corporation and Department of Agriculture and Fisheries.

Special thanks go to the contribution from:

- Graham Spackman, Richard Routley, Michael Braunack, Maurice Conway and Anne Shepherd for editorial assistance.
- Chris Dowling (Back Paddock Company) for technical content
- Bruce Forster (Department of Natural Resources and Mines) for input regarding the soils of central Queensland
- Jo Owens and Giselle Whish for input regarding soils of southern Queensland
- David Lloyd, Brian Johnson, Sue O'Brien for information derived from their LeyGrain™ Pastures Workshop package
- Alastair Doherty and David Owens for assistance in preparation of the spreadsheet calculator.
- Heather Lees for desktop publishing and production.

CS4138 06/15

First published 2009
Republished 2015

Executive summary

Declining soil fertility through soil erosion and/or product removal has created the need for nitrogen addition. Vertosol soils have proved resilient to decline for lengthy periods but inevitably actions to improve or maintain soil physical, chemical and biological integrity will be required. Because of geological history, soil depth, length of farming, some soils are more vulnerable than others and need more immediate remedial activity. In some instances, return to pasture may be the most appropriate strategy.

Sustainable soil management currently focuses on the important issues of retention of surface residues, minimising wheeled traffic and managing soil nitrogen and other nutrients. Agronomic practices impact soil biota largely by the quantity and quality of crop residues returned to soil. Large amounts of residue increases the size of the potential carbon energy source for soil biota, since plant residues contain about 40% carbon. Periodic pasture leys and reduced tillage practices can benefit soil biota. Any short-term detrimental effect on biota from the increased use of herbicides, insecticides or fungicides appears to be out-weighed by wider ecological and crop production benefits. There is a large body of evidence demonstrating that many commercial fertilisers and animal manures increase rather than decrease soil biota and their activities in soil.

The major farming soils of the region comprise the grey, brown and black vertosols, black, red or brown sodosols, red and brown chromosols and ferrosols and some areas of red kandosols and black dermosols. The vertosols are generally of high clay content, chemically well buffered and appear resilient to negative impacts of nitrogen loss due to cultivation, crop removal and soil erosion. The other soil types are more 'fragile' and in some cases will support cropping only for short periods. Alternatively they are best utilised with livestock production.

Stored soil water and rainfall directly determine crop grain yield and crop grain yield determines the demand for nitrogen. This region has highly variable rainfall and hence yields and nitrogen demand will vary widely. There are two actions that can be taken to improve the management of risk involving nitrogen fertiliser application.

1. Measure or estimate soil water and soil nitrogen levels close to planting.
2. Use the *WhopperCropper* program to view the full range of potential yield outcomes. Scenarios are easily created for inputs (including nitrogen fertiliser). Having access to the full range of potential outcomes is superior to a calculation involving a single 'district average' because the user can readily evaluate how their attitude to risk is matched by the effect of different input on potential yield and gross margin outcomes.

Deep soil testing is the best method available for determination of soil nitrate-nitrogen but is still prone to inaccuracy. Carefully consider the number of cores that are used to get the representative sample.

After determining the soil nitrate level, the nitrogen fertiliser rate required can be calculated by the difference between the expected crop nitrogen demand and the soil nitrogen supply. Remember there may be an extra contribution from the soil in the time between soil sampling and planting.

WhopperCropper can also be used directly to view the potential yield ranges from different soil nitrogen levels and several potential nitrogen fertiliser rates.

After the soil nitrogen fertiliser rate is determined, source the cheapest or most convenient form of nitrogen that can be applied with the available equipment. Decide on application timing based on equipment available and work load requirements. The timing of the nitrogen application appears less important than satisfying the crop nitrogen demand. Observe the recommendations of the maximum quantity of fertiliser that can be placed with the seed or place nitrogen fertiliser at an appropriate distance from the seed row.

The cheapest form of nitrogen fertiliser should be sourced that is also appropriate to use with available equipment and for the timing of the application. Application at or prior to sowing is the most effective means of ensuring that the crop is able to readily access applied nitrogen. Nitrogen applied after sowing, when the soil surface is dry, may remain unavailable for crop uptake until the surface soil moisture has been re-wetted. When nitrogen fertiliser is applied at sowing, only a moderate rate should be applied with or in close proximity to the seed; recommendations of maximum quantity of nitrogen fertiliser that can be placed with the seed are provided. Where pre-sowing nitrogen application cannot be achieved, post-sowing application results in increased financial risks which are probably greater than revealed by outputs of *WhopperCropper*.

Pulse crops may contribute much less nitrogen (usually <30 kgN/ha) than a ley, particularly if grain removal is large by comparison to the vegetative material produced by the pulse crop. Leys or pulse crops may provide rotational benefits, like disease suppression, in addition to the variable accretion of nitrogen in soil.

1. Introduction

This manual covers important aspects of soil types, soil biology, organic carbon and management of factors pertinent to southern Queensland and northern New South Wales cropping industries. The nitrogen cycle and the flows between pools are described in detail. Losses and potential losses are detailed with the aim of maximising production whilst reducing fertility rundown.

High rainfall variability is well recognised in this region. In terms of managing risk it is important to understand how this rainfall variability affects crop yield and hence nitrogen demand. In this manual, outputs from simulation models, via the *WhopperCropper* program, are used extensively to generate seasonal production for wheat and sorghum crops including financial returns using weather data available from locations across the region over the past 100 years.

Whilst far from perfect, soil testing has been recognised as the primary means to determine plant-available nitrogen supply. Calculation of crop demand and soil supply of nitrogen and supplementary fertiliser requirements can be automated using the electronic calculator supplied with this manual.

Management options for timing, placement and safe application of nitrogen fertiliser are outlined. Also discussed is the integration of pastures with a legume component into cropping systems to offset soil nitrogen decline with continuous cropping.

Risk management with nitrogen fertilisers

To manage nitrogen nutrition and financial risk, two features of southern Queensland cropping need to be recognised;

- a) soil types differ in their original nitrogen status and in their capacities to supply plant-available nitrogen
- b) seasonal rainfall variation has a major impact upon financial risk associated with fertiliser application.

A successful strategy to manage nitrogen nutrition and financial risk must embrace these aspects of cropping as well as providing a platform from which advice can be modified/ fine tuned as new research findings come to hand. For this reason the current state of knowledge about use of nitrogen fertilisers and ley pastures in the region has been captured.

2. Understanding and managing soil biology

Key messages

- An enormous diversity of microbes exists in the soil, most in massive numbers.
- Because the majority of soil biota relies on carbon as their energy substrate, concentration of bio-available organic carbon in soil is associated with most changes in soil biota.
- Agronomic practices impact soil biota largely by the quantity and quality of crop residues returned to soil.
- The quantity of plant residues returned impacts the magnitude of potential energy source for soil biota.
- Nitrogen flows in the soil are complex and involve continuous movement of nitrogen between pools including biota.

Soil organic matter

Schwenke (2004) defined soil organic matter (SOM) as everything of biological origin whether living or dead. Both humus and organic carbon are components of the soil organic matter (Table 1). The above-ground portion of living plants is excluded. Humus is the most stable part of soil organic matter and is slow to break down.

Soil organic matter influences the biological, physical and chemical properties of soils which in turn provide ecological benefits. These include:

- Nitrogen storage, supply and cycling
- Food for microbes
- Cation exchange capacity
- Water-holding capacity infiltration and soil porosity
- Aiding soil aggregation
- Phosphorus storage
- pH buffering
- Chelation of micronutrients
- Pesticide degradation (substrate for microbes and chelation)
- Carbon sequestration
- Weed suppression (soil cover and allelopathy)
- Nematode and other disease organism suppression.

However, there can be downsides that include:

- allelopathic (growth inhibiting) chemicals
- hydrophobic (water repelling) substances
- nutrient tie up during decomposition of plant residues, particularly nitrogen and sulphur.

Soil organic carbon is an indicator of soil nitrogen fertility as well as an indicator of soils ability to support microbial populations.

Table 1. Components of soil organic matter (SOM)

Component	Percentage	Comment
Active SOM (living)	2 to 12%	see Table 2
Active SOM (non-living)	8 to 24%	Stubble, dead roots, manure, bones, sugars, amino acids, organic acids)
Stable SOM (humus)	70 to 90%	Sugars, amino acids, proteins, fats, lignin, other humic substances, charcoal

Table 2 Types and functions of soil microbes.

Type of microorganism	Function in soil
Organisms that add nutrients to soil	
Nitrogen-fixing micro-organisms Symbiotic N ₂ -fixing bacteria e.g. <i>Rhizobium</i> and <i>Bradyrhizobium</i> species	Fix atmospheric nitrogen in symbiosis with legume plants
Non-symbiotic N ₂ -fixing bacteria e.g. <i>Azospirillum</i> , <i>Azotobacter</i> species	Fix atmospheric nitrogen in bulk soil, near crop residues and in rhizosphere
Organisms that transfer nutrients into plant available forms or facilitate their uptake by plants	
Nitrifying microorganisms e.g. <i>Nitrosomonas</i> and <i>Nitrobacter</i> species	Convert ammonia nitrogen into plant available nitrate form
Sulfur-oxidizing micro-organisms e.g. <i>Thiobacillus thiooxidans</i> , most heterotrophic bacteria and fungi	Convert elemental sulfur and organic sulfur into plant-available sulfates and 'solubilise phosphates' (unlikely to occur in alkaline soils of CQ)
Mycorrhizae e.g. <i>Vesicular Arbuscular Mycorrhizae</i> (VAM) (except for crops such as canola)	Facilitate the uptake of phosphorus and zinc by most agricultural crops
Phosphorus-solubilising micro-organisms e.g. <i>Penicillium</i> species	Solubilise plant-unavailable inorganic and organic phosphorus into available forms
Organisms whose action results in the loss of nutrients from soil	
Denitrifying micro-organisms e.g. <i>Thiobacillus denitrificans</i>	Convert nitrate nitrogen into nitrogen and nitrous oxide gas
Sulfur-reducing bacteria e.g. <i>Desulfovibrio</i> species	Reduce sulfate sulfur into hydrogen sulfide gas
Organisms involved in the decomposition of crop residues	
Cellulolytic bacteria and fungi e.g. <i>Cellulomonas</i> species	Decompose cellulose and like compounds in crop residues
Organisms that promote above-ground and/or below-ground plant growth	
Plant growth promoting rhizobacteria e.g. <i>Pseudomonas</i> species, <i>Bacillus</i> species <i>Streptomyces</i> species	Promote above-ground and/or below-ground plant growth through hormone production or other mechanisms
Organisms involved causing plant diseases	
<i>Rhizoctonia solani</i> , <i>Pythium ultimum</i> , <i>Fusarium</i> species, <i>Verticillium</i> species, Ggt) <i>Bipolaris sorokiniana</i> ; <i>Fusarium graminearum</i> ; <i>Pyrenophora tritici-repentis</i>	Rhizoctonia barepatch, take-all, damping-off diseases. Common root rot Crown rot Yellow spot
Organisms involved in the control of plant diseases	
Bacteria: <i>Pseudomonas fluorescens</i> , <i>Bacillus subtilis</i> Fungi: <i>Trichoderma koningi</i> , <i>Fusarium oxysporum</i> Actinomyces: <i>Streptomyces rimosus</i>	Control soil-borne plant diseases

Source: Gupta and Roget 2004

Soil micro-organisms

Soil organisms (biota) are involved in transformation processes that are essential for crop production, soil quality and environmental health. There is a two way relationship between the soil biota and agricultural production; soil biota plays a key role in a number of nutrient transformation processes and crop residues supply carbon and nutrients to the soil biota. Soil biota also provide the following benefits:

- increased carbon and nitrogen transformations
- direct benefit to plants (nitrogen-fixing rhizobia)
- aiding soil stability (fungi filaments bind aggregates)
- competing with plant pathogens (disease suppression)
- providing short-term immobilisation of inorganic fertilisers.

Soil organisms can be grouped according to their size, morphological characteristics, function and food preference (Gupta and Roget 2004). Soil biota are also combined into groups based on their role in specific soil functions (Table 2). For example nitrifying micro-organisms are those that convert ammonia nitrogen into nitrate nitrogen, making it available to plants. Soil organisms range in size from microscopic (bacteria) to centimetres (earthworms).

The four major groups of soil biota, based on size are:

- microflora (bacteria, fungi, algae and actinomycetes)
- microfauna (protozoa, nematodes)
- mesofauna (collembola, mites)
- macrofauna (earthworms, beetles, termites).

In addition, soil fauna are also classified into various groups based on their principal food source, e.g. bacterial-feeding, fungal-feeding, plant parasitic or predatory fauna.

Micro-organism populations are generally enormous. The population will vary dependent on temperature, moisture and food supply. Only in extremely dry situations will most of the micro-organisms desiccate and die. Examples of population numbers are:

- bacteria: millions or trillions/gram of soil (400 to 5000 kg/ha)
- actinomycetes: (similar to bacteria) millions/gram of soil (400 to 5000 kg/ha)
- fungi: 1000 to 20 000 kg/ha
- algae: 1 to 10 billion/gram soil (10 to 500 kg/ha).

(from Brady and Weil 1996)

Specific bacteria associate with the particular conditions present in the soil e.g. wet or dry, with or without oxygen. Exchangeable calcium is important for their survival, as is soil pH.

Reduced tillage supports a fungal based system, whilst conventional tillage favours a bacterial-dominant system (Gupta and Roget 2004).

Carbon and nutrient cycling and nutrient availability

Gupta and Roget (2004) report the following summary of the contribution of microbial biomass:

“Organic matter in soil is the most important fraction that supports microbial populations, especially the biologically available portion of soil organic matter.”

Microbial biomass, the living component of soil organic matter, constitutes 2 to 7% of the organic carbon in soils. Microbial biomass acts as the engine for organic matter turnover and nutrient release. The size of microbial biomass carbon in the surface soil may range from 250 mg C/kg in a sandy soil to 1100 mg C/kg in a clay soil rich in organic matter. Microbial biomass carbon may only represent a small portion of soil organic matter (2 to 7%), but it is dynamic and living and thus is more sensitive to management practices than total soil organic matter.

Microbial biomass is a storehouse of plant-essential nutrients. For example, nitrogen levels in microbial biomass range from 15 to 150 kg N/ha. Microbial biomass also holds 5 to 15 kg/ha of sulphur and 10 to 45 kg/ha phosphorus. Nutrients held in microbial biomass are not prone to leaching, are tied up only temporarily, and are released for plant uptake as a result of predation by microfauna and the death of microbes during soil drying. It is the interactions between micro-organisms and organic matter in the soil that largely determine the fertility and overall quality of the soil. Therefore it is extremely important to use farm management practices that maintain organic matter levels, especially biologically available organic matter, in our soils’.

Effect of pastures

Plants are the major source of available carbon for biological activity, so soil biodiversity and biological activity depend on the quality and quantity of carbon inputs from plants, through root exudation and above and below-ground plant residues, and plant-induced changes in soil physical and chemical properties.

Pastures composed of mixtures of plant types (legumes, grasses) are considered to have a greater potential to influence diverse biological processes. This is due to the mixture of exudates and quantity and quality of residues. However, the quantity of carbon in grazed systems is affected by grazing management.

Effect of agronomic practices on soil biota

Because soil biota are dominated by heterotrophic organisms (those which rely on carbonaceous materials as an energy substrate), the concentration of organic carbon in soil, will in turn, affect the soil biota (Bunemann and McNeil 2004).

Australian soils are inherently low in biologically available carbon, so carbon inputs have a major influence on soil biological activity (Gupta and Roget 2004).

Agronomic practices impact soil biota largely by the quantity and quality of crop residues returned to soil. The quantity of plant residues returned directly affects the magnitude of the potential carbon energy source for soil biota, since all plant residues contain about 40% carbon. Residues of different chemical composition tend to favour different soil biota; for example, bacteria are favoured by inputs with high nitrogen concentration such as legume residues whereas cereal residues tend to increase populations of fungi. Crop residues decompose at varying rates so nitrogen availability will vary. There may even be large differences in nitrogen availability across various residues of the same species; woody or lignin containing residues are much less bio-available than herbaceous residues.

Several agronomic practices of northern farming systems are likely to impact soil biota and their activities in soil and some are discussed below:

- the occurrence of tillage and/or stubble retention
- the use of chemicals such as herbicides, insecticides and fungicides
- the application of fertilisers or manures.

Effect of tillage and stubble

Bell *et al.* (2004) found that overall microbial activity was less in soil continuously cropped (with a fallow between crops) than in soil with periodic leys. Reduced tillage results in less soil erosion, less exposure of soil organic matter to oxidation, and no dilution with subsoil material (Dalal and Chan 2001). However, reduced or no-till systems can also concentrate stubble and its contained nutrients within the uppermost layer of surface soil; only in the top 2.5 cm layer was there an increase in soil carbon detected during 8 years of no-till wheat with 75 kg N/ha applied annually on a Vertisol at Warra (Dalal *et al.* 1998). Also, it is likely that long fallows and/or bare areas in widely-spaced sorghum rows would result in reduced soil carbon inputs. Whether microbial populations are impacted by tillage and/or stubble retention will depend upon the quantity and quality of residues returned and weather conditions during its decomposition.

Effect of pesticides

Van Zwieten (2004) provides evidence that soil biota are affected by some but not all pesticides, ranging from negligible to large negative impacts but also with some positive impacts.

Immediate impacts can include short term stimulation of enzymatic activity, and bacterial numbers through to elimination of earthworm populations. It must be remembered that while pesticide effects may be evident in short-term laboratory tests, they may have little if any lasting effects in the field. Any short-term detrimental effect to soil biota may be out-weighed by wider ecological and crop production benefits. Specific examples of the effect of herbicides, insecticides and fungicides include the following (as cited by van Zwieten 2002):

- **Glyphosate:** bacterial numbers were reduced, fungi and actinomycetes were increased, overall increase in microbial activity of 9 to 19% (Araujo 2003); short term effects but no lasting changes to microbial community (Busse *et al.* 2001).
- **Glyphosate and paraquat:** activation of urease, suppression of phosphatase, enzymes that might impact upon action of micro-organisms, particularly soil bacteria (Sannino and Gianfreda 2001).
- **Atrazine:** urease activation, suppression of invertase (Sannino and Gianfreda 2001).
- **Atrazine and metolachlor:** altered community structure of bacteria and actinomycetes (Seghers *et al.* 2003).
- **Chlorpyrifos:** reduced bacterial numbers and slightly increased fungal numbers (Pandey and Singh 2004).
- **Copper:** reduced microbial biomass and OM breakdown; earthworms avoid soil with concentrations as low as 34 mg/kg (van Zwieten *et al.* 2004).

Effect of fertilisers

Effects of fertiliser additions on soil biota have been reported by Bunemann and McNeil (2004). They present a large body of evidence demonstrating that applications of many commercial fertilisers and animal manures increase rather than decrease soil biota and their activities in soil. Most effects of fertiliser addition on soil biota are immediate but may last up to 3 months or longer after the addition. The greater influence of organic versus inorganic fertilisers was emphasised in a study of soil respiration, acid phosphatase and dehydrogenase activity in maize. Three months after application of 200 kg N/ha as ammonium nitrate, the above components were greater than in the non-fertilised control. However, all these changes were more pronounced when the same amount of nitrogen was added as dairy manure or composted sewage sludge (Marinari *et al.* 2000).

Long-term application of phosphorus fertiliser often has little if any effect on soil biota. However, mineral nitrogen fertilisation has been reported to have a negative effect on soil biota in acid soils of South Australia and elsewhere. These negative effects are due to increased soil acidity created by addition of fertilisers such as ammonium sulphate and urea. However, addition of these fertilisers is unlikely to affect pH of central Queensland's well buffered neutral to alkaline vertosol soils and therefore should have little impact on soil biota.

A reduction in specific organisms such as arbuscular mycorrhizal (AM) fungi by phosphorus fertilisation appears to be fairly well established. In a comparison of Australian pastures under conventional and biodynamic management, a negative relationship existed between available phosphorus and colonisation rates of clover roots with AM fungi (Ryan *et al.* 2000), but AM colonisation rates of ryegrass were not affected by phosphorus addition (Ryan and Ash 1999). These findings agree with the variable effect of nitrogen, phosphorus, potassium (NPK) fertilisation on percent root colonisation by AM in different grassland species observed by Rillig *et al.* (1998). Research suggests that mycorrhizal symbioses are affected primarily by indirect effects of fertilisers through changes in plant growth and metabolism rather than by any direct effects on AM fungi.

Managing soil organic matter

Maintaining or increasing the amount of soil organic matter comes down to well recognised current best management practices:

- reduce soil erosion (results in less physical removal of nutrients)
- maintain stubble cover (Dalal and Chan 2001, Bell *et al.* 2004)
- avoid long fallows where possible (Bell *et al.* 2004) (maintains microbial population)
- grow healthy crops (adds biomass). May require use of inorganic fertilisers and pesticides
- apply well composted manure if available
- retain crop residues (no removal by burning or baling).

Can soil organic matter be increased?

The soil contains a massive reserve of organic matter. One hectare of soil to a depth of 10 cm weighs at least 1000 tonnes (assuming a soil bulk density of 1 g/cm³). If soil organic carbon (OC) is 1%, SOM is 1.7% or 17 t/ha in the top 10 cm. To increase OC to 2% (SOM = 3.5% or 35 t/ha) would require an additional 18 t/ha of OM. This could NOT be in the form of fresh manure because 80 to 90% of the product is lost over time (Schwenke 2004) although manures can be considered as valuable nutrient sources (especially for P and K).

A more successful way to increase soil organic matter is to incorporate a pasture phase into the cropping system. At *Warra* in south Queensland soil organic matter was increased by 20% (650 kg C/ha/year) after a 4-year pasture phase although this effect was likely to be short-lived because of the greater proportion of active materials added.

Building soil organic matter should be an aim in broadacre agriculture for soils that have very depleted soil organic carbon levels (less than 0.8% organic carbon). Growers with soils with this level or more may employ strategies to at least maintain moderate soil organic carbon levels assuming no physical constraints are limiting production. A pasture ley phase may suit some mixed farmers to build nutrient reserves. Pulse crops may 'save' some nitrogen depletion when conditions and prices favour their incorporation in the rotation.

References

- Abbott, L, and Murphy, D. 2004. Overview of 'soil biology' tests. In: Soil biology in agriculture. Proceedings of a workshop on current research into soil biology in agriculture. Tamworth Sustainable Farming Training Centre 11-12 August 2004. Ed Lines-Kelly R pp. 8-14. NSW Department of Primary Industries, Orange 2800.
- Araujo A.S.F, Monteiro R.T.R, Abarkeli R.B., 2003. Effect of glyphosate on the microbial activity of two Brazilian soils. *Chemosphere*, 52, 799-804.
- Bell, M., Seymour, N., Stirling, G., Van Zwieten, L., Sutton, G, and Moody, P (2004). Impact of management practices on activity of soil biota and productivity constraints in Vertosols of the northern grains region. In: Soil biology in agriculture. Proceedings of a workshop on current research into soil biology in agriculture. Tamworth Sustainable Farming Training Centre 11-12 August 2004. Ed Lines-Kelly R pp. 18-24. NSW Department of Primary Industries, Orange 2800.
- Brady N. C and Weil R. R 1996. Organisms and ecology of the soil. In: Brady N C and Weil, R. R (1996). The nature and properties of soils. Prentice-Hall International.
- Bünemann, E.K and McNeil, A (2004). Impact of fertiliser on soil biota. In: Soil biology in agriculture. Proceedings of a workshop on current research into soil biology in agriculture. Tamworth Sustainable Farming Training Centre 11-12 August 2004. Ed Lines-Kelly R pp. 64-69. NSW Department of Primary Industries, Orange 2800.
- Busse, M.D., Ratcliff, A.W, Shestak C.J, and Powers RF 2001. Glyphosate toxicity and the effects of long-term vegetation control on soil microbial communities. *Soil Biology & Biochemistry* 33, 1777-1789.
- Dalal, R.C and Chan K.Y 2001. Soil organic matter in rainfed cropping systems of the Australian cereal belt. *Australian Journal of Soil Research* 39 435-464.
- Gupta VVSR and Roget David K 2004. Understanding soil biota and biological functions: Management of soil biota for improved benefits to crop production and environmental health. In: Soil biology in agriculture. Proceedings of a workshop on current research into soil biology in agriculture. Tamworth Sustainable Farming Training Centre 11-12 August 2004. Ed Lines-Kelly R pp. 72-78. NSW Department of Primary Industries, Orange 2800.
- Dalal, R.C., Strong, W. M., Weston E. J., Cooper J. E., Wildermuth, G. B., Lehane K. J, King, A. J. and Holmes C. J. (1998) Sustaining productivity of a vertisol at Warra, Queensland, with fertilisers, no-tillage or legumes 5. Wheat yields, nitrogen benefits and water-use efficiency of chickpea-wheat rotation. *Australian Journal of Experimental Agriculture*, 1998, 38, 489-501.
- Marinari S, Masciandaro G, Ceccanti B, Grego S 2000. Influence of organic and mineral fertilisers on soil biological and physical properties. *Bioresource Technology* 72, 9-17.
- Pandey, S., Singh, D.K. 2004. Total bacterial and fungal population after chlorpyrifos and quinalphos treatments in groundnut (*Arachis hypogaea* L.) soils. *Chemosphere* 55, 197-205.
- Rillig, M.C, Allen, M.F, Klironomos, J.N, Chiariello, N.R, and Field, C.B. 1998. Plant species specific changes in root-inhabiting fungi in a California annual grassland: Responses to elevated CO₂ and nutrients. *Oecologia*, 113, 252-259.
- Ryan, M.H., Small, D.R and Ash, J.E. 2000. Phosphorus controls the level of colonisation by arbuscular mycorrhizal fungi in conventional and biodynamic irrigated dairy pastures. *Australian Journal of Experimental Agriculture*, 40 663-670.
- Ryan, M and Ash, J 1999. Effects of phosphorus and nitrogen on growth of pasture plants and VAM fungi in SE Australian soils with contrasting fertiliser histories (conventional and biodynamic). *Agriculture Ecosystems & Environment*, 73, 51-62.
- Sannino F, Gianfreda L 2001. Pesticide influence on soil enzymatic activities. *Chemosphere* 45 417-425.
- Schwenke, G 2004. Soil organic matter, biological activity and productivity: myths and realities. In: Soil biology in agriculture. Proceedings of a workshop on current research into soil biology in agriculture. Tamworth Sustainable Farming Training Centre 11-12 August 2004. Ed Lines-Kelly R pp. 72-78. NSW Department of Primary Industries, Orange 2800.
- Seghers D, Verthe, K, Reheul, D, Bulcke, R, Siciliano, S.D, Verstraete, W, Top E,M. 2003. Effect of long-term herbicide applications on the bacterial community structure and function in an agricultural soil. *FEMS Microbiology Ecology*, 46 139-146.
- Van Zwieten, L.M 2004. Impact of pesticides on soil biota. In: Soil biology in agriculture. Proceedings of a workshop on current research into soil biology in agriculture. Tamworth Sustainable Farming Training Centre 11-12 August 2004. Ed Lines-Kelly R pp. 72-78. NSW Department of Primary Industries, Orange 2800.
- Van Zwieten, L., Rust, J., Kingston, T., Merrington, G., and Morris, S 2004. Influence of copper fungicide residues on occurrence of earthworms in avocado orchard soils. *Science of the Total Environment*, 63, 59-68.

3. Gains and losses of soil nitrogen

Key messages

- Soil that had brigalow trees as its native vegetation usually had a high level of nitrogen fertility (organic carbon of usually greater than 2%) before being used for farming.
- Soil that supported predominantly grass vegetation had a lower nitrogen fertility (organic carbon levels of 1 to 1.5%) because of the lack of a nitrogen fixing vegetation.
- The rate of decline of nitrogen fertility is often faster in other soil types and hence will require nitrogen addition sooner than Brigalow/softwood scrub soils.
- The onset of nitrogen fertiliser response could occur after 5 to 30 years of cropping.
- Soils will mineralise 50 to 100 kg N/ha per year.
- Flow rates of nitrogen between pools vary from very slow (immobilisation or mineralisation) to very fast (denitrification), are all affected by weather conditions, and are difficult to measure independently.
- Vertosol soils are chemically well buffered and appear comparatively resilient to negative impacts of nitrogen loss due cultivation, crop removal and soil erosion.
- Nitrogen fertiliser application in conjunction with reduced tillage practices can slow the decline of soil nitrogen.
- Loss of nitrogen due to denitrification can occur quickly removing up to 40% of nitrate nitrogen from top-soil, but only if three conditions coincide; waterlogging, nitrate and fresh crop residues.
- Soil organic carbon level is an indicator of current nitrogen fertility status, reflecting the capacity of the soil to supply plant-available nitrogen and therefore the potential to respond to nitrogen fertiliser.

Cropping soils of southern Queensland

Grain-growers and advisers frequently describe soils and their fertility by association with the original vegetation. Below are listed, in approximately descending order of original nitrogen fertility, four major soil/vegetation associations:

- Mixed brigalow soils
- Basaltic uplands
- Alluvial soils
- Sodosols and sands.

The original vegetation type distinguishes the brigalow scrub soils, with generally higher initial fertility, from the basaltic upland and alluvial soils with generally lower initial fertility status. The difference in soil fertility reflects the ability of brigalow softwood scrub to maintain a higher total soil nitrogen level than the grass vegetation of alluvial plain soils. However, the plains soils are highly productive because of the massive water-holding capacity and have slower rundown of nitrogen fertility.

Soils need to be able to store at least 120 mm of plant available water within their rooting depth for reliable dry-land cropping. Rooting depth can be limited by the depth of soil to an impervious barrier such as rock or to a zone in the subsoil with a high concentration of soluble salts (greater than 0.8 dS/m) and/or sodic conditions (greater than 15% exchangeable sodium percentage). These zones may be referred to as the salt bulge or sodic bulge. Crops are unable to use the water present in layers of soils with these undesirable conditions for deep root penetration.

Cracking clay soils


Cracking clay soils are the most important and widespread soils used for dry-land cropping in southern Queensland. They occur on a variety of landscapes and parent materials, and are suitable for cultivation provided they are deep enough, do not have large melon holes (gilgais) and are not prone to flooding. The shrinking and swelling nature of cracking clay soils assists with recovery from compaction.

Soil types vary within districts and there is a broad variation from east to west in south Queensland. For greater detail readers should access the appropriate Land Management Manuals for their area. These publications are generally available from Natural Resource and Water offices.

The soils of south Queensland can be divided into the following land resource areas (Harris *et al.* 1999, Maher 1996):

(a) Alluvial plains

Alluvial plains support large areas of cropping in the south Queensland region. They are level to gently sloping. In the Condamine floodplain there are two subclasses:



Map 1. Distribution of major dryland cropping soils of southern Queensland

Source: Department of Natural Resources and Mines

Older alluvial plains that are elevated and not subject to overland flow but are continually eroded from adjacent catchments. Some soil types include: Waco, Cecilvale and Oakey.

Recent alluvial plains are dominated by levees, terraces, oxbows and prior streams. Some soil types include: Condamine, Anchorfield, Mywybilla. Vegetation consists of very open grassy woodland of poplar box with coolabah, river red gum on drainage lines.

(b) Brigalow uplands and poplar box walloons

Formed from weathering of the soft walloon coal measures resulting in long undulating plains and occasional undulating rises. Some common soil include: Moola, Diamondy, Acland, Kenmuir and Elphinstone. Vegetation of shrubby open forest of brigalow, belah and wilga with some areas of poplar box occur.

(c) Brigalow plains

Brigalow plains occupy large areas to the west of the central Darling Downs including into the Tara-Chinchilla district; generally these are flat to gently undulating with shallow to deep gilgais. Vegetation can include: brigalow, belah, wilga and false sandalwood. Some poplar and black tea tree on poorer drained areas. Soil types include Kupunn and Tara.



Typical depiction of recent or older alluvial land type narrow valleys of the broad level plains ($\leq 2\%$ slope) of mixed basaltic and sandstone alluvium

Source: Department of Agriculture and Fisheries (2008). Land Types of Queensland – Darling Downs. Version 1.0. Prepared by the Grazing Land Management workshop teams, Department of Primary Industries and Fisheries, Brisbane.



Gently undulating to steep low hills and rises typical of the brigalow uplands land type

Source: Department of Primary Industries and Fisheries (2008). Land Types of Queensland – Darling Downs. Version 1.0. Prepared by the Grazing Land Management workshop teams, Department of Primary Industries and Fisheries, Brisbane.



Flat to gently undulating to steep low hills and rises typical of the brigalow plains land type

Source: Maher (1996) Understanding and managing soils in the Murilla, Tara and Chinchilla Shires, Department of Primary Industries and Fisheries Training Series QE960001, Brisbane.



Mountain coolibah open woodland with grassy understorey typical of the undulating rises and rolling low hills on basaltic uplands

Source: Department of Primary Industries and Fisheries (2008). Land Types of Queensland – Darling Downs. Version 1.0. Prepared by the Grazing Land Management workshop teams, Department of Primary Industries and Fisheries, Brisbane.

(d) Basaltic uplands

Is a complex landuse area that includes the plateau around Toowoomba. Mostly undulating plains and rises with some steep hills. Soil types include: Craigmore, Irving, Charlton, Purrawunda, Kenmuir and Drayton.

Sodosols and sands

These soils are unsuitable for cultivation and include areas of ironbark/bull oak sodosols, Cypress pine sands, sandstone forests, granite hills and traprock hills. Soil types include Weranga, Braemar, Davy, Banca and Gammie.

Indicators of soil nitrogen fertility

Soil organic matter content

Most southern Queensland soils contained large quantities of organic matter at the commencement of cropping. This quantity has declined over the duration of cropping, particularly in the topsoil from which crops derive most nutrients and much of the crop's water supply. Because of the high clay content (50 to 70%) of most southern Queensland soils, the organic matter decline is less than would be for other soil types located in a tropical region. Decline in soil organic matter of brigalow or softwood scrub soils appears initially faster than for open grassland soils, due possibly to the lower clay content which is known to offer some protection from decomposition of soil organic matter.

Most arable southern Queensland soils contain 170 t/ha or more of organic matter to a depth of 1 m, of which 58% is organic carbon. Soil organic matter is most frequently measured by the quantity of soil organic carbon present. Up to 50% of the organic matter resides in the upper 30 cm of soil.

Soil organic carbon

Soil organic carbon (OC) is the laboratory test that measures the quantity of soil organic matter (SOM). Hence organic carbon can be used as an indicator of soil fertility status. Because OC % can be measured using a relatively inexpensive oxidation procedure it is commonly used as a surrogate measure of soil nitrogen status.

SOM contains approximately 58% carbon (C) thus its concentration in soil can be estimated from OC content as follows:

$$\begin{aligned} \text{SOM} &= \text{OC} * 1 / 0.58 \\ &= \text{OC} * 1.72 \end{aligned}$$

Total nitrogen content

Total nitrogen content of soil in the top 10 cm can be measured directly using a standard laboratory process but is also frequently estimated from soil organic carbon assuming an approximate ratio of C:N of 10:1. Soils of southern Queensland may vary from less than 0.1 to more than 0.2% total nitrogen. The lower values are generally from open grassland soils, whilst the higher values are usually indicative of brigalow/softwood scrub soils.

Total nitrogen identifies the soil's potential to supply plant-available nitrogen. More than 90% of the soil's total nitrogen may be initially unavailable to the crop because it exists in organic forms. Total nitrogen therefore is a measure of the soil's capacity to supply plant-available nitrogen over the long term.

Plant-available nitrogen (mineral-nitrogen)

Plant-available nitrogen is best determined by soil tests that measure the forms of soil nitrogen referred to as mineral nitrogen. Mineral nitrogen is principally nitrate-nitrogen because in the northern region, transformation of organic-nitrogen to nitrate-nitrogen through an intermediate ammonium-nitrogen, phase is very rapid.

Ammonium nitrogen is a very temporary phase and detected in field moist soil at only very low concentrations (<1 mg/kg). Disregard any ammonium soil test value conducted on air-dried (40°C) soil because the value of ammonium-nitrogen will usually be artificially high. This occurs because air-drying promotes transformation of organic-nitrogen to the ammonium nitrogen form. Ammonium-nitrogen tests, if required, must be conducted quickly on field-moist soil.

Because the level of ammonium-nitrogen present in unfertilised field-moist top-soil is usually very low (<1 kg N/ha) it is generally disregarded in calculating plant-available nitrogen supply in a nitrogen budgeting calculations.



Figure 1. The complete nitrogen cycle

The nitrogen cycle

The full nitrogen cycle is shown in Figure 1. The flows are continuous and rapidly change in size. Of particular interest to farming systems is the flow from organic matter to mineral-nitrogen because of the large quantity of nitrogen involved.

Nitrogen transformations

The process of mineralisation involves:

- The decomposition of soil organic matter by microbes to release inorganic (mineral) forms of nitrogen (initially ammonium) and water (Figure 2). Soil nitrate is the inorganic form of nitrogen that is available for plant uptake whilst organic nitrogen is not.
- A reverse process from mineral nitrogen to organic nitrogen can occur (called immobilization)
- The enzymatic oxidation from ammonium to nitrite is undertaken by the nitrosomonas bacteria whilst the conversion from nitrite to nitrate occurs via the nitrobacter bacteria
- In southern Queensland soils approximately 50 to 100 kg N is mineralised annually




Figure 2. Major nitrogen transformations in the soil-plant system




Figure 3. Major outputs/losses of nitrogen in the soil-plant system

- The mineralisation rate is greatest under warm (25 to 35°C), moist (neither dry nor water-logged) conditions with sufficient aeration
- After a phase when nitrogen rundown is initially rapid, the amount of inorganic nitrogen released is approximately proportional to OC% (SOM) content
- i.e. soil with 2% OC will mineralise twice as much nitrogen annually as a soil with 1% OC.
- Removal in produce (grain, fibre, meat and wool), erosion of topsoil, gaseous losses as ammonia (NH₃) and oxides of nitrogen (chiefly N₂O, N₂)
- Gaseous loss of nitrogen as ammonia may occur from surface applied ammonium-forming nitrogen fertilisers (urea, ammonium nitrate, anhydrous ammonia). Ammonia may also be lost from cereal crops during the grain filling stage when nitrogen is being translocated from vegetative parts to grain.
- Nitrous oxide (N₂O), dinitrogen (N₂), and other nitrogen oxides may be emitted from soil to the atmosphere when soil is waterlogged, resulting in a denitrification process mediated by soil-borne organisms.

Losses of nitrogen

Nitrogen removal in crop produce, and that lost by soil erosion and denitrification can account for the largest amounts of nitrogen loss (Figure 3 and Table 3). The quantity nitrogen leached through the profile is usually quite small (Radford *et al.* 2008).

The quantity removed in crops will depend upon the rainfall which is a significant driver of crop yield. Knowledge of the inherent soil nitrogen fertility will determine if nitrogen application is urgently required or can be postponed. On soils that are depleted in nitrogen, it may be advisable to supply nitrogen at a rate equal to potential removal rates.

With the exception of high intensity rainfall events, soil erosion has been significantly reduced by adoption of reduced tillage and controlled traffic systems, retention of stubble and contour banks.

Denitrification can be a significant loss under conditions where all three criteria of influence occur coincidentally; presence of carbon residues, waterlogged soil and presence of quantities of nitrate from soil or fertiliser sources (Table 4).

Other losses from the soil are negligible compared to the losses from crop removal, erosion and denitrification.

Processes that result in depletion of soil nitrogen include (Figure 5 and Table 3):

Denitrification

The loss of plant available nitrogen from topsoil usually occurs only occasionally from heavy clay soils in southern Queensland. Nitrate–nitrogen may be lost when heavy clay soil is waterlogged for periods of 24 hours or longer, at high soil temperatures and when nitrate-nitrogen and plant residues are both present. Obviously this combination of factors does not occur frequently, so loss of plant available nitrogen should be a low risk to growers. The factors that must be present to result in a high risk of denitrification loss are shown in Table 4. Because of the requirement for easily decomposable organic matter, denitrification mainly occurs in the surface soil.

It can be seen from the Table 4 that when nitrogen fertiliser is applied during summer closely following harvest of a previous cereal crop, and when all 3 risk-factors coincide, only then will this loss be of greatest concern to growers. At other times, at least one risk-factor is likely to be absent or of little influence.

Even then, application of nitrogen may be so essential to supplying the needs of a double cropped cereal, the grower may be prepared to take the risk and apply nitrogen fertiliser. The need to apply nitrogen to double crop a cereal has not

Table 3. Sources, frequency and magnitude of nitrogen loss from the soil

N loss or process	Source	Frequency	Magnitude	Comment
Removal of produce	Grain / silage / hay	Regular	Gross	
Displacement of soil	Soil erosion by water	Episodic	Gross	
Leaching of solute	Soil nitrate and fertiliser	Episodic	Variable – unknown below rooting	Dominated by soil permeability
Gaseous N loss Denitrification	Topsoil nitrate soil or fertiliser	Episodic	Variable >10%	Dominated by soil drainage
Ammonia emissions from soil	Ammoniacal fertilisers	Regular	Variable <25% applied	Application method and weather
Ammonia emissions from plants	Plant N content	Regular	<1-15 kg/ha/yr	High N% / senescence -net loss

Source: Dr W Strong (pers. comm.)

been a common occurrence, so rarely will the grower have to balance the risk of losing some nitrogen (up to 40%) to increase the plant-available nitrogen requirements for the next cereal.

Alternatively, growers may choose to apply nitrogen during late summer or early autumn for the forthcoming winter crop to spread the work load or to obtain discounted fertiliser prices. The risk factors in doing so should be taken into account.

It is sometimes stated that there is an advantage in applying nitrogen fertiliser early as it will allow applied nitrogen to move with rainfall to deeper soil layers and hence lengthen the window of availability to a dryland crop. This concept is flawed for two reasons. Firstly, rainfall events necessary to move applied nitrogen deeper into heavy clay soil can also create loss of applied nitrogen as described above. Secondly, more than 200 mm of rain is generally required to displace applied nitrogen to depth greater than 20 cm in clay soils.

Quantifying potential losses from nitrogen applied in late summer

The highest likelihood of nitrogen loss occurs from nitrogen fertilised soil between November and March when a high level of crop residues is present and when the likelihood of waterlogging is high.

On the Darling Downs (Strong and Cooper 1992) reported that anhydrous ammonia was completely converted to nitrate-nitrogen in 11 days after application. The nitrogen was applied in February and only 8 mm of rain received after application.

Losses as high as 71% have been reported with waterlogging created by irrigation but in dryland systems, substantial loss (30 to 60%) of applied nitrogen has been measured during summer fallow (Avalakki *et al.* 1995).

Nitrogen application during drought periods

In periods of unpredictable rainfall it might be economical to withhold the application of nitrogen to crops with high risk of crop failure, principally those with low soil water at planting and/or moderate soil nitrate levels. Applying nitrogen to crops at lower risk or in more favourable seasons (high soil water at planting and low soil nitrate) may yield a more economical result. Expect some carryover benefit of applied nitrogen to crops following failed fertilised crops but inevitably losses of up to approximately 15% can occur.

Simulated crop outputs are useful to demonstrate the range of potential yield outcomes. The concept of the 'expected' yield relies on the correct 'guess' regarding the rainfall of the forthcoming season. Thus there is no guarantee that using crop simulations will result in less financial loss unless the grower chooses the 'correct' grain yield expectation. Low expected yields generally discourage nitrogen fertiliser application. Thus, using low input levels should reduce financial loss where actual yield outcomes are moderate to low, but may sacrifice potential profit when infrequent high yields occur.

Nitrate leaching

At 13 sites in southern Queensland drainage under cropping averaged 8 mm/year (Tolmie *et al.* 2003). The annual rate of deep drainage under native vegetation was lower, averaging 0.3 mm/year on grey vertosols and 1 mm/year on black vertosols. In a similar study in central Queensland, nitrate-N had been leached below crop rooting depth at only three of seven sites. Thus it appeared that although water appears to be moving through soil profiles there is little chance that significant nitrogen will be lost via this process.

It was also reported that drainage for farming systems currently practiced in southern Queensland (less tillage, more summer/opportunity crops) was about half that of farming systems involving high level of tillage, less stubble retention and a wheat-dominant cropping system (Silburn *et al.* 2008).

Gains of nitrogen

Several processes result in soil nitrogen accumulation:

- dominant potential sources of nitrogen input into southern Queensland cropping soils are nitrogen fertilisers and legumes (Figure 4 and Table 5)
- a small quantity (<10 kg/ha/year) may derive from lightning strikes
- a small (unknown) quantity may derive from processes of stubble decomposition and asymbiotic nitrogen fixation, processes mediated by soil-borne organisms.

Other contributing sources are complex and very difficult to measure. They contribute only a small amount of nitrogen to the system and operate in a continual state of flux with the quantities contributed being small compared to the gross amounts that are required by crops or lost by erosion or denitrification.

Soil contains from 2% to 12% of its organic matter as living microbes; equivalent to 20 to 120kg/ha. Populations of the native biota will flourish under conditions of adequate water and energy (carbon) supplies. Management that facilitates these conditions will promote microbial populations even without additional amendments.

Note: Soil nitrate is NOT a soil fertility measure. Soil nitrate levels constantly change in the soil, due to losses and gains that occur throughout a fallow and whilst soil nitrate does estimate supply of plant-available nitrogen for the forthcoming crop, it is not a measure of the inherent soil nitrogen fertility.

Soil fertility decline

There is worldwide recognition that the fertility of arable soils is in decline.

Within Australia, evidence of declining soil fertility, crop production, and grain quality has been reported by Dalal and Mayer (1986a) (Figure 5). Soil fertility declines with increasing time in cultivation. Using anaerobic mineralisable N as an indicator, Dalal and Mayer (1990) proposed that crops would respond to applied nitrogen fertiliser after 5 to 15 years of commencing grain cropping.

Growers in southern Queensland appear well aware of the issues of fertility rundown and nitrogen fertiliser requirement. Lawrence *et al.* (2000) stated all attendees at a nitrogen fertiliser management workshop reported using nitrogen fertiliser at an average rate of 37 kgN/ha. Admittedly these were farmers motivated to attend such a workshop but in most areas of southern Queensland some form of nitrogen application is considered necessary in most years. The dilemma for southern Queensland growers is how to continue profitable cropping regardless of soil fertility decline.

Table 4. Risk factors that may promote denitrification N losses from soil, waterlogging, nitrate and crop residues that may occur during summer and winter seasons.

Factor	Summer		Winter	
	Nil N	+N fert	Nil N	+N fert
Waterlogging	High	High	Low	Low
Nitrate supply	Low	High	Low/med	High
Residues	High	High	Low/med	Low/med
Overall losses	Low	Potentially high loss	Low	Low

Source: Strong *et al.* (1996)

Cornish *et al.* (1998) and Spackman and Garside (1995) have reported that wheat yield in 39 shires was related to low nitrogen fertility and in some cases phosphorus deficiency. Furthermore, improved soil fertility is linked to increased water use efficiency, an essential requirement of Australia's grain production systems (Cornish *et al.* 1998).

Restoring soil organic matter

Soil nitrogen fertility decline will generally occur with continuous cereal cropping. The introduction of a grass/legume phase is the only way to significantly reverse the soil fertility decline. Addition of a legume (pulse) crop may slow the process but is unlikely to eliminate it. Agronomic practices such as no-till and controlled traffic that reduce soil erosion have been shown to increase soil organic matter only when nitrogen fertiliser was applied and stubble was retained. (Wang *et al.* 2004).

Dalal (1995) reported that an increase of 650 kgC/ha/year was recorded in a vertosol by a grass+legume pasture for four years. The organic C increase was attributed to input from the grass root biomass (10 t/ha/year compared the continuous wheat of 2 t/ha/year). Nitrogen fixed by the legume plays an important part in this increase because of the increase biomass production of the grass and the 'locking up' of nitrogen in the grass, thus reducing potential losses from denitrification.

Conversely, short (two year) phases of lucerne-wheat or medic-wheat had a negligible effect on soil organic carbon.

Fate of applied nitrogen

The quantity of in-crop rainfall affects the removal, loss and apportionment of nitrogen fertiliser applied to a cereal crop in farming systems of northern Australia

Results from a trial quantifying the recovery of nitrogen fertiliser labelled with N15 and its loss from the system when applied to wheat at sowing at Warra, in southern Queensland are shown in Figure 5.



Paddock conditions likely to induce denitrification

- In a dry year (1990) grain removal was only slightly lower but loss to the atmosphere was very low and the amount recovered in soil higher.
- In a wet year (1988) loss was much higher and the amount recovered in soil was reduced. In all seasons, fertiliser recovered in soil was mainly in organic forms.
- Total loss of the nitrogen fertiliser was 5 to 25% depending upon the seasonal rainfall.

Carryover of fertiliser nitrogen after failed crops

If conditions remain dry following a failed crop, it is likely that a significant amount of applied nitrogen fertiliser will be available for the next crop. Armstrong and Halpin (1993) reported that following wheat crop sown in June and killed immediately after emergence, that 75% and 87% of the original nitrogen fertiliser was recovered in the soil in September. Rainfall during the growing season totalled only 29 mm. Similarly, sorghum in 1993 took up as much ¹⁵N-labelled fertiliser applied 12 months previously as it did for freshly applied fertiliser (Armstrong *et al.* 1996).

Interaction between nitrogen and other elements

In this section, the effect of phosphorus, zinc, and potassium are described briefly in relation to effective nitrogen management. The general principle is that other nutrients must be in a non-limiting supply in order for the full response from nitrogen fertiliser to be realised. Nitrogen fertiliser is usually the most expensive input and the response is rate-related i.e. up to a point the more that is applied the greater the yield (water not limiting).

- Is important that the supply of other elements is non-limiting to enable a crop to respond to applied nitrogen fertiliser.
- Crop response to applied N should not be affected unless soil bicarbonate test phosphorus level is below 25 mg/kg, (Darling Downs or eastern areas, or 15 mg/kg western regions). These levels indicate need for P application, but response will still depend upon rainfall received by the crop.

- The critical soil potassium level is 0.4 meq/100g. Most Queensland vertosol soils currently contain higher levels of potassium and so N response should not be negatively impacted.
- The critical soil sulphur level is approximately 5 mg/kg. Soil surface levels are sometimes less than this but crops may access large sulphur reservoirs deeper in the soil. Low sulphur levels are likely to negatively impact crop N response only when cereals are double cropped.
- The zinc critical level is usually reported as 0.8 mg/kg for soils with pH greater than 7. Grain yield responses to zinc are highly variable and not always reflected by soil zinc test. Plant tissue analysis is accepted as a more reliable indicator of zinc adequacy for the plant. After a long period of continuous cropping, a strategy to apply modest application of zinc may avoid negative impact on crop response to applied N; where applied P is required, zinc can be efficiently co-applied.

Classifying soils according to original vegetation does not usually help to identify most other soil nutrients and characteristics that may impact on long-term crop and pasture performance. Soil testing for the elements in question is necessary to identify inadequate plant-available supplies of most nutrients or other soil dysfunctions in southern Queensland.

Phosphorus soil testing

In many southern Queensland soils phosphorus levels may be in the low to moderate range. Moderate success using soil tests to recommend phosphorus application is evident for south Queensland.

Results from more than 200 phosphorus fertiliser trials on the Darling Downs revealed higher success in identifying phosphorus responsive soils by two soil tests:

- 80% of soils with bicarbonate phosphorus below 25 mg/kg and BSES acid phosphorus below 50 mg/kg responded to applied phosphorus.

Frequency of response of yield to phosphorus fertiliser

After 20 to 30 years of cropping, soil testing of most southern Queensland soils is recommended because of the inevitable depletion of soil phosphorus reserves.

However, both the bicarbonate extractable phosphorus test and/or the BSES acid extractable phosphorus tests may not necessarily identify soils that may demonstrate a yield response to applied phosphorus fertiliser. This is because crop responses at moderate to high nutrient supplies sometimes occur and vice versa. This occurs because of the highly variable rainfall. Thus in a similar way to nitrogen soil tests, phosphorus soil tests of nutrient supply are at best indicative of response frequency to applied nutrient rather than a categorical indication of a yield




Figure 4. Major inputs of nitrogen into the soil nitrogen mineral-nitrogen pool

response. The challenge for the grower is to establish when the response frequency or supply of nutrient from soil is adequate to sustain the expected level of crop production. Hence it is also important therefore to monitor phosphorus requirements of crops from time to time with on-farm experiments or seek computer simulation outputs of yield response frequency.

Sampling deeper soil layers for phosphorus

There is some evidence of relocation of phosphorus from sub-soil to the surface especially under zero-till practices. The surface soil test (0-10 cm) remains standard industry practice. However it is advisable to test the sub-surface level of phosphorus every 8 to 10 years to monitor changes in that region. The main outcome of an indication of low soil levels would be support for fertiliser application to crops at rates in line with the expectation of the response frequency.

When sowing chickpeas into the 10 to 15 cm soil layer it would be advisable to sample from this layer and apply phosphorus fertiliser if required.

Potassium

Potassium supply is potentially limiting in southern Queensland. Potassium is removed in substantial quantity by most cereal crops and may be in moderate to low supply.

The key points are:

- 0.4 meq/100g level, which is reported as being the critical level for wheat crops. Legume crops have critical levels as low as 0.25. Currently many current soil tests are returning levels in excess of this amount so potassium applications may not be warranted
- High clay content of most southern Queensland soils would suggest that potassium supply should not be quickly exhausted by continuous grain cropping.
- However, where cropping for silage production or where hay is removed, available potassium levels should be monitored at intervals of 3 to 5 years.

Table 5. Potential sources, frequency and magnitude of nitrogen gains in the soil

N gain or process	Source	Frequency	Magnitude	Comment
Fertiliser	Inorganic/organic	Regular	Gross	Residual in soil OM
Symbiotic N fixation	Legume plants	Regular	Gross	Growth dependent
Associative N fixation	Particular plant sp.	Regular	Slow rate	Indirect measurement
Other non-symbiotic fixation	Microbial available carbon (straw)	Irregular	Variable <1-20 kg/ha/yr	Dependent on C inputs
Electrical discharge	Lightning	Irregular	5-10 kg/ha/yr	Tropical storms
Plant ammonia absorption	Atmospheric ammonia	Irregular	usually <ammonia emissions	Net emissions usually greater

Source: Dr W Strong (pers. comm.)




Figure 5. Total N concentrations in relation to the period of cultivation.

Key: Soil and vegetation associations: 1. Waco (older alluvial black earth, grassland); 2. Langlands-Logie (clays, Brigalow); 3. Cecilvale (Poplar box); 4. Billa Billa (clays, Belah); 5. Thallon (clays, Coolabah); 6. Riverview (red-earth, Ironbark). Source: Dala and Mayer 1986.

Sulphur

Sulphur is usually released in soil and used by crops in the same way as nitrogen.

The key points are:

- Only about one tenth as much sulphur is required by crops as nitrogen
- In spite of this difference in quantitative requirement, it is anticipated that response to applied sulphur might occur when cereal response to applied nitrogen is frequent; in such soils sulphur may be required, particularly when switching to a leguminous crop
- In many southern Queensland soils, the existence of gypsum (CaSO_4) in subsoil layers will mean that a ready supply of sulphur will be available for crops even when sulphur supply from its release into topsoil is low. A surface soil test is of very limited value for assessing sulphur sufficiency.

Zinc

Zinc is used in small amounts by all crops but soil zinc supplies may also require supplementation in southern Queensland soils. Important points regarding zinc nutrition are:

- Low chemical availability of zinc in soils with high pH, e.g. vertisols
- Reduced colonisation of mycorrhiza-dependent plants after protracted fallow periods due to drought may combine to reduce plant-available zinc levels
- Crops not dependent on mycorrhiza are recommended to be sown after long fallow or failed crops
- Zinc-coated phosphate fertilisers provide a practical means of applying zinc to mycorrhiza-dependent crops like maize.




Figure 6. Fate of 75 kgN/ha nitrogen fertiliser (urea) labelled with ^{15}N applied to wheat at sowing over 4 seasons at Warra, in southern Queensland. Recovery in grain, straw, soil and that lost or not recovered are shown for each season; rainfall was for June to October.

Source: Dr W Strong (pers. comm.)

Soil testing procedures

It is vital that rigorous sampling procedures are adhered to and soil samples are handled appropriately. Soil testing companies supply such protocols. Another useful reference is 'Soil Matters' – monitoring soil water and nutrients in dryland farming (Dalgliesh and Foale 1998).

Some important issues are:

- Take sufficient samples for the paddock area in a randomised sampling pattern but avoiding atypical areas such as old fencelines, close to trees etc.
- Ensure that sampling equipment is clean
- Avoid contamination of the sample with other materials e.g. do not use galvanised buckets
- Avoid touching soil samples in hot weather (use a trowel)
- Sample to the correct depth
- Be aware that paddock history may be atypical e.g. after prolonged drought
- Collect the required quantity prescribed for laboratory testing that is bulked from the multiple cores from the paddock
- Be wary not to mix soil between the different depth increments
- Store samples in cool place (e.g. esky with ice) until they reach the lab
- Refrigerate samples if they cannot be sent immediately, air dry or dry at up to 40°C in an oven
- Record the position and time of collection on a map or with a GPS
- Despatch samples early in the week to avoid transit delays during weekends.

References

- Armstrong RD and Halpin NV. (1993). Potential residual value of nitrogen fertilisers in central Queensland. In: Proceedings 7th Australian Agronomy Conference, 108-110.
- Armstrong RD, Halpin NV, McCosker, K, Standley J and Lisle AT. (1996). Applying nitrogen to grain sorghum in central Queensland: residual value and effect of fallowing and tillage practice. *Australian Journal of Agricultural Research*. **47**, No.1, 81-95.
- Avalakki UK, Strong WM and Saffigna P G. (1995). Measurement of gaseous emissions from denitrification of applied Nitrogen-15. III Field measurements. *Australian Journal of Soil Research* **33** 101-111.
- Cornish PS, Ridge P, Hammer G, Butler D, Moll J and Macrow I. 1998. Wheat yield trends in the northern grains region – a sustainable industry. Proc. 9th Australian Agronomy Conference Wagga Wagga. pp. 649-652.
- Dalal RC (2005) Soil organic matter, farming organic C and the role of N. In: Proceedings of workshop, Long Term Fertiliser Use – better decisions for sustainability and profit, GRDC, Toowoomba.
- Dalal RC, and Mayer, RJ (1986). Long-term trends in fertility of soils under continuous cultivation and cereal cropping in southern Queensland. V rate of loss of total nitrogen from the soil profile and changes in carbon:nitrogen ratios. *Australian Journal of Soil Research* **24**, 493-504.
- Dalal RC. and Mayer R. (1990). Long-term trends in fertility of soils under continuous cultivation and cereal cropping in southern Queensland. VIII Available nitrogen indices and their relationships to crop yield and N uptake. *Australian Journal of Soil Research* **28**, 563-575
- Dalgliesh NP and Foale MA (1998). A guide to soil sampling. In: Soil Matters –monitoring soil water and nutrients in dryland farming, Eds. N Dalgliesh and M Foale, CSIRO, Australia.
- Harris PS, Biggs A J and Stone BJ. (eds) (1999). Central Darling Downs land Management Manual. Department of Natural Resources, Queensland DNRQ990102.
- Maher JM (ed) (1996). Understanding and Managing Soils in the Murilla, Tara and Chinchilla Shires. Department of Primary Industries Training Series QE96001, Brisbane
- Lawrence DN, Cawley ST and Hayman PT. (2000). Developing answers and learning in extension for dryland nitrogen management. *Australian Journal of Experimental Agriculture*, 2000, **40**, 527-539
- Millar G and Armstrong R (1999) How much have soil fertility levels declined in soils used for cropping in Central Queensland? Department of Natural Resources Queensland.
- Radford BJ, Silburn DM and Forster BA. (2008). Soil chloride and deep drainage responses to land clearing for cropping at seven sites in central Queensland. In preparation, *Journal of Hydrology*.
- Silburn DM, Cowie B A and Thornton CM. (2008). Brigalow catchments revisited – effects of land development on salt leaching and deep drainage determined from non-steady chloride profiles. In preparation, *Journal of Hydrology*.
- Spackman GB and Garside AL. (1995). In: Major factors affecting grain production in central Queensland and their implications for sustainable grain farming systems. A review prepared for the Grains Research and Development Corporation. Canberra, Australia.
- Strong WM, Dalal RC, Weston EJ, Cooper JE, Lehane KJ, and King AJ (1996). Nitrogen fertiliser residues for wheat cropping in sub-tropical Australia. *Australian Journal of Agricultural Research* **47**, 695-703.
- Strong WM., and Cooper JE. (1992). Application of anhydrous ammonia or urea during the fallow period for winter cereals on the Darling Downs, Queensland. I. Effect of time of application on soil mineral N at sowing. *Australian Journal of Soil Research* **30**, 695-709.
- Tolmie PE, Silburn DM, Biggs AJW 2003 'Estimating deep drainage in the Queensland Murray-Darling Basin using soil chloride.' Department of Natural Resources and Mines, Coorparoo, Queensland. QNRM03020.

4. Determining crop demand for nitrogen

Key messages

- Total crop water supply (rainfall plus stored soil water) determines grain yield. Grain yield determines the crop demand for nitrogen.
- Measure or estimate stored soil water at planting.
- Take soil water at planting into account when estimating target yield either ‘informally’ or with the *WhopperCropper* program.
- Use *WhopperCropper* to estimate yield ranges and select the midpoint of the season type of choice.
- Calculate crop nitrogen demand from a grain yield target and an optimal grain protein value; use the value for grain protein that will optimise grain yield with the available water supply (for wheat that grain protein value is 11.5%).
- Use the spreadsheet-based Nitrogen Fertiliser Calculator (Smart N Decisions) included with this manual.

Factors that affect crop grain yield

The main determinant of grain yield is water supply, namely:

- Stored soil water
- Rainfall received during crop growth.

Highly variable rainfall will result in highly variable quantities of stored soil water and in-crop rainfall. There are two important aspects for managing this variability:

- Measure or estimate stored soil water
- Use the probability concepts described in this manual to work with potentially variable yield outcomes
- The computer program, *WhopperCropper* directly provides the effect of soil water at planting and in-crop rainfall on yield.

Indicator of rainfall variability


For southern Queensland and northern New South Wales, the rainfall variability is rated from moderately to highly variable (Table 6 and Map 2).

- ANNUAL rainfall is rated as *moderately variable* (Map 2a)
- WINTER rainfall is *extremely variable* (Map 2b)
- SUMMER-growing periods have *high variability* in rainfall (Map 2c and d).

Table 6. Summary of rainfall variability ratings for south Queensland

Season	Rating scale #	Variability rating
Map a) – Annual mean	0.75 to 1.0	low to moderate
Map b) – July to September	> 2.0	moderate to very high
Map c) – November to January	1.0 to 1.25	moderate
Map d) – January to March	1.25 to 1.5	moderate to high

Variability = $(90p-10p) / 50p$ where 90p, 50p and 10p are annual 90th, 50th and 10th rainfall percentiles respectively.



Map 2. Regions of similar rainfall variability for periods, a) annual, b) July to September, c) November to January and d) January to March.

Source: Bureau of Meteorology <http://www.bom.gov.au/climate/averages/variability>

Factors affecting soil 'plant available water capacity' (PAWC)

The quantity of plant-available water that a soil can store depends upon its physical and chemical make-up.

a) Physical factors include:

- depth to bedrock materials
- texture (higher clay percentage stores more water)
- organic carbon percentage (hold water to a minor extent)
- structure (macro and micro pores hold water in clay soils but sandy soils allow considerable drainage)
- physical barriers such as compaction (reduced water entry)
- surface crusting (reduced water entry).

b) Chemical factors can also affect the 'effective' water-holding capacity by reducing root exploration or water uptake. The factors are:

- Salinity (presence of dissolved salts reduces water uptake, but not when gypsum is the cause of high electrical conductivity)
- sodicity (excess sodium ions – poor water infiltration, increased runoff).

Stored soil water as a factor affecting grain yield

The quantity of **stored soil water** has a significant effect on crop production, and will be the prime reason to adjust target yield. Thus the grower should have some knowledge of the expected stored water status of the paddock at planting.

Stored soil water at planting has considerable effect on crop yields and hence on crop nitrogen demand. This is especially apparent for *winter* crops. Thus it is important to:

- Maximise the storage of rainfall over the previous fallow
- Estimate or measure the stored soil water especially prior to winter crops by using either:
 - A push-probe to measure the depth of wet soil
 - Using the *Howwet?* computer program
 - The '20% rule'
 - Use *WhopperCopper* to indicate
 - The effect of stored water on grain yield
 - The effect of plant-available water-holding capacity (PAWC) has on potential yield.

Maximising the storage of rainfall in the previous fallow

Summer fallow efficiency (percentage of rainfall that is stored) is most commonly in the range of 18 to 22%. Very short fallows can have a fallow efficiency as high as 50%. The remainder is lost to evaporation, runoff or deep drainage. The key to maximising infiltration is to maintain a minimum of 30% soil stubble cover. Many growers have recognised this need and implemented zero or minimum tillage practices. This is even more effective when combined with fully matched two-centimetre accuracy controlled traffic farming systems.

Fallow efficiency decreases as fallow length increases so many growers have adopted ‘opportunity cropping’; crops are sown when stored soil water is adequate to sustain a dryland crop. This reduces loss of soil water to evaporation and dramatically reduces runoff and erosion when high intensity rainfall occurs when the soil profile is approaching fully recharged. Individual crop yields may be slightly reduced because of lower stored water reserves grain production is often higher in the longer term because of increased cropping frequency. Research is ongoing to model optimum crop frequency for Queensland farming systems.

Critical components of stored soil water

Not all of the water in the soil is available for use by a crop. The crop can only reduce water content to the level of ‘crop lower limit’ (CLL) or ‘wilting point’. The maximum amount of water a soil can hold is the ‘drained upper limit’ (DUL). The difference between these two values is the ‘plant available water capacity’ (PAWC). As soil water is recharged, the quantity of water stored is the ‘plant available water (PAW) (Figure 7).

As plant roots grow, soil water will be extracted progressively from soil until the soil reaches the CLL (approximately 50% of the total water in clay soil may be extracted by plants). This ‘drying-front’ progresses downwards as the roots grow into the soil.

The surface soil can dry to less than the crop lower limit due to the evaporation effects of sun and wind. If this occurs, this deficit will need to be overcome before water is available for crop use. During rainfall, after the cracks in the soil are closed, water recharge occurs from the top down.

Methods to estimate the amount of soil water at planting

All the methods have inaccuracies, but all should enable estimation of soil water especially to the broad categories required for *WhopperCropper* (one-third, two thirds and full profile).

Depth of wet soil

The amount of water held in the soil can be roughly determined by the depth of wet soil using a ‘push probe’. Table 7 describes typical rooting depths and plant available water contents.




Figure 7. Representation of critical components of soil water. Likely location of plant-available water in a typical vertosol of 150 mm water-holding capacity (PAWC) when a) one third recharged (=50 mm plant-available water PAW), b) two thirds recharged (=100 mm PAW) and c) fully recharged (= 150 mm PAW).

Use the Howwet? software program

Howwet? allows the user to enter actual rainfall data for a paddock of interest. Using selectable factors such as:

- soil type
- stubble cover
- paddock slope
- fallow start and end date.

The program estimates the proportion and total amount of the rainfall that should be stored in the soil.

Figure 8 illustrates simulated accumulation of soil water between 1 October 2005 and 15 April 2006 at Dalby, south Queensland. It shows that favourable early fallow rainfall (October and November 2005) quickly recharged the soil profile. Rainfall in January and February 2006 probably moved water to deeper soil layers, adding to that portion of water storage that is protected from subsequent evaporation. Efficiency with which water can be stored early in the fallow can be quite high (up to 40 to 50%) because of structural cracks in dry soil and the short time over which evaporation can occur. The downward slope of the line indicates loss of soil water through evaporation. The red columns illustrate water runoff.

Howwet? also provides a table of outputs (e.g. Figure 9). Some example details are:

- 441 mm (75%) of the 588 mm of rain received during this period evaporated
- The gain in soil water was 120 mm from a total rainfall of 588 mm (about 20% of fallow rainfall stored)
- The overall fallow efficiency of 20% is fairly typical for fallow periods at this time of year in south Queensland
- The estimate for soil water stored at the end of the fallow, 137 mm (83% of a fully wet profile) could be described as an 'excellent' prospect for subsequent cropping in spite of the high overall quantity of rainfall that was evaporated.
- 31% of rainfall occurred in events of less than 15 mm that quickly evaporated (high percentage of small rainfall events to reduce efficiency with which rainfall is stored).

Table 7. Soils and typical plant available water contents in Queensland

Soil type	Rooting depth (m)	Available soil water (mm/m)	Typical PAWC
Heavy alluvial, flooded brigalow or yellowwood	1.25	170	212
Friable alluvial	1.25	185	230
Shallow open downs	0.75	180	135
Deep open downs	1.0	160	160
Light alluvial	1.2	185	222
Heavy mixed alluvial	1.1	180	198
Deep/heavy brigalow or brigalow-belah clay	1.2	160	192
Scrub walloon soils	1.05	180	189
Brigalow/softwood scrub	1.0	160	160
Light Callide alluvial	1.2	150	180
Shallow/light brigalow or box clay	0.9	160	144
Black upland soils	1.05	180	189
Light box clay	0.85	160	119
Softwood brigalow	0.85	140	119
Brigalow/Dawson gum brigalow duplex	0.95	150	143
Forest walloon soils	1.0	120	120
Red brown/red earths	1.05	125	131
Red upland basalt	1.0	100	100

Source: from Wheatman v4



Figure 8. Graphical representation of soil water accumulation at Dalby from rainfall events during the fallow period, 1 October 2005 to 15 April 2006, as simulated by Howwet?



Figure 9. Soil water storage simulated for the summer fallow, 1 October 2005 to 15 April 2006, using Howwet?.

The '20% rule'

Most grain growers have access to rainfall records, so when all other methods to estimate fallow water storage are unavailable, a rough rule of thumb could be derived from fallow rainfall using a figure of 20% of the rainfall being stored.

The '20% rule assumes that the single fallow efficiency value of 20% accounts for the average losses through runoff and evaporation that occurs following fallow rainfall. It is a reasonable first approximation of the amount of rain that may be accumulated over a seasonal fallow (e.g. from winter crop harvest to the next winter crop planting). A weakness in this approach is that fallow efficiency usually varies with the length of the fallow and patterns of rainfall. Typically, actual values are higher at the start of a fallow (around 40%) when the soil is dry and cumulative evaporation is low. As the fallow period is extended cumulative evaporation (and sometimes runoff) can increase. Thus in a double crop situation the fallow efficiency value can be very high. Average fallow efficiency values of between 18 and 25% (average 23%) are typical for southern Queensland for a full summer fallow (winter crop to winter crop).

Note: In a test of the same set of fallow data (Figure 10) it was found that improved accuracy was gained if values of 18% were used for western Queensland and 23% for south eastern regions. When this was done, accuracy of the result was similar to *Howwet?*


Using WhopperCrapper to indicate the effect of soil water-holding capacity and soil water at planting

Because of rainfall variability, a field nutrition experiment over one or two years will not be a good indicator of the possible yield outcomes. For this reason, crop simulation modelling is used to demonstrate the full range of potential yields. This enables the user to act upon their 'attitude to risk' when choosing input levels.


When the soil water at planting has been calculated by one of the methods previously described, the potential yield ranges can be generated by the *WhopperCrapper* program.

Because the data is generated from 100 years of rainfall data, the graphs are 'probabilistic' in nature. This has the advantage of providing a realistic range of potential yields to 'target' rather than a single yield as used in previous nitrogen budgeting techniques.

Figure 11 indicates the large effect of soil water at planting on wheat yield at Dalby. The median yield (heavy dark line) increases from 1200 kg/ha to almost 3500 kg/ha as the soil water at planting was increased from one third full to completely full (in every year of the simulation). Because the soil water is reset to the indicated value **every** year, the output indicates the broad insights rather than a yield for any particular year.



a) Modified '20%' rule. (Note: 18% fallow efficiency used for central and western Queensland and 23% fallow efficiency for south Queensland) $R^2 = 0.75$



b) Howwet? $R^2 = 0.74$

Figure 10. Observed vs predicted plant available water in the root zone at the end of fallows

Source D Freebairn (pers comm.)




Figure 11. Seasonal distribution of wheat yields at Dalby with stored water at planting equal to one third, two thirds or full at planting. Soil N assumed to be 100 kgN/ha + 750 kg/ha fertiliser N applied at planting. Planting date of 15 May, 1 million plants/ha, medium maturity variety, soil PAWC=190 mm.

How to read this *WhopperCropper* boxplot

The black line in the 'red box' is the median yield (50% of all years have this 'yield' or less). The dashed line within the 'red box' is the mean yield. The upper edge of the 'red box' is 75% probability. This is read as 'in 75% of years, yields will be less than this value'. The lower edge is the 25% probability value i.e. 'in 25% of years, yields will be less than this value'. The upper and lower short horizontal lines represent 100% and 0% probabilities respectively.

In contrast, the effect of soil water at planting is slightly less evident on sorghum yield because of the more dominant summer rainfall (Figure 12). However the riskiness of the yield outcome is significantly increased if planting was conducted with one-third full profile every year as indicated by to absolute range of yields from 0 kg/ha to 7500 kg/ha depending upon the seasonal rainfall.

Probability concepts

Whilst it is widely recognised that yield variability occurs, understanding and working with yield ranges may assist input planning and risk management compared with using a 'commonly-accepted' farm, district or state average.

Probability is derived from the concept that a 'theory' (hypothesis) is tested more than once to ensure the result has not occurred simply by chance and/or is an 'unusual' result. In terms of crop yields, we know that many factors can affect yield in a particular year. Some of the factors are controllable e.g. nitrogen fertiliser rate but many are not e.g. rainfall and the timing of that rainfall. Some factors also interact e.g. yield response to nitrogen fertiliser rate with rainfall. Several factors can interact in ways that increase or decrease yield.

Generating 'target yield' ranges

In order to calculate crop nitrogen demand, a **target yield** range for the forthcoming season must be determined. This can be achieved from:

- Farmer experience (paddock history)
- Crop simulation models (as used in *WhopperCropper*).

Farmer experience

Farmer experience and knowledge of paddock performance over many seasons can indicate an achievable level of crop production. Farmer field trials can also test principles but it must be remembered that they only represent a small percentage of the possible outcomes. Crop models are developed from accurately conducted research trials and then tested in other trials.

WhopperCropper

The APSIM Crop simulation model (the building block for *WhopperCropper*) can be used to demonstrate the distribution of yields achieved over the full length of rainfall records from a site. After they are generated, the simulated outcomes provide




Figure 12. Seasonal distribution of sorghum yields at Dalby with stored water at planting equal to one third, two thirds or full at planting. Soil N assumed to be 100 kgN/ha + 75kg/ha fertiliser N applied at planting. Planting date of 15 October, 60 000 plants/ha, medium maturity variety, soil PAWC=190mm, solid 1m rows.

a yield distribution from which a target yield might be selected. Selecting a target yield from the full possible range is a more informed decision because of the inherent seasonal variability but also because it embraces the grower's attitude to risk.

WhopperCropper is an easy-to-use package that enables scenarios to be created to generate yield ranges with appropriate practical setup parameters. The full range of input parameters in *WhopperCropper* are:

- Crop type – common summer and winter
- Soil water-holding capacity – up to five levels
- Soil water at planting – one third, two thirds and full
- Planting date – up to five dates
- Maturity length – three categories
- Plant population – usually three levels
- Row configuration – skip in sorghum and cotton

- Effect of soil nitrogen content – usually three levels
- Nitrogen fertiliser rate (planting and in-crop) usually six nitrogen rates – usually six nitrogen rates.

Simulating long-term yield ranges

The first step in generating yield probability graphs is to generate annual crop yields from a desired scenario or scenarios. This graph type is an output from *WhopperCropper* (Figure 13).

To generate a *probability distribution*, the simulated yield data is arranged from lowest to highest and then subdivided into (for example four) categories of equal number of seasons representing low, low-moderate, moderate-high and high yields.

Each of these categories is represented for simulated wheat yields at Dalby and shown as a box plot in Figure 14. When expressed in terms of probabilities the yield ranges are as indicated.

The sections of the boxplot are effectively subdividing the yield range into sub-groups (Figure 15).

These are read as (for example):

- In '75% of years, yields will be less than 2939 kg/ha'
- The lower edge is the 25% probability value i.e. 'in 25% of years, yields will be less than 1074 kg/ha'
- The upper and lower short horizontal lines represent 100% and 0% probabilities respectively (5665 and 608 kg/ha respectively).

These outputs provide reference for the grower to judge the likelihood of the expected yield target, as well as enabling adjustment of the expected yield target for potentially different water supplies to the crop.

This approach allows yield and financial risks to be made quite evident.




Figure 13. Wheat grain yields (600 kg/ha to 5500 kg/ha) were simulated for the period 1890 to 2005 at Dalby, each with monoculture cropping with two thirds of a full profile stored soil water




Figure 14. Diagram of how 'boxplot' yield results relate to four season type ranges




Figure 15. Diagram of how individual yield results are grouped as a boxplot with the lowest and highest yields being represented as 0 to 100th percentile of the range respectively

Choosing a single target yield

The components for calculating crop nitrogen demand are simply:

- target crop yield
- grain protein content.

The basis for the nitrogen demand calculation is that a single target yield be selected. Selecting a high target yield might expose a grower to unspecified financial risk because of a high frequency of poor to moderate crop yields. Provision of the likely yield range for long-term seasonal outcomes enables the target to be selected with full knowledge of the associated risk or likelihood of achieving that yield. Selecting from the full potential range of yield outcomes is a more informed way to select a target yield because of the high inherent seasonal variability in regions of northern Australia.

Nitrogen fertiliser needs to be managed in the context of districts rainfall variability. Maximising profits in the good seasons whilst reducing losses in the more frequent moderate to poor seasons is the key to optimising profits in this environment. A suggested process is as follows:

- Use *WhopperCropper* to provide the long-term range of yields that occur for a district
- Use the default *WhopperCropper* 'boxplot' to demonstrate the four categories of equal number of seasons that may occur (low, low-moderate, moderate-high and high yields)
- Select a target yield from a category knowing the likelihood with which that yield might be achieved; 25% high to very high yields, 25% moderate-high yield, 25% low-moderate yields and >25% zero to low yields
- Calculate nitrogen demand for the target yield using the formulae below.

Selection of target yield knowing that yield can vary considerably enables a grower to incorporate their 'attitude to risk' in the decision. Anticipation of the forthcoming season or a 'desired-bet' attitude can be reflected in the choice of target.

The mid-point of the yield range for each category could be a reasonable choice for the single 'target yield' value. Naturally the final choice of target yield may be influenced by financial constraints but knowledge of the likely long-term outcomes will enable the grower to avoid unnecessary financial risks.

These outputs provide reference for the grower to judge the likelihood of the expected yield target, as well as enabling adjustment of the yield target for expected water supply for the crop and for an acceptable financial risk.

This is probably the most informed approach of all to evaluate fertiliser needs because financial risk is made evident in model outputs.

Calculating crop nitrogen demand from target yield

To estimate nitrogen demand for the target grain yield, a final grain protein is assumed that reflects a crop in which the nitrogen supply has not restricted grain yield. Yields of cereal crops will not have been limited by nitrogen supply when grain protein concentrations are more than:

- 11.5% for wheat
- 9.5% for sorghum
- 10.5% for barley.

Source: Dr W Strong

These proteins refer to grain moisture levels of 12%, 13% and 0% respectively.

These are considered the optimum grain protein levels to target and indicate the adequacy of the nitrogen supply for the seasonal outcome. Crops with resultant grain proteins below these critical contents may have been adversely affected by inadequate nitrogen supply. Crops with proteins above these concentrations can usually access an adequate supply of nitrogen to respond to the water available, including water stored at planting and rainfall during crop growth. At these grain protein levels there is confidence that the nitrogen supply was adequate for the optimum economic production.

Equation to calculate nitrogen demand

Crop nitrogen demand can be readily estimated by calculating grain nitrogen (kg/ha) and converting this to crop demand using a simple multiplier that has been derived from many field trials with wheat and sorghum.

The multiplier to obtain total amount of nitrogen that is required for wheat and sorghum (grain + vegetative matter) at the optimum grain protein is 1.7.

Note: Constants in the equations below describe the fraction of nitrogen in wheat protein (10/5.7), conversion of grain yield t/ha to kg/ha (*1000), nitrogen concentration as a percentage (1/100) and the multiplier to convert grain nitrogen to crop nitrogen demand (1.7). Calculations for sorghum nitrogen demand are identical to those for wheat except the fraction of nitrogen in sorghum protein is (10/6.25).

The 1.7 conversion factor, and the given grain protein percentages, are those considered to give the most economic use of nitrogen.

Table 8. Crop nitrogen demands (kg/ha) estimated using the above formula for a range of expected wheat and sorghum yields (t/ha) and targeting 11.5 and 9.5% protein respectively.

Target grain yield	Wheat N demand	Sorghum N demand
1.0	34	26
1.5	51	39
2.0	69	52
2.5	86	65
3.0	103	78
3.5	120	90
4.0	137	103
4.5	154	110
5.0	171	129
6.0	206	142
7.0	240	155
8.0	274	181

Note: remember that these values represent the total crop demand not the fertiliser application required. The soil nitrogen supply needs to be subtracted from this value to obtain the fertiliser required (see Chapter 5 and 6).

Table 9. Efficiency with which plant-available nitrogen in soil is transferred to grain of wheat, barley or sorghum. The shaded area represents the most economic target level.

Protein (%)	Wheat	Barley	Sorghum
9	1.35	1.32	1.58
10	1.49	1.43	1.94
11	1.64	1.56	2.50
12	1.81	1.74	

Source: Dr W Strong (pers. comm.)

Hence nitrogen demand for a **wheat** crop, with a target yield of 2.5 t/ha can be calculated as follows;

Equation 1.
 Nitrogen demand (kg/ha)
 $= (\text{Grain yield (t/ha)} * \text{Grain protein percent} * 10/5.7) * 1.7$
 $= (2.5 * 11.5 * 10/5.7) * 1.7$
 $= 85.7 \text{ kg/ha}$

Similarly, nitrogen demand for a **sorghum** crop, with a target yield of 2.5 t/ha can be calculated as follows;

Equation 2.
 Nitrogen demand (kg/ha)
 $= (\text{Grain yield (t/ha)} * \text{Grain protein\%} * 10/6.25) * 1.7$
 $= (2.5 * 9.5 * 10/6.25) * 1.7$
 $= 64.6 \text{ kg/ha}$

Table of crop nitrogen demand as determined by grain yield

Estimated crop nitrogen demand for wheat and sorghum at various yield expectations have been calculated using equations 1 and 2 and are shown in Table 8. It can be used to directly determine nitrogen demand at 9.5% and 11.5% protein for wheat and sorghum respectively.

Table 8 was generated using the same nitrogen transfer efficiency factors of 1.7 which corresponds to grain protein levels of 11.5 and 9.5 for wheat and sorghum respectively. If a different grain protein is to be targeted a different transfer factor should be used (Table 9). Notice that for each crop the 'efficiency factor' varies with final grain protein content. Shaded areas denote grain nitrogen factors at protein ranges of most economical production.

If different grain protein targets are required, substitute values from Table 9 into equations 1 or 2.

Using the SOI phase system to modify the target yield

To this point we have considered that simulated outputs be categorised only by yield, with equal number of seasonal outcomes assigned to each yield category. However, if a reliable seasonal forecasting tool was available, different yield distributions may be apparent for different seasonal forecasts. The SOI phase may provide some skill to forecast the coming season. Figure 16 shows the shift in wheat yield distribution for contrasting phases of SOI at Dalby, south Queensland.

Note the modelled planting date was 30 May with an April/May SOI phase used. This was because at earlier planting dates, the SOI phase has no skill to modify yield probabilities. The 'locking-in' of the SOI phase is often after the desired winter crop planting date and thus of no value to a winter-planting decision. However, late winter/early spring rain can also be impacted by SOI which may influence a spring or summer crop planting and nitrogen fertiliser decisions.




Figure 16. Yield ranges for wheat at Dalby 15 June planting for Positive and Negative SOI phases and stored water two-thirds full. The positive and negative SOI distributions are significantly different at the 5% level using the Kruskal-Wallis test

Source: WhopperCropper

Case study

Effect of sorghum row spacing on yield


Results from modelling and field experiments have indicated that skip row configurations can result in increased yields compared with solid plant configurations at yield levels below about 2.6 t/ha (Routley *et al.* 2003). This is presumed to be a result of conservation of soil water in the centre of the skip area for use by the plant in the grain filling stage. At higher potential yield levels, a yield reduction can occur with skip row configurations (Figure 17). Hence the choice of row configuration for a particular paddock situation will depend on available soil moisture at planting, likely in crop rainfall and the producer's attitude to risk.

Additional factors that should be taken into account are:

- Possible reduced rainfall infiltration because of low stubble cover in the inter-row space;
- Weed control difficulties because of lack of crop competition in the inter-row space.



A wide row sorghum configuration demonstrating low cover in the inter-row area which may impact on rainfall infiltration in this zone



Data from these experiments

Figure 17. Relationship between solid plant yield and skip yield (Source Routley *et al* 2003, adapted from Butler *et al* 2001)

References

Butler G, Cawthray S, Castor M, Yeates S and Christian T. 2001. Improving the reliability of sorghum production in the farming system. In: Proceedings 10th Australian Agronomy Conference, 2001. Australian Society of Agronomy.

Routley R, Broad I, McLean G, Whish J. and Hammer G. (2003). The effect of row configuration on yield reliability in grain sorghum: 1. Yield, water use efficiency and soil water extraction. In: Proceedings of the 11th Australian Agronomy Conference, 2003, Australian Society of Agronomy.

5. The soil's capacity to supply nitrogen

Key messages

- Deep soil testing for nitrate-nitrogen is the best method available but still prone to inaccuracy.
- Refer to the soil sampling rate table to determine a sampling rate for accuracy:
 - For example to be correct 8 times in 10 with an accuracy of $\pm 10\%$ would require 18 cores per 40 ha
 - Avoid obvious non-representative areas like old fence lines and different soil types.
- Soil organic carbon will generally reflect the period the paddock has been growing crops and can give an approximation of the potential to supply nitrogen:
 - Use organic carbon or the yield and protein of previous crops to gauge ability to supply nitrogen.

Calculation – Soil nitrogen supply (from soil test)

For each depth layer, plant-available soil nitrogen is calculated from the quantity of nitrate-nitrogen measured in that layer.

Available nitrogen = soil test value (mg/kg)
 * soil bulk density (g/cm³)
 * number of 10 cm increments.

For example, depth layer 1 (e.g. 0 to 60 cm)

e.g. Available nitrogen = 8
 * 1.1 (average bulk density over the layer)
 * 6 (six ten cm layers)
 = 52.8 kgN/ha

Smart N decisions – a nitrogen fertiliser calculator

Supplied with this manual is an easy-to-use computerised calculator that has a section for calculation of crop nitrogen demand. On a single page, the relevant data are entered for:

- Soil sampling date
- Expected planting time
- Stored soil water
- Expected season type (e.g. poor, average etc)
- Soil test results.

The calculator estimates:

- Nitrogen mineralisation from soil sampling to planting
- Crop demand
- Soil supply
- Nitrogen fertiliser rate required.

By using the Seasonal Outlook and Desired Yield graph ('Expected Season Type'), in conjunction with the 'Seasonal Comparison' table (where gross margins are calculated), the optimum nitrogen fertiliser rates for each season type can be calculated.

In addition, an estimate of the losses incurred if the season turns out differently to that targeted or anticipated can be calculated.



Figure 18. The single page on which the calculations are made

Direct measurement of plant-available nitrate-nitrogen in the soil

Direct measurement of the pool of nitrate-nitrogen to rooting depth, usually by soil layer and close to sowing, provides the most accurate measure of plant-available nitrogen supply in soil.

Applying nitrogen fertiliser without measuring soil nitrogen supply may lead to oversupply of nitrogen due to build up of plant-available soil nitrogen after extended fallow periods as a result of drought. Measurement of soil nitrogen supply may avoid unnecessary fertiliser application.

Table 10. Relationship between the number of soil cores taken on areas up to 40 ha, the accuracy of the results and the confidence that the mean value will fall within the level of accuracy.

Confidence level #	Number of cores required	
	Nitrate	Water
Med level of accuracy \pm 20% of mean		
66%	3	2
80%	5	3
90%	8	5
High level of accuracy \pm 10% of mean		
66%	10	7
80%	18	12
90%	29	20

the confidence level indicates how often the result would be within the level of accuracy i.e. 80% = correct 8 in 10 times.

Source: Soil Matters – monitoring soil water and nutrients in dryland farming, Eds. N Dalgliesh and M Foale, CSIRO, Australia (derived from Jones 1994).

Principles of soil sampling

Unfortunately, soil testing is time consuming and prone to error due to inherent variability in soil nitrate-N in most soils. This is especially so in brigalow scrub soils, in which the original vegetation may have been patchy, and due to a generally shorter duration of cropping.

This contrasts with more uniform open grassland soils that were originally predominantly grass vegetation and have been farmed for longer, resulting in mixing of the organic matter and depletion of soil nitrogen to uniformly low levels.

The only option is to take as many soil samples as is practically possible. Samples should be kept cool (4°C) and sent to the laboratory as soon as possible. Table 10 demonstrates the trade-off between accuracy and the number of samples that are bulked to make a test sample. These values were formulated for soils of south Queensland for areas up to 40 ha. Soils of higher variability such as in central Queensland may require higher sampling intensity than listed here. Sampling for soil water requires fewer cores because it is slightly less variable.

Sampling patterns

Typical soil sampling patterns for fallow paddocks are diagonal, circular or random positions (Dalgliesh and Foale 1998). None of these techniques make it easy to identify where the samples originally came from. GPS locating of sample sites may be useful if there is a need to return to the same spot for repeated sampling as may be required where soil properties such as organic carbon are to be monitored.

Depth of sampling

For nitrate sampling to determine plant-available nitrogen, coring to the depth of the wet soil is advisable in order not to over-estimate the potential supply. An estimation of the rooting depth of the crop is required when soil sampling is well ahead of planting. The supply of plant-available nitrogen for the next crop will most likely be derived from the depth of soil water recharge. Table 11 details the current accepted depths of sampling.

Table 11. Suggested sampling depths (cm) for chemical and water analysis for deep soils without sub-soil constraints

Crop	Rooting depth	Water, chloride, E.C and pH	Nitrogen ¹	Phosphorus, zinc, organic carbon
Sorghum, cotton	180	180	90	10
Wheat, chickpea ²	150	150	90	10
Mungbean	120	120		10

1. Sample to full rooting depth if a nitrogen bulge is suspected.

2. If 'deep' planting, sample to expected planting depth.

Sample depth increments

The number of increments is a compromise between the cost of sampling and analysis and the benefits gained from knowing the position of water, nitrogen and other nutrients in the profile. Too few increments may miss a dry layer thus overestimating the effective amount of water available. Too few or too shallow sampling may also miss a nitrogen bulge that may be able to supply a crop in a good season or accurately identify the position and concentration of chemical subsoil constraints to root growth and water extraction in the profile. However, a judgement must be made as to whether nitrogen in deep soil layers would be available in the majority of seasons.

Limitations of soil nitrate-nitrogen testing

Soil nitrate-nitrogen tests should only be used as a guide to indicate potential nutrient supply. Predicting supplementary nutrient needs with any degree of accuracy relies on capability to predict **total crop requirements** at the time fertiliser is applied, usually before sowing.

Possible errors in using soil testing to predict crop nitrogen demand include:

- Variability of distribution of plant-available N creates difficulty in collecting representative soil samples
- Dry conditions in the previous fallow may underestimate soil potential to supply nitrogen for the subsequent crop
- In dry seasons, fertiliser nitrogen may be stranded in dry top soil
- In wet seasons, applied or native nitrogen in topsoil may be lost to the atmosphere.

Nevertheless, frequency of response to applied nutrient over a sequence of crops is still a useful parameter that can be derived from most soil tests in regions of unreliable cropping. As this is expensive and time consuming the easiest way is use simulation modelling tools such as *WhopperCropper*.

Using soil nitrate as an indicator of the potential of a soil to respond to nitrogen fertiliser

Season-to-season variation in crop production is usually related to variable water supplies (stored soil water and in-crop rainfall), and may create huge variation in crop nutrient demand. When used against a background of such extreme variation in season-to-season nutrient demand, soil tests can never be expected to separate responsive from non-responsive seasonal outcomes. If the soil inherently contains a considerable quantity of nitrogen, additional nitrogen fertiliser will only increase grain yield in years of highest rainfall. An example of this is shown for 11 wheat crops at *Warra* on the Western Downs where the frequency of response to applied nitrogen decreased with increasing soil test (plant

available nitrogen) from 90% at a low total available soil nitrogen (50 kg/ha) to 36% at 150 kg/ha total plant available soil nitrogen (Figure 19).

The challenge for applying soil tests wisely to cropping in unreliable regions is to establish a soil test and frequency of response to fertiliser that is profitable while maintaining an appropriate level of soil nutrient for future crop production.

For nitrogen, where application costs per crop are high:

- applying nutrient only when there is high response frequency (low current nitrogen levels and high soil water availability) may be profitable but may not sustain cropping in the long term
- applying nutrient when the expected response frequency is low may be less profitable over the short term but may improve sustainability in the longer term
- when a soil test indicates a low soil level of nutrients like P and Zn, fertiliser application may be advisable even though the response frequency may be fairly low. This is because the application cost per crop is moderate but large yield gains can be obtained in some years. More importantly, non-limiting supply of the other nutrients will facilitate a more reliable yield response to applied nitrogen.

Soil organic carbon as an indicator of response to nitrogen fertiliser

Soil organic carbon is frequently used as a surrogate measure of soil fertility status because it is a proportional measure of the amount of organic matter in the soil (58% of soil organic matter is carbon). It is a relatively inexpensive laboratory test. However, this has limitations as indicated in both Tables 12 and 13. For soils of the Darling Downs with organic carbon content below 1.0%, response frequency did not exceed 70 percent (Table 12). At organic carbon levels greater than 1.0%, frequency of response was only 50% or less. Total N percent is also a measure of soil fertility but is a much more expensive test.

The management practices on the Darling Downs during the 1960s and 1970s when these trials were conducted, utilised a high proportion of long fallows (14 to 16 month) and mechanical tillage. High levels of nitrate-nitrogen in long fallows would have masked potential response to nitrogen fertiliser due to longer duration of soil nitrogen mineralisation. However, within shorter fallows, soil organic carbon can be a useful indicator of potential crop response frequency to applied nitrogen.

Organic carbon and total nitrogen levels are indicative or surrogates of the soil's capacity to supply nitrogen and are therefore useful to monitor trend in soil nitrogen supply or frequency of crop response within various crop rotations.

Soil nitrate level, on the other hand is a consequence of these soil properties (organic carbon and total nitrogen) as well as duration and conditions during the fallow when organic




Figure 19. Demand for nitrogen (kg/ha) for 11 rain-fed wheat crops at Warra indicated by vertical bars. Horizontal lines indicate 3 soil test levels of available nitrogen, 50, 100 and 150 kg/ha. This sequence of wheat crops would have responded to fertiliser nitrogen with a frequency of 90%, 63%, and 36% respectively, where 50, 100 or 150 kg/ha of nitrate-nitrogen was measured at sowing. Notice the imprecise nature of predicting crop response from soil nitrate-nitrogen with such unreliable cropping.

Table 12. Response frequency to applied nitrogen for Darling Downs soils cropped with wheat in 1965-1971, categorised by soil organic C level or total soil nitrogen level. Neither soil test totally separates responsive from non-responsive sites but each is useful to monitor trend in crop response frequency within various crop rotations.

Organic carbon percent	<1.0	1.0-1.4	>1.4
Response frequency (per cent)	70	50	30
Total N percent	<0.1	>0.1	
Response frequency (per cent)	60	30	

Table 13. Simulated monthly release of plant-available nitrogen (kg/ha) for soils with a range of organic carbon levels at Dalby on a soil of 240mm PAWC that is 67% full prior to planting and contains 50kgNO₃/ha. Planting date is 15 May for wheat and 15 October for sorghum. The shaded area indicates mineralisation during a typical cropping period of either wheat or sorghum crops.

Organic carbon	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
With wheat crop												
0.8	9.4	8.7	9.1	7.2	0.9	1.7	1.9	2.3	2.7	2.5	5.4	8.2
0.9	12.4	11.5	11.8	9.2	0.7	2.7	2.9	3.4	3.8	3.7	7.0	10.9
1.0	13.5	12.5	12.8	9.9	1.3	3.2	3.2	3.8	4.0	4.2	7.7	11.7
1.1	14.4	13.2	13.7	10.6	1.9	3.6	3.6	4.1	4.6	4.7	8.7	12.7
1.2	15.3	14.0	14.5	11.2	2.5	4.0	4.0	4.4	4.9	5.0	9.4	13.5
1.3	16.4	15.0	15.5	12.1	3.1	4.4	4.4	4.8	5.2	5.4	10.0	14.5
1.4	17.5	16.0	16.4	12.8	3.8	4.9	4.7	5.2	5.5	5.8	10.0	14.5
1.5	18.3	16.8	17.3	13.4	4.3	5.2	5.0	5.5	5.7	6.1	11.3	16.2
1.6	19.0	17.4	17.9	13.9	4.7	5.5	5.3	5.7	5.9	6.5	11.7	16.8
With sorghum crop												
0.8	9.1	2.4	4.8	5.7	4.9	3.8	3.7	4.6	6.5	1.9	7.0	8.9
0.9	10.0	2.7	5.4	6.6	5.6	4.3	4.2	5.1	7.2	2.7	8.0	9.9
1.0	10.8	3.0	6.4	7.4	6.2	4.7	4.5	5.5	7.8	3.5	8.7	10.8
1.1	11.5	3.2	7.2	8.1	6.8	5.1	4.9	5.9	8.3	4.3	9.8	11.7
1.2	12.1	3.5	7.9	8.8	7.3	5.5	5.2	6.3	8.8	5.0	10.7	12.6
1.3	12.9	3.8	8.7	9.6	7.8	5.8	5.5	6.8	9.4	5.9	11.7	13.6
1.4	13.5	4.1	9.5	10.3	8.3	6.2	5.9	7.2	10.0	6.8	12.7	14.5
1.5	14.1	4.4	10.1	10.9	8.7	6.5	6.2	7.5	10.4	7.5	13.5	15.3
1.6	14.5	4.7	10.7	11.4	9.1	6.8	6.4	7.8	10.8	8.3	14.4	16.0

Source: APSIM version 6.1, Keating et al (2003).

Table 14. Selectable inputs and parameter choices for WhopperCropper ‘fallow’

Input factor	Options
Soil organic carbon levels	0.8, 0.9, 1.0, 1.1, 1.2, 1.4 (per cent)
Districts	Dalby, Goondiwindi etc.
Soil PAWC's	80, 120, 150, 190 mm
Date of start of fallow	15th of every month
Fallow length	2, 5, 7, 12 months
Soil water at start of fallow	0, 25% and 50% full
Soil nitrogen at start of fallow	0, 25, 100 kgN/ha




Figure 20. Range of simulated soil nitrates (kgN/ha) at end of 2, 7 or 12 month fallows commencing 15 October, Dalby, 150 mm PAWC, 25% full, 25 kgN/ha at commencement of fallow. Simulations involved 100 years of weather data. The solid line is the median value, dashed line is the mean. Other points are 100%, 75%, 25% and 0% values respectively.

Source: WhopperCropper.




Figure 21. Example of fallow nitrogen mineralisation estimate from Howwet? The setup is partially described above and uses Dalby weather data, soil organic carbon of 1.0% (assumed 20 years of farming), 30% cover and 10% profile refill at fallow start, 20% cover at fallow end, slope = 1%.

nitrogen is converted into the mineral form, principally nitrate-nitrogen. Hence level of nitrate-nitrogen in soil is dynamic and timing of its measurement is very important if it is used to indicate plant-available nitrogen supply.

Simulation tools use for estimating nitrogen mineralisation during a fallow

The practicalities of soil testing require a time lag between sampling and planting to enable laboratory analysis. Hence, an estimate of the amount of nitrogen that might be mineralised between sampling and planting is required. Thus when soil sampling is much earlier than planting time, plant-available nitrogen determined by an early soil test can be adjusted to estimate the level at planting. An estimate of the additional nitrogen that is mineralised in that period can be obtained from a number of simulation tools.

Using a ‘look-up’ table generated using the APSIM model

Estimated **monthly** nitrate-nitrogen releases for soils of differing organic carbon levels at Dalby, south Queensland derived from APSIM are shown in Table 13. These outputs were derived using the following set-up options:

- Soil organic carbon range 0.8 to 1.6%
- 170 mm soil PAWC
- Soil water approximately two thirds full every year
- 15 May sow date (wheat), 15 December sow date (sorghum)
- 50 kgN/ha in the soil at sowing.

WhopperCropper ‘fallow’

Using the ‘fallow’ option (under the crops selection menu), WhopperCropper provides the capability to estimate the following outputs

- soil nitrogen mineralisation (and soil nitrate at end of fallow)
- storage of water (and stored soil water at end of fallow)
- runoff (total)
- drainage (total)
- evaporation (total)

using any combination of the selectable inputs shown in Table 14.

Example output - soil nitrate-nitrogen at the end of various fallow lengths

Soil nitrate available at the end of the fallow increases with fallow length and will vary widely in response to differing soil water and temperature conditions in the fallow (Figure 20).

Howwet?

The *Howwet?* program has a fallow mineralisation calculator (Figure 20). It is a simplified version of the nitrogen module from APSIM program used to create *WhopperCropper* 'fallow'. However, it uses actual start and end dates of the fallow and actual rainfall records for the paddock in question, so can be a useful indicator of nitrate-nitrogen mineralisation.

Using yield and protein values of previous cereal crops to estimate soil nitrogen supply

Table of yield and protein results

Using the efficiency factors tabulated in Table 9 it is possible to estimate the minimum supply of plant-available nitrogen to recently grown cereal crops retrospectively using grain yield and protein values to estimate nitrogen (kg/ha) removed in grain. In this way, grain yield and protein for previous wheat or sorghum crops can be used to estimate a minimum supply of **soil nitrogen** for the next crop.

However, such estimates of likely nitrogen supply must be interpreted with care. Position of crop in the rotation could have considerable influence on estimating the future minimum nitrogen supply. The estimated future soil nitrogen supply might be over-estimated if:

- the preceding fallow was longer than 6 months
- the preceding crop was a grain legume.

However, even with these limitations, previous cereal crop production figures can be useful to obtain an approximate value for the soil's capacity to release plant-available nitrogen. Minimum soil nitrogen supplies, calculated from previous sorghum and wheat crops are tabulated in Tables 15 and 16.

Note that Tables 15 and 16 show how much nitrogen was removed by various grain yield and protein combinations. The nitrogen amounts shown must be factored up by an appropriate efficiency factor to account for the total amount of nitrogen required to grow the crop: grain and stubble (refer to Tables 8 and 9).

Recording grain yield and grain protein outcomes of recent cereal crops is a valuable way to qualitatively evaluate the conditions of soil nitrogen supply and soil water that the crop experienced (Table 17).

Critical grain protein values have been used to identify probable onset of nitrogen deficiency or to monitor the adequacy of nitrogen supply for the cropping system. High frequency of crop production of grain proteins below the critical value is indicative of the need to either commence regular application of nitrogen fertiliser or increase the level of nitrogen applied to a particular rotation. Low cropping intensity or drought periods may prevent widespread use of this strategy.

Table 15. The minimum supply of nitrate-nitrogen that was available to a crop estimated retrospectively from grain yield (t/ha) and protein content (%) of a recent sorghum crop. These values are based on grain nitrogen only and thus must be factored up by an appropriate efficiency factor to account for the total amount of nitrogen required to grow the crop; grain and stubble.


Grain yield	Sorghum protein					
	8	9	9.5	10	11	12
1.0	13	14	15	16	18	19
1.5	19	22	23	24	26	29
2.0	26	29	30	32	35	38
2.5	32	36	38	40	44	48
3.0	38	43	46	48	53	58
3.5	45	50	53	56	62	67
4.0	51	58	61	64	70	77
4.5	58	65	68	72	79	86
5.0	64	72	76	80	88	96
5.5	70	79	84	88	97	106
6.0	77	86	91	96	106	115

See Tables 8 and 9.

Table 16. The minimum supply of nitrate-nitrogen that was available to a crop estimated retrospectively from grain yield (t/ha) and protein content (%) of a recent wheat crop. These values are based on grain nitrogen only thus must be factored up by an appropriate efficiency factor to account for the total amount of nitrogen required to grow the crop; grain and stubble.

Grain yield	Wheat protein						
	9	10	11	11.5	12	13	14
1.0	16	18	19	20	21	23	25
1.5	24	26	29	30	32	34	37
2.0	32	35	39	40	42	46	49
2.5	39	44	48	50	53	57	61
3.0	47	53	58	61	63	68	74
3.5	55	61	68	71	74	80	86
4.0	63	70	77	81	84	91	98
4.5	71	79	87	91	95	103	111
5.0	79	88	96	101	105	114	123
5.5	87	96	106	111	116	125	135
6.0	95	105	116	121	126	137	147

See Tables 8 and 9.



Source: Strong pers.comm.

Figure 22. Wheat protein outcome and grain yield relative to yield with nitrogen unlimited



Figure 23. Frequency of response to nitrogen fertiliser for > 200 wheat crops categorised by protein content of unfertilised grain

Values of grain protein percentage alone can indicate nitrogen sufficiency

Results of numerous multi-rate nitrogen fertiliser experiments with cereals in Australia’s northern region indicate robust relationships between the grain protein outcome and potential for cereal crops wheat, barley and sorghum, to respond to applied nitrogen.

Relative wheat yield (defined as crop yield relative to the yield with unlimited nitrogen supply) has a relationship with grain protein which is depicted in Figure 22. According to this relationship, wheat grain protein of 11.5% is produced at approximately 90% relative yield, which is normally accepted as reflecting the most economic grain yield in multi-rate fertiliser experiments.

Frequency of response to nitrogen fertiliser

As further evidence of a relationship between grain protein and low crop nitrogen supply, results of more than 200 dryland nitrogen fertiliser trials showed that wheat protein of 11.5% or less was produced on sites which have high frequency of response in grain yield to nitrogen fertiliser (Figure 23). This means that if a wheat grain protein of less than 11.5% was produced, grain yield would have been increased if additional nitrogen was available. Similarly, critical levels of grain protein to that indicated for wheat (11.5%) have been discovered for barley and sorghum using trial data available from multi-rate N fertiliser experiments. The critical grain proteins for barley (dry grain) and sorghum (moist grain) are 10.5 and 9.5% respectively.

Table 17. Using past grain yield and protein outcomes to reflect qualitatively on the water and nitrogen supplied to cereal crops of northern Australia.

Qualitative yield and protein outcomes	Low protein	High protein
Low grain yield	Likely N deficiency	Low water supply or other limiting factor
High grain yield	Higher than average water supply	Rarely produced

References

Dalgliesh N P and Foale M A (1998). A guide to soil sampling. In: Soil Matters –monitoring soil water and nutrients in dryland farming, Eds. N Dalgliesh and M Foale, CSIRO, Australia.

Jones P N (1994). Sample size for on-farm soil sampling. CSIRO. Brisbane. IPP&P Biometrics Unit Report IPPP/BU/94/15.

Keating BA, Carberry PS, Hammer G L, Probert ME, Robertson M J, Holzworth D, Huth N I, Hargreaves JNG, Meinke H, Hochman Z, McLean G, Verburg K, Snow V, Dimes JP, Silburn M, Wang E, Brown SL, Bristow K, Asseng S, Chapman S, McCown RL, Freebairn DM and Smith CJ. (2003). An overview of APSIM, a model designed for farming systems *European Journal of Agronomy* **18** 267-288.

6. Determining nitrogen fertiliser requirement

Nitrogen fertiliser requirement = crop nitrogen demand – soil nitrogen supply

see Chapter 4

see Chapter 5

Crop demand

As described previously, **crop nitrogen demand** may be estimated from a target yield and a relevant grain protein.

Soil supply

As described previously, **soil nitrogen supply** may be estimated by field soil sampling or computer programs (*Howwet?* or *WhopperCropper*).

Nitrogen fertiliser calculator

Also supplied with this manual is an easy-to-use computerised calculator (Figure 24). On a single page the relevant data are entered for:

- Soil sampling date
- Expected planting time
- Stored soil water
- Expected season type (e.g. poor, average etc)
- Soil test results.

The calculator estimates:

- Nitrogen mineralisation from soil sampling to planting
- Crop demand
- Soil supply
- Nitrogen fertiliser rate required.

Also included in the calculator is an estimate of the losses incurred if the season turns out differently to that targeted or anticipated.



Figure 24. Nitrogen fertiliser rate calculator (included with this manual)

7. Alternative methods to calculate nitrogen fertiliser requirement

Key messages

- *WhopperCropper* is an easy-to-use program for calculating the effects of varying input levels (including nitrogen fertiliser) on potential yields using the full range of historical rainfall data.
- Having access to the full range of potential outcomes is superior to a calculation involving a single 'district average' because the user can readily evaluate how their attitude to risk is matched by the effect of different input on potential yield and gross margin outcomes.
- *WhopperCropper* is the best way to analyse the long-term financial aspects of varying input levels. A small amount of data entry is required.
- Grain nitrogen removal is 20 kgN/ha and 17 kgN/ha for each tonne of wheat and sorghum respectively. The total soil nitrogen requirement to grow a crop with adequate grain protein is typically 1.7 times this amount.

An example simulation output is shown below but by using the product personally, scenarios of particular interest to the user can be derived.

Simulating crop response to applied nitrogen with WhopperCropper

Crop responses simulated with *WhopperCropper* can be used to compare effects of nitrogen fertiliser application rate, stored water at planting, soil nitrogen supply, planting time, plant population and crop maturity, SOI phase and combinations of all of these factors. The program uses 100 years of weather data from selected sites and consists of a database of scenarios of all possible combinations of practical input levels that farmers must consider.

The potential range of responses to applied nitrogen fertiliser is demonstrated in Figure 25. The analysis applies to wheat yield at Dalby with five levels of applied nitrogen, 0, 25, 50, 100 or 150 kg/ha where soil contains 50 kg/ha plant-available nitrogen and the soil profile is fully wet (240 mm) every year. Other setup parameters: medium wheat maturity, 30 May planting, density of 100 pl/m².

In the absence of fertiliser nitrogen, 75% of all yields would be expected to be less than 1250 kg/ha. With the addition of nitrogen, the range of potential yields increased with both 25 and 50 kgN/ha. Rates of N higher than 50 kg/ha increased the average yield slightly because of the potential to achieve high yields in the years of high rainfall. However median yields increase very little because the yields remain evenly spread across low and high yields.

WhopperCropper also has the facility to enter costs and prices and hence gross margins can be calculated. In Figure 26 it can be seen that the median gross margin declines with rates greater than 100 kgN/ha. In addition, the proportion of 'lower-end' gross margins increase with the 100 kgN/ha rate.

Nitrogen fertiliser rate based on grain removal rate

Table 18 indicates the typical nutrient removal rates per tonne of grain. Use this as a guide only because there can be considerable variation around these values because of different grain protein levels that may occur. Nutrient replacement rates below crop removal will increase soil fertility decline. Nutrient replacement while continuing to crop will slow fertility decline rather than arrest the decline. Because of highly variable seasonal rainfall, application strategies that rely on nutrient removal rates of previous crops make nutrient applications a risky option in the short-term because:

- the supplied nitrogen fertiliser may not be ultimately required when rainfall is low and a low yield occurs




Figure 25. Wheat yield at Dalby with five levels of applied nitrogen, 0, 25, 50, 100 or 150 kg/ha where soil contains 50 kg/ha plant-available nitrogen and profile is fully wet (240 mm). Other setup parameters: medium maturity, 30 May planting, density of 100 pl/m².




Figure 26. Gross margins for wheat at Dalby with five levels of applied nitrogen, 0, 25, 50, 100 or 150 kg/ha where soil contains 50 kg/ha plant-available nitrogen and profile is fully wet (240 mm). Other setup parameters: medium maturity, 30 May planting, density of 100 pl/m², wheat grain price \$350/t, total non-nitrogen fertiliser costs \$160/ha, nitrogen fertiliser costs \$2.50/kg (\$1150/t of urea – priced at Oct 2008).

- the cost of supplementing the supply of the nutrient may not be economical in relation to market returns for the crops
- replacement of the nutrient may not match the original form or distribution, thus failing to simulate the original supply of plant available nutrient
- environmental sustainability may be affected by either over or under supply of nutrients in some situations.

Table 18 provides information to estimate nutrient removal by a crop or a sequence of crops. However, use the procedures described elsewhere in this manual and the electronic calculator as the preferred means to estimate crop nutrient needs.

Table 18. Quantitative removal (kg/t grain) of nutrients by grain crops common to Queensland

Crop	Typical nutrient removal (kg/t of product)						
	N	P	K	S	Ca	Mg	
Sorghum	19	3.7	4.8	2.8	0.5	1.4	
Wheat	23	3.4	4.5	1.8	0.5	1.4	
Maize-grain	24	3.3	5	3.8	0.2	0.9	
Sunflower	26	4.1	8.0	4.0	2.0	2.2	
Barley	20	2.7	5.0	2.0	0.4	1.2	
Chickpea	42	2.0	5.3	-	1.6	0.6	
Peanut-pods	52	5.1	7.1	-	0.6	2.0	
Peanut – hay	13	0.5	-	-	-	-	
Soybean	73	5.9	15.9	-	-	-	

Source: Dalal, R. C. and Probert, M, E (1997)

References

Dalal, R. C. and Probert, M, E (1997). Soil nutrient depletion. In: Sustainable crop production in the tropics, an Australian perspective. Eds. A.L Clarke and P.B Wylie. Department of Primary Industries, Queensland.

8. Evaluating the financial risk of applying nitrogen fertiliser in south Queensland

Key messages

- Without addition of nitrogen, decline in soil nitrogen fertility is inevitable. At some point in time, nitrogen deficiency will reduce profitability.
- Accurately predicting crop nitrogen demand for an individual crop is not possible due to significant seasonal variability.
- The decision-support package *WhopperCropper* can display the range of seasonal yields with the aim of choosing an appropriate target yield. The aim is to reduce grower exposure to financial risk although there can be no guarantee that a result in a single year will be as desired.
- Use *WhopperCropper* to display the full range of potential gross margins using the selectable range of nitrogen fertiliser rates.
- Estimates of plant-available soil nitrogen and water are needed to minimise financial risks where soil nitrogen supply is likely to be very low or very high.

The role of field trials to diagnose nitrogen requirement

Since the 1960s when crop responses to nitrogen fertiliser were first identified in southern Queensland, numerous field trials were conducted to advance our capacity to predict nitrogen requirements of cereals (Little *et al.* 1969, Strong *et al.* 1978, Dalal *et al.* 1998, Strong *et al.* 1996 a and b). Another aim of early field research was to develop a diagnostic technique to reliably predict future crop nutrient requirements. Soil testing was the primary diagnostic technique that held most promise at that time.

Soil tests for nitrogen, phosphorus and zinc were soon discovered to provide less skill for determining crop fertiliser requirements in southern Queensland farming systems than for more reliable rainfall environments such as the USA and UK. Other diagnostic approaches were therefore sought. In 1980s use of crop production information (grain yield and protein concentration) to estimate cereal crop nitrogen supply was promoted for use as a monitoring tool for grain growers to estimate when more frequent responses to applied nitrogen could be obtained. Results from previous multi-rate fertiliser experiments resulted in the use of grain protein as a nitrogen sufficiency indicator. Applications of these data is presented in this manual.

Evaluating the financial risk of applying nitrogen fertiliser to grain crops

Financial risk is important in a decision to apply nitrogen fertiliser in south Queensland, because of its variable rainfall. Recognition of this variability, and its effect on seasonal crop production in dryland farming systems of northern Australia, led to development of crop simulation models. Many field trials were used to derive principles of crop nutrition that provided algorithms for the crop simulation models. Field trials were used, and continue to be used, to validate or justify simulation outputs. Simulation models such as *Wheatman*, APSIM and its derivative products (*WhopperCropper*, *Yield Prophet* and *Howwet?*) have found applications in decision making for many grain growers in the region.

WhopperCropper (Nelson *et al.* 2002) is very relevant for nitrogen management decisions for cereal crops of northern Australia. Using *WhopperCropper*, the likelihood of unprofitable fertiliser use can be assessed as well as the frequency of profitable outcomes from any nitrogen management strategy. *WhopperCropper* provides yield outputs that are estimates over long-term cropping that are impossible to acquire by any other means. By adding costs and prices, the full range of potential gross margins can be generated. Hence the risk of negative gross margins can also be displayed.

The effect of soil water at planting on gross margin (with no nitrogen limitations)

As demonstrated previously (Figures 11 and 12) the soil water available at planting significantly affects the potential yield range. Figure 27 demonstrates the impact on the gross margins of wheat and sorghum at Dalby. The simulation assumes wheat price of \$300/t, variable costs (excluding N fertiliser) of \$140/ha, sorghum price of \$200/t, variable costs (excluding N fertiliser) of \$160/ha, and a nitrogen fertiliser cost of \$2.50/kgN (approx. \$1150/t urea) when applicable to the simulation.

Note: Figure 27 demonstrates the scenario using prices relevant at October 2008. The reader should use *WhopperCropper* with current prices and costs to construct comparative scenario analyses.

Other input options include using soil with a total water-holding capacity equal to 240 mm (PAWC) with a one third full (80 mm), two thirds full (160 mm) and full (240 mm) profile of plant available water.

How to read this *WhopperCropper* boxplot

The black line in the 'red box' is the median yield (50% of all years have this 'yield' or less). The dashed line within the 'red box' is the mean yield. The upper edge of the 'red box' is 75% probability. This is read as 'in 75% of years, yields will be less than this value'. The lower edge is the 25% probability value i.e. 'in 25% of years, yields will be less than this value'. The upper and lower short horizontal lines represent 100% and 0% probabilities respectively.




Figure 27. Effect of soil water at planting of wheat and sorghum yields at Dalby with stored water at planting equal to one third, two thirds or full at planting. Soil nitrogen set to 150kgN/ha at planting. Planting dates as shown. Density: Wheat 100 pl/m², sorghum 6 pl/m², medium maturities, soil PAWC=240 mm, sorghum solid 1 m rows.




Figure 28. The effect of varying N fertiliser rate at different levels of starting soil water on wheat gross margin at Dalby. Soil nitrogen was set to 25 kg/ha. Other parameters are the same as in Figure 27.

Figure 27 shows that:

In wheat:

- Simulated median gross margin for wheat increases by approximately \$400/ha for each 80 mm increase in plant-available water at planting
- Approximately 20% of the gross margins for wheat are negative when the soil water recharge is only one third full at planting. Planting with an extra 80 mm of stored water (two thirds full = 160 mm) eliminates the risk of negative returns from wheat. This demonstrates the crop reliance in south Queensland of water stored in the soil rather than the small quantity of in-crop rainfall that occurs during winter.

In sorghum:

- Filling the soil to two thirds full significantly reduces the risk of negatives gross margins from almost 20% to less than 5%.
- A full profile increases sorghum gross margin and dramatically reduces the risk of low returns.

The interaction of stored soil water and nitrogen fertiliser rate

Figure 28 demonstrates the impact on wheat gross margin at Dalby of a soil with a total water-holding capacity equal to 240 mm (PAWC) with a one third full (80 mm), two thirds full (160 mm) and full (240 mm) profile of plant available water AND three rates of nitrogen fertiliser (0, 50, 100 kgN/ha). Soil nitrogen level was set to a low value of 25 kgN/ha. The same wheat variable costs as above are assumed.

- When the profile is only one third full at planting, gross margin is negative in 50% of years when there is no nitrogen fertiliser applied. This reflects the high risk of the low soil water at planting as well as nitrogen deficiency that negates most positive returns in better seasons. With 50 kgN/ha applied a negative return occurs in only 20% of years. Because of the high cost of the 100 kgN/ha rate, the average and median returns are less than for the 50kgN/ha rate. High returns are evident only in the better seasons when in-crop rainfall interacts positively with the high nitrogen supply.
- With the profile two thirds full at planting, a nil nitrogen rate severely limits yield potential. 50 kgN/ha (75 kg/ha total available N) produces positive gross margin outcomes but nitrogen could still be limiting in good seasons. 100 kgN/ha (125 kg/ha total available N) provides equal or greater gross margins than the 50 kgN/ha rate in 75% of seasons. Only in the worst 25% percent of seasons is the gross margin less.
- A full profile at planting provides high returns with any N rate.

The decision to plant on a two thirds full profile rather than wait for a full profile may be influenced by factors such as the time in the planting window (yields reduce with later plantings) and the need to guarantee cash flow.

The nitrogen rate selected can be influenced by financial factors and the grain price, the amount of stored soil water, and individual opinion on the outlook for coming season.

Incorporating the SOI seasonal climate forecast into decisions to minimise the financial risk of applying nitrogen

Determining if the SOI phase system has skill

The SOI phase system uses a two-month indicator period to change the rainfall probability for the next three months. Yield ranges may therefore be modified according to SOI phase. However the skill with which the SOI phase system may accurately forecast the yield range differs by location and by time of year. The Rainman v 4.3 program has tables, graphs and maps that indicate when the SOI phase has skill for rainfall occurrence. The *WhopperCropper* program has a statistical section that indicates if the scenarios, including SOI, are statistically different from each other.

Using *WhopperCropper* to produce a gross margin analysis of a three way interaction of soil water at planting, nitrogen fertiliser rate and SOI phase

Figure 29 describes a three-way interaction of:

- soil water at planting (one third and two thirds full)
- nitrogen fertiliser rate (0, 50 kgN/ha)
- SOI phase (positive and negative).

The other setup factors are as for previous scenarios.

Figure 29 shows that knowledge of the soil water at planting and SOI phase might be used to reduce risks associated with the application of nitrogen fertiliser. Note that there are large differences in the median and variance of the potential outcomes.

For a **negative phase of SOI** (over April/May) the following outcomes are indicated:

- the predicted gross margin range with one third full (80 mm) stored water at planting indicates a 75% chance of negative returns without fertiliser
- applying 50 kgN/ha with a one third full profile results in less frequent though greater downside risks as well as acceptable returns in better seasons
- A soil profile that is two thirds full reduces frequency of downside risks, eliminating negative returns altogether with application of 50 kgN/ha. The outcome is very similar to that of the positive SOI phase.

In summary, when the SOI phase is negative, application of nitrogen fertiliser would present a high and perhaps unacceptable risk if the soil water levels at sowing were below 80 mm.

However, with a **positive SOI phase** (during April/May):

- Supplying 50 kgN/ha fertiliser (giving a total soil N supply of 75 kgN/ha), but with a one third full soil profile, creates a wide range of possible outcomes. Crop returns are higher in better seasons but the cost of the 50 kgN/ha of fertiliser reduces the gross margins in poor seasons. Hence the scenario is extremely variable but the median and average returns are moderately positive
- the predicted gross margin range for the one third full profile (80 mm) with nil nitrogen applied at planting indicates that the low soil water reserves will limit yield in low rainfall seasons. In the better seasons, the low soil nitrogen levels will limit grain yield response. Hence there are no positive gross margin outcomes
- Applying 50 kgN/ha when the soil is two thirds full provides the most favourable (low) range of yields with a vary favourable median gross margin
- Even higher gross margins would be expected with higher nitrogen fertiliser rates if the soil was fully wet at planting (240 mm) (data not shown), particularly in positive SOI years.

In summary, at Dalby, stored soil water plays a bigger role in reducing variability and increasing potential returns than the SOI phase. However, the SOI phase indicates even greater financial risk in applying nitrogen fertiliser when the phase is negative than when it is positive especially when stored soil water is low.

Additional risk analysis options

Using the spreadsheet ‘Nitrogen Calculator’


On the accompanying CD is a nitrogen fertiliser rate calculator, the ‘Smart N Decision Calculator’.

This automates the fertiliser calculation process described in previous chapters and also has a section that automates the process of analysing the yield and gross margin outcomes if the season turns out differently to the one targeted or anticipated.

The nitrogen fertiliser rate is calculated based on user-entered soil water at planting and targeted (or anticipated) season type. The spreadsheet calculates:

- A target yield
- Crop nitrogen demand
- Soil nitrogen supply is calculated from soil sample test results
- The difference between the demand and supply is the recommended nitrogen fertiliser rate.

There is also a ‘risk analysis’ worksheet. This calculates the yield across ALL seasons with the chosen nitrogen fertiliser rate. This is contrasted with the yield expectation in each season type with the fertiliser rate that would have been ‘more appropriate’ for that season.



Profile refill, applied N fertiliser and SOI phase

Figure 29. Using SOI phase (negative or positive) to reinforce decisions to apply or not apply nitrogen fertiliser to wheat at Dalby with either low (one third full = 80 mm) or moderate (two thirds full = 160 mm) soil water at planting. Other setup factors as indicated above.

Using a 'set' nitrogen fertiliser rate as indicated by crop nitrogen removal

Experience of the full potential yield range combined with the knowledge that 20 kgN/t of grain is removed by each crop can give a starting point for a nitrogen fertiliser rate. For example, if the 'average' grain yield is 2 t/ha the average nitrogen removal will be 40 kgN/ha per crop. However, using this average value disregards the variability that occurs in soil water and nitrogen at planting due to soil type, seasonal and fallow length effects, and in-crop rainfall, and hence to potential yield ranges. Whilst the value will be 'roughly right' there will be cases of financial losses due to under- and over-fertilising that may have been otherwise avoided.

References

- Dalal, R.C. W. M Strong, E. J. Weston, J. E. Cooper, G. B. Wildermuth, K. J. Lehane, A. J. King, C. J. Holmes (1998). Sustaining productivity of a vertisol at Warra, Queensland, with fertilisers, no-tillage or legumes 5. Wheat yields, nitrogen benefits and water-use efficiency of chickpea-wheat rotation. *Australian Journal of Experimental Agriculture*, 38, 489-501
- Littler, J.W., Kelso, W.T., and Marley, J.M.T. (1969). Effects of nitrogen fertiliser on the yield and malting quality of barley on the southern Darling Downs, Queensland. *Queensland Journal of Agricultural and Animal Sciences* 26, 309-317.
- Nelson, R.A., Holzworth, D.P., Hammer, G. L., and Hayman, P.T. (2002) Infusing the use of seasonal climate forecasting into crop management practice in North East Australia using discussion support software, *Agricultural Systems*, 74 393-414.
- Strong, W.M., Best, E.K. and Cooper, J.E. (1978). Yield response of dryland wheat to nitrogen and phosphorus fertilisers. Queensland *Wheat Research Institute, Biennial report*, 1976-1978.
- Strong, W M, Dalal, R C, Weston, E J, Cooper, J E, Lehane, K J, King, A J, and Chicken, C J. (1996a) Sustaining productivity of a Vertisol at Warra, Queensland, with fertilisers, no-tillage or legumes 2. Long-term fertiliser nitrogen needs to enhance wheat yields and grain protein. *Australian Journal of Experimental Agriculture*, 36, 665-74.
- Strong, W M, Dalal, R C, Weston, E J, Cooper, J E, Lehane, K J, King, A J, and Chicken, C J. (1996b) Sustaining productivity of a Vertisol at Warra, Queensland, with fertilisers, no-tillage or legumes 3. Effects of nitrate accumulated in fertilised soil on crop response and profitability. *Australian Journal of Experimental Agriculture*, 36, 675-82.

9. Applying nitrogen fertiliser in south Queensland

Key messages

- Calculate the elemental nitrogen rate required (previous chapters).
- Determine the cheapest or most convenient form of nitrogen that can be applied with the available equipment.
- Decide on application timing based on equipment available and work load requirements. The timing of the nitrogen application appears less important than satisfying the nitrogen demand.
- Observe the recommendations of the maximum quantity recommended with the seed or place urea at an appropriate distance from the seed row.

Calculating the cost of elements in nitrogen fertiliser products

Most producers will apply the cheapest source of fertiliser that is in a form suitable for their application equipment. When comparing the cost of fertilisers it is necessary to calculate the cost of the 'element' of interest.

When the tonnage price of the fertiliser is known, the actual cost of the elemental content can be calculated. The calculation is:

$$\text{Elemental cost (\$/kg)} = (\text{product cost (\$/t)} / 1000) / (\text{percentage of element} / 100)$$

e.g. cost of nitrogen in urea where urea cost = \$550/t,
percentage nitrogen = 46%
nitrogen cost = $(550 / 1000) / (46 / 100)$
nitrogen cost = \$1.20/kg

Phosphorus cost in MAP, percentage P = 22%
MAP cost = \$780/t
P cost = $(780 / 1000) / (22 / 100)$
P cost = \$3.55/kg

Table 19 indicates comparative prices, as at June 2015. Prices of the nitrogen fertilisers are usually linked to oil prices so updated prices should be obtained.

Readers are advised to check current fertiliser prices when making this calculation.

Common forms of nitrogen fertiliser

Urea

Urea is usually the cheapest form of solid nitrogen fertilizer. A bigger application boot is needed, so more soil disturbance can occur during application. However, while there is potential for soil moisture loss, damage to emerging seedlings is potentially less because of the larger quantity of soil into which the urea is mixed. Urea is most commonly applied prior to planting, or during planting if it can be placed away from the seed row. A three-bin cart is needed if applying both urea and starter fertiliser at planting time.

Anhydrous ammonia gas (NH₃)

This is usually the cheapest form of nitrogen fertilizer, and is widely used in irrigation and some dryland areas of south Queensland. The necessary plumbing is relatively cheap and easy to set up but it requires an extra trailing or mounted tank (see photo on the following page).

UAN (Urea ammonium nitrate)

This liquid product is currently more expensive (\$/kgN) than urea, although can be competitive at times. It is extensively used in Western Australia and is especially useful for in-crop applications in sandy soils. Some farmers are trialing Queensland. It can be placed relatively close to seed. There is a need for a mounted or trailed liquid tank, the necessary plumbing, and on-farm storage.

Table 19. Approximate comparative nitrogen fertiliser prices as at June 2015

Product	%N	Cost (\$/t, bulk)	\$/kg N
Urea	46	550	1.20
Anhydrous ammonia	82	950	1.16
Ammonium nitrate	34	Now 'dangerous goods'	
Ammonium sulphate	20.2	480	2.38
Feedlot manure	approx 3		
MAP	10	780	3.55
MAP + Zinc Compound	10.5	800	3.64
DAP #	18	780	4.33

'starter' (P and Zn) fertilisers are not used as sole sources of nitrogen but the nitrogen content can be included in a nitrogen budget

Timing of nitrogen application

In northern Australia, the traditional time to apply nitrogen fertiliser to cereal crops is before planting, usually after soil water has been recharged during a fallow period. Without follow-up rain, nitrogen applied immediately before or during planting may remain trapped in topsoil as it dries. If dry conditions continue, the crop may not access the applied nitrogen until topsoil water has been recharged, but in such situations crop yield and demand for N is usually lower anyway. If soil water recharge occurs late during grain filling the crop may still access applied nitrogen and respond with increased grain protein.

Research shows that although in any one year there may be an advantage due to applying nitrogen fertiliser either during the fallow or at planting time, over a period of time there is likely to be little difference. An adequate N supply to meet crop demand it is more important than the timing of application.

If soil water recharge occurs too late to benefit the fertilised crop, a high proportion of applied nitrogen will be carried over for use by subsequent crops in the rotation. Similarly where nitrogen is applied before planting and a planting rain does not eventuate, significant carryover. In south Queensland the losses from the system were found to be 5 to 25% percent depending on the season (Strong pers. com.).

Nitrogen deficiency in the early stages can affect the number of grains that are formed in the embryonic head. Subsequent nitrogen demand is driven by the rapidly developing biomass prior to flowering. Under favourable early conditions, high nitrogen supply may promote high vegetative biomass which in turn can use large amounts of soil water. Restriction of nitrogen supply can theoretically reduce this early demand but is unlikely to work in practice unless soil nitrogen is very low. In addition, the restriction in potential yield (crop sink) may be a disadvantage if the season becomes favourable.

Pre-plant nitrogen application

Advantages

- More opportunity for nitrogen to move into the profile
- Gets the nitrogen application job out of the way
- Only option for many farmers and planter set-ups.

Disadvantages

- May cause excessive moisture loss during application, which on occasions can jeopardize planting opportunities
- Requires earlier nitrogen fertiliser decision (and soil testing)
- Fertiliser cost is incurred without a guarantee of when you will be able to plant the next crop
- Increased risk of nitrogen losses due to waterlogging in fallow.



Anhydrous fertilizer applicator

A strategy to apply nitrogen well before planting has been commonly used by growers in northern Australia including south Queensland, in an attempt to separate nitrogen application from planting for logistical reasons mentioned above. This method presents growers with the dilemma of deciding to apply nitrogen under the assumption that soil water will be recharged sufficiently after the application to support a rain-fed crop. Where nitrogen is applied early and a planting rain does not eventuate, significant carryover of applied nitrogen to subsequent crops can occur.

Nitrogen applied at planting

Advantages

- Ensures expenditure on nitrogen fertiliser only occurs when a planting opportunity arises
- More time to decide if nitrogen is needed relative to soil water recharge
- More easily done in summer than winter cereals (because of wider rows)
- No loss of moisture or planting opportunity
- Less risk of waterlogging losses.

Disadvantages

- Higher workload at planting time – increased labour needs and lower efficiency of planting operation
- Requires specialised planter setup; a three-bin seed cart if starter fertilizer is also required, otherwise a anhydrous or UAN cart is required
- Risk of nitrogen being stranded in dry soil therefore lower nitrogen availability to the crop (but lower crop demand if it stays dry).

Application of nitrogen in-crop

- Last resort if nitrogen not applied earlier
- Needs to be applied in first 35-40 days (by end of tillering in winter cereals). Follow-up rainfall is needed for benefit to accrue (this makes it a risky practice in Queensland)
- Side-dressing
 - requires row crop equipment, straight rows
 - rainfall soon after application less critical

- Top-dressing
 - Needs rain immediately after application to avoid losses (of urea, ammonium sulphate), and soon after application of other products (ammonium nitrate)
- Foliar application
 - Only small amounts of nitrogen can be applied otherwise leaf burn may occur
 - Relies on rain soon after application for best response
 - Most of the nitrogen uptake occurs via the soil after being washed off the leaves.

In spite of the additional workload, nitrogen application at planting would appear worth pursuing to optimise its efficient use by the crop. The trend towards zero and reduced tillage, which usually extends the planting window, is another reason to delay nitrogen application until planting to avoid topsoil disturbance and soil moisture loss.

Nitrogen fertiliser placement

Placement and timing of nitrogen fertiliser will depend upon the type of available equipment and the need to match the nitrogen demand for the crop.

Pre-plant placement of nitrogen

- Generally band at less than 2 x seed row spacing. Not wider than 1 m spacing
- Minimise soil disturbance (and moisture loss)
 - Coulters/discs ideal
 - Narrow tyne and point
 - Only place deep enough to get coverage.

Placement of nitrogen at planting

Preferably nitrogen fertiliser should be placed at least 30 mm away from seed (unless applying very low rates or using wide points).

Maximum nitrogen fertiliser rate with seed.

Winter cereals 9 kgN/ha in 50 cm rows,
 18 kgN/ha in 25 cm rows

Sorghum 4.5 kgN/ha in 1 m rows

Placement options include:

Winter cereals – place nitrogen in every second inter-row space. This will enable a substantial N rate (40-75 kg/ha) to be safely applied at sowing and ensures that each row has access to nitrogen. It also minimizes soil moisture loss, stubble handling and machinery setup issues although it does require extra applicators.

Summer cereals – ideally place 50 cm away from row.

How much fertiliser can be placed in the seed row?

The maximum application rate of fertiliser in the seed furrow is primarily influenced by the susceptibility of the crop species to ammonia and salt (osmotic) effects, the chemistry of the fertiliser, soil conditions and application equipment. Hence, the safe nitrogen rate with seed will be lower in dry conditions and using narrow tynes, points or discs on wide row spacings. Conversely, under cool conditions and in very wet soil higher rates with the seed may be possible. The rates in Table 20 would indicate safe application rates in most conditions in south Queensland.

The effects of nitrogen fertiliser on crop germination can be seen in the photo above.

Example: for a narrow point opener for wheat on 36 cm row spacing, the safe urea rate with the seed is approximately 27 kg/ha (Table 20).

Foliar applied nitrogen

Foliar applications are of limited use because of the small amount of nitrogen that can be safely applied. Leaf ‘burn’ can occur at high rates (see photo below). Urea can be used as a foliar spray. A 30% solution (30 kg in 100 L of water) applied at 110-120 L/ha applies an equivalent of 15 kg N/ha. Repeat every 2-3 weeks as needed. Some leaf burn may be expected, but this generally does not affect subsequent grain yield. If there is no prior experience with foliar application to the crop, it may be wise to conduct a test strip to test its sensitivity.

This can give a short term benefit to the crop for example immediately following waterlogged conditions when there are signs of root growth but cannot be relied on to supply the full crop requirement.

Minimising nitrogen fertiliser costs

Using a strategic process, such as that described in this manual will help to optimise nitrogen fertilizer use and avoid excessive, risky or unnecessary expenditure.

In particular, use of soil tests will more accurately identify when nitrogen fertiliser is necessary and provide a basis for calculating a nitrogen fertilizer rate.

If the required nitrogen rate is excessive and/or the fertilizer cost is prohibitive, consider planting a pulse crop on a proportion of the farm to reduce nitrogen fertiliser costs. The use of legumes in the crop rotation to help manage nitrogen fertility is discussed in the next chapter. In general, pulses would be expected to add 0-30 kgN/ha, and grazed lab lab enough nitrogen, for the next grain crop. Butterfly pea will be of benefit if soil water is replenished prior to the next cereal and the BFP residues have decomposed.



Evidence of leaf 'burn' from foliar applied nitrogen



Evidence of reduced emergence when too much fertiliser was applied close to the seed in oats planted in March and subject to high temperatures that caused rapid soil drying.

Photo B Radford.

Gearing up for bulk fertiliser will further reduce costs.

Increasing length of fallow to accumulate more nitrate N is discouraged; savings in N fertilizer costs are likely to be offset by the more rapid decline in soil fertility as well as a decrease in cropping opportunities.

Applying spatial information to nutrient management

Two applications of precision agriculture are being explored in northern Australia to enable grain growers to maximise their returns:

- increasing capacity to monitor crop nitrogen requirements; and
- increasing capacity to distinguish areas within the crop of similar or contrasting grain protein for improved segregation of grain during harvest.

Grain yield and grain quality are rarely uniform over large areas of crop in south Queensland. Differences in soil type and soil depth are possible causes of variation in crop outcomes, although other soil and management factors also contribute to variation in crop performance. Production zones could be managed differently so as to optimise nutrient application where:

- similar variation in crop production occurs every cropping season
- components of 'precision agriculture', such as yield monitors and aerial imagery are available to gain knowledge of spatial variation.

Managing fertiliser input by zones would assist growers to produce grain of a target protein that attracts premium market returns. Even higher returns could eventuate if grain within the premium protein window could be identified prior to crop harvest, enabling better segregation of grain during harvest or blending at receipt to maximise the quantity of premium grain delivered.

In addition, monitoring nutrient management by zone is important for systems of continuous cropping because of the likelihood that fertiliser requirements may increase with continued cropping. Mapping grain yield and protein content of cereal crops (see Figure 30) should provide a guide to nitrogen requirements of subsequent cereal crops in the rotation.

Application of imagery captured from aerial or satellite platforms should increase grower capability to achieve these goals. At this point in time, protein monitors are not considered sufficiently accurate for widespread use.

Table 20. Approximate recommended maximum rates of actual nitrogen and urea (kg/ha) when applied in the seed rows in winter and summer cereal crops in Queensland. Rates are for typical heavy clay soils with very good seedbed soil moisture in the Queensland region in wheat, triticale, barley, oats¹, sorghum and maize¹ crops

	Seeder opener type					
	Disc opener#		Narrow point#		Sweep#	
	Safe rates with seed					
Seed/fertiliser row spacing	N (kg/ha)	Urea (kg/ha)	N (kg/ha)	Urea (kg/ha)	N (kg/ha)	Urea (kg/ha)
10" (25 cm)	14	30	18	39	23	50
12" (30 cm)	13	28	15	33	20	43
14" (36 cm)	11	24	12.5	27	17	37
20" (50 cm)	7	15	9	20	12	25
40" (100 cm)	3.5	8	4.5	10	6	13

Note: These rates have been adapted from data supplied by Dr C Dowling and data from R Heller, Alberta Reduced Tillage Linkages Canada (www.reducedtillage.ca) that has been modified in accordance with knowledge and experience of Queensland conditions. In general, these recommendations account for a wet seedbed that may subsequently dry quickly after planting, thus increasing the risk of damage from applied nitrogen.

refers to the 'relative' width of disturbed soil into which the fertiliser is placed and is approximated at 25 mm, 50 mm and 75 mm respectively. The actual degree of mixing can vary widely because of variations of soil texture, implement speed, tyne movement etc. Checking dispersion of crop seed in the disturbed soil can give an idea of how closely the fertiliser may be placed with the seed. The greater the mixing, the greater the margin of safety.

¹there is anecdotal evidence that these crops can tolerate approximately 10% more nitrogen with the seed than the table indicates.




Figure 30. Maps of yield, grain protein and grain moisture for a farm in southern Queensland

10. Nitrogen contribution of ley legumes and pulses

Key messages

- The nitrogen contribution from a legume depends upon its effective nodulation by the appropriate strain of root nodule bacteria and the amount of leaf (and root) material returned to the soil as litter or recycled through livestock.
- Nitrogen is removed when grain and hay is removed from the paddock. Grazing animals return much of the nutrients to the soil but may concentrate nutrients in patches and around watering and resting areas.
- A rough rule of thumb for nitrogen fixed by legumes is 2 to 2.5% of biomass i.e. 20 to 25 kg/tonne of biomass (under conditions of low soil N and effective rhizobial nodulation). The net contribution to the soil nitrogen pool is from 1 to 1.5%.
- Ley pastures generally add 0 to 100 kgN/ha to the soil N pool, depending on climatic conditions experienced during the pasture phase.
- Pulse crops generally add 0 to 30 kgN/ha and contribute most when biomass is high and grain yield is low.
- Lablab is a productive short term ley pasture and can contribute large amounts of N to the soil (40 to 80 kgN/ha/year).
- Lucerne will contribute 60 to 90 kg N/ha/year in the first 1 to 2 years but less thereafter unless seasonal conditions are very good.
- Soil water depletion by the ley can be a major constraint when returning to cropping after a ley phase. A ley phase, as with any phase within the rotation, depletes soil water that must be restored before sowing the next crop. Lucerne can dry soils to depth and slow soil water replenishment can be a constraint when returning to cropping.

Introduction

The provision of adequate nutrients to optimise crop performance, either by applying fertilisers or by exploiting inherent soil fertility, is a fundamental requirement of any sustainable cropping system. Over time, the inevitable nitrogen fertility decline associated with cultivation and nitrogen removal by crops has occurred.

A number of factors combine to make the use of ley pasture (including legumes) attractive as a low cost method of restoring soil nitrogen fertility in southern Queensland cropping systems, they include:

- the relatively high cost of nitrogen fertiliser
- high variability in seasonal rainfall, making prediction of optimum nitrogen fertiliser application rates difficult.

However, incorporating pastures into cropping land may be difficult because of:

- infrastructure costs (fences, watering points) where infrastructure is not already present
- pasture establishment costs, depending on species sown
- pasture establishment difficulties with small seeded species in heavy clay soils
- possible soil compaction if stock are not removed before rain
- weed management issues when re-cropping
- difficulty in re-charging the soil water profile after using deep rooting pasture species prior to re-cropping.

Factors affecting the nitrogen contribution from ley pastures and pulse crops

Ley pasture may consist of grass, legume or a mixture of grass and legume species, which is usually introduced between phases of cropping, to restore soil chemical and physical fertility, primarily by increasing soil carbon and/or nitrogen during the ley period. Pulse crops are leguminous crops that produce high protein seed used for human or livestock production.

A (ley) legume is a plant that grows with symbiotic root nodule bacteria that use ('fix') atmospheric nitrogen gases to create new nitrogen compounds for use by the plant. When residue from these legumes is returned to the soil, mineralisation occurs that makes extra nitrogen available for subsequent crops. Thus, legume residues may partially or totally replace the nitrogen requirement of subsequent cereal crops. Nitrogen contribution from decaying legume plants will generally become available to subsequent crops at a faster rate than decaying grass pastures or cereal stubbles. This is because

the lower carbon to nitrogen (C:N) ratio of the legume material better matches the C:N ratio of the food for micro-organisms. The decay rate is optimised as no nitrogen has to be accessed from the soil. If material with a high C:N ratio (such as cereal stubble or grass residues) is incorporated, nitrogen for cell synthesis in micro-organisms is obtained from the soil, temporarily 'tying-up' nitrogen that would otherwise be available to plants thus reducing availability of nitrogen in soil.

However, combining a grass with a legume species is an important strategy because the grass becomes a reservoir for the nitrogen fixed by the legume growing in combination with the grass. This encourages continued nitrogen fixation by the legume. Thus, a greater quantity of nitrogen is ultimately returned to the soil, when legumes are in mixtures with grass. The additional fibrous grass material also contributes positively to soil physical structure and a higher soil carbon level.

Nitrogen contribution from decaying pasture may become available to subsequent crops at a slower rate than a comparable rate of mineral fertiliser. This may better match crop nitrogen requirements as the crop grows. Nitrogen (mainly from decaying roots) may also be placed deeper in the soil. Thus cereal crops will access this nitrogen later in their development when demand for nitrogen is high. If dry matter material does not mineralise rapidly, nitrogen may be 'tied-up' in microbial biomass creating a temporary nitrogen deficiency in the cereal crop.

Balance of nitrogen removed and nitrogen retained

The nitrogen contribution of legumes will depend upon the quantity of vegetative biomass produced that remains in the paddock – either as litter or recycled through livestock – for decomposition. Biomass production is effected by climatic conditions, and biomass production and nitrogen fixation are greater in wetter conditions. Individual seasons can vary widely with a subsequent wide variation in biomass production. Table 21 details the variation that can occur across regions.

Other soil nutrients

Biomass production and nitrogen fixation will only be maximised if soil nutrients such as phosphorus, zinc and sulphur are in adequate supply. Soil testing prior to planting will give an indication of soil nutrient levels and fertiliser requirements. On highly alkaline soils, phosphorus, zinc and sulphur fertilisers may be required. Sulphur soil tests may indicate low levels in surface soils but often adequate quantities at depth. Superphosphate is a convenient source of phosphorus and sulphur.

Key factors to consider before planting pastures

Note: These management processes are described in detail in the LeyGrain Manual (Lloyd *et al.* 2007). Some factors are listed below.

Before sowing

- What are the reasons for returning a cropping paddock to pasture?
 - Low organic carbon status and hence low nitrogen fertility (a common indicator is consistently low grain protein from cereal crops)
 - Physical deterioration of surface soil e.g. crusting
 - Marginal water-holding capacity of the soil makes cropping risky
 - Economic returns from livestock vs cropping
 - Crop diseases or weeds that may be controlled by a break crop
- Consider what long-lasting residual herbicides were used in the previous cereal crop and observe plant-back periods
- Fallow to accumulate soil water, a critical issue that is often overlooked with pasture sowings
- Prepare soil seed-bed appropriately
 - Similar for a grain crop using conventional tillage techniques is usually most appropriate
 - Consider sowing into stubble after a winter grain crop, providing the soil profile water requirements have been met
- Purchase high quality seed but compensate sowing rate if germination rate of seed is low
- Consider the potential weed species that may have to be controlled.

Overall – Prepare for establishing a pasture as you would a crop!!

At sowing

- Use the appropriate seeding rate
- Sow at the correct time
 - Tropical grasses – mid January to mid March
 - Tropical legumes – spring/summer
 - Temperate species – autumn (April/May)
- Apply fertiliser if required (especially P fertiliser for legumes)
- Inoculate with specific Rhizobium bacteria
- Sow at the correct depth
 - Small seeds – 1 to 2 cm
 - Larger seeds – no more than 7 cm

Table 21. Increase in soil N (kg/ha/year) following legume leys in southern inland Queensland.

Region	Soil type	Forage species	Mineral N provided (kgN/ha)
Western Downs	Heavy clay	Butterfly pea	53
		Burgundy bean	45
		Lablab	65
		Lucerne	56
Balonne	Heavy clay	Medic	35-55
		Lucerne	45-95
		Grass/legume	30-50
	Sandy loam	Medic	45
		Lucerne	45
		Grass/legume	15

Source: LeyGrain- Pastures in Farming Systems. Sustaining Profit and the Environment.

- The advantage of seed pelleting is not clear
 - Pelleted fluffy grass seeds will flow better through conventional planters.
 - Remember to compensate for the weight of the pelleting material when calculating the seeding rate
- Consider applying an ant protection coating
- Scarify hard-seeded medics.

Undersowing with cover crops

- Most commonly used when sowing temperate species though there are indications that some tropical grasses and legumes can be undersown in autumn/winter
- Decreasing order of competitiveness of cover crops: oats > barley > wheat
- Reduce sowing rate of cover crop compared to grain or forage crop by at least 50%.

Nitrogen accumulation from leguminous pasture

Nitrogen accrual from pasture systems is generally positive although widely variable. The quantity of nitrogen fixed is generally in the range of 2-2.5% of above ground dry matter production and the amount accrued in the soil is from 1-1.5% of above ground biomass production.

Lloyd *et al.* (2007a) cite the range of nitrogen contributions of 15 kgN/ha for grass/legume pasture on a sandy loam soil, up to 95kgN/ha from lucerne on a heavy clay soil (Table 21). Losses of nitrogen of 10 and 20 kg/ha in faeces and urine were reported by Steele and Wallis 1988, cited by Doughton and Holford (1997). The nitrogen returned by animals is often accumulated in patches in the paddock and in shade lines.

Most nitrogen fixed by the legume is transferred from nodules/roots to plant tops; chickpea roots and nodules contained only 6% of the quantity of N contained in plant tops. Therefore, fate of N contained in legume tops is critical to the role legumes play in the N economy of the farming system. Hence, there is likely to be a greater N contribution from a grazed ley legume than from a harvested grain legume, since a very high proportion of pulse N is contained in its seed; 66-89% of N fixed by chickpea crops was removed in seed (Doughton and Holford 1997).

Medic

Over eight rotations of medic-wheat tested at Warra on the Darling Downs, the nitrogen contribution from medic was directly related to March to September rainfall and was equivalent to 40 kgN/ha/t DM produced (4.0% of above ground biomass) (Weston *et al.* 2002). Water available in the soil after a medic phase was similar to levels after continuous wheat.

Lucerne

Nitrogen produced by above ground lucerne biomass was directly related to total annual rainfall (26 kgN/ha/t DM produced or 2.6% of above ground biomass production) and to October to September rainfall (36 kg N/ha/t DM produced or 3.6% of above ground biomass production) (Dalal *et al.* 2004). Water available in the soil after a lucerne phase is almost always less than after continuous wheat.

Grass/legume pastures

Nitrogen produced by different pasture is also directly related to rainfall. (Medics, lucerne and Rhodes grass produced 17.2 to 20.5 kg N/ha for each 100 mm rainfall with the grass contributing 5.2 to 7.0 kg N/ha and the legume 10.6 to 13.2 kg N/ha for every 100 mm rainfall (Strong *et al.* 2006) .

Potential losses of contributed nitrogen

The nitrogen mineralised from pasture legumes is subject to the same potential losses as that from nitrogen fertiliser. Mineralisation from a pure legume pasture can be quite rapid and hence the accumulated nitrate may be subject to loss. The risk of loss may be lower after a grass/legume ley phase because of the slower rate of mineralisation of organic to mineral nitrogen.

Soil water considerations when returning to cropping

It is a common experience that deep-rooted perennial pasture will deplete soil water stores more than an annual crop. In dry seasons it may take some time for the soil moisture profile to be recharged after a pasture phase. Hence, the first crop following a pasture phase may incur a yield penalty because of less available water at planting. In very dry seasons crop planting may need to be delayed a season or more until the profile is recharged. This has occurred following lucerne-based pastures in southern Queensland and following

butterfly pea in trials in central Queensland. Removing ley pasture when soil moisture is high or is likely to be recharged may minimise the risk of a yield penalty in the next grain crop.

Ley legumes suited to southern Queensland

The material from this section is derived from the LeyGrain Manual (Lloyd *et al.* 2007a).

In southern Queensland, the dominant ley legumes and grasses are best classified according to rainfall and soils. Tables 22 and 23 summarise the adaptation and attributes of those pasture species. It is advisable to seek advice from agronomists working in the regions in order to clarify current experience and management practice.

Details of legumes suited to southern Queensland (from Lloyd *et al.* 2007b)

Temperate annual legumes

Annual medics (*Medicago spp.*) that are adapted to the neutral to alkaline soils include: **Barrel** (*M. truncatula*), **Snail** (*M. scrutella*) and **Spineless burr** (*M. polymorpha*) (Lloyd *et al.* 2007b).

Species that are adapted to the acid to neutral soils include: **subclovers** (*Trifolium subterraneum*). These are appropriate for the granite belt and traprock soils. **Serradella** is adapted only to the sandy friable soils of the granite belt and west of the Condamine River.

Woolly pod vetch (*Vicia villosa*) vetch is widely adapted but seed set has had variable reliability.

Temperate perennial legumes

Lucerne (*Medicago sativa*) is widely adapted but is best suited to deep alkaline soils. It is relatively tolerant of sodic and alkaline soils and is easy to establish. It is highly nutritious and promotes high liveweight gains. Bloat can be an issue, though this can be alleviated through management. It is a medium term ley that thins over time owing to its susceptibility to disease in wet and dry conditions. Soils can be dried to depth and soil water replenishment can be slow in dry seasons.

Sulla (*Hedysarum coronarium*) is a new short-term perennial that is best adapted to calcareous soils. It is non-bloating and is extremely productive in wet winters. Its persistence is enhanced by management that allows seed set. The root rotting diseases *Rhizoctonia solani* and *Sclerotium rolfsii* can affect sulla in the subtropics.

Tropical annual (and self-regenerating annual) legumes

Lablab (*L. purpureus*) is a high-producing annual adapted to alkaline clay and clay-loam soils.

Wynn Cassia (*Chamaecrista rotundifolia*) is a self regenerating annual for drier environments that is adapted to sandy acid soils.

Tropical perennial legumes

Burgundy bean (*Macroptilium bracteatum*) is a short-term perennial that can be highly productive and regenerate from seed.

Desmanthus (*D. virgatus*) is a deep rooting perennial, suited to long term pastures, that is persistent and drought hardy though competition can cause slow early development. This legume is well adapted to clay soils.

Caatinga stylo (*Stylosanthes seabrana*) is also suited as a long term component of permanent pastures and as a short-term ley species on a wide range of clays, clay-loams, loams and sandy-earths. It is adapted to colder areas where it has better frost survival, seedling regeneration and production capabilities than other stylos.

Details of grasses suited to southern Queensland (adapted from Lloyd *et al.* 2007b)

The most important species of grasses for production in southern Queensland have included:

Buffel (*Cenchrus ciliaris*) – the most widely adapted summer-growing species and the cornerstone of the beef industry. Decline in pasture productivity is evident and under investigation.

Panic (*Panicum maximum*) Green panic colonised large areas of fertile brigalow soils but new sowings include cultivars Gatton that has wider soil adaptation.

Bambatsi panic (*Panicum coloratum*) is particularly well adapted to high-clay soils. Establishment is more reliable than all other grasses except Purple pigeon grass.

Purple pigeon grass (*Setaria incrassata*) was released principally as a solution to poor establishment of grasses on heavy clay soils.

Creeping blue grass (*Bothriochloa insculpta*) is well adapted to alkaline loams, clay loams and friable clays.

Premier digit grass (*Digitaria eriantha*) is very widely adapted but best adapted to clay-loams, loams and sandy soils

A summary of the persistence and ground cover of grasses sown in adaptation trials, seven years after a sowing in 1983 is detailed in Table 24.

Table 22. Legumes for the Northern grain belt. Source: Lloyd et al (2007a) LeyGrain – Pastures in Farming Systems. Sustaining Profit and the Environment

Minimum rainfall	Legume species	Growing season	Tolerances					Soil preferences					Pasture duration
			Water logging	Alkaline	pH		Salt	Black earth	Heavy clay	Friable clay	Loam	Sand	
650 + mm	Arrowleaf clover (<i>Trifolium vesiculosum</i>)	W	1	x	✓	✓	1	x	x	✓	✓	✓	SA
	Balansa clover (<i>Trifolium michelianum</i>)	W	4	x	✓	✓	4	x	x	✓	✓	✓	SA
	Butterfly pea (<i>Clitoria ternatea</i>)	S	1	✓	✓	x	1	✓	✓	✓	x	x	P
	Persian clover (<i>Trifolium resupinatum</i>)	W	3	✓	✓	✓	1	x	✓	✓	✓	x	SA
	Strawberry clover (<i>Trifolium fragiferum</i>)	W	3	x	✓	✓	3	x	x	✓	✓	✓	P
600 + mm	Biserrula (<i>Biserrula pelecinus</i>)	W	1	x	✓	✓	1	x	x	x	✓	✓	SA
	Burgundy bean (<i>Macroptilium bracteatum</i>)	S	1	✓	✓	x	1	✓	✓	✓	✓	x	SP
	Caatinga stylo (<i>Stylosanthes seabrana</i>)	S	1	✓	✓	x	1	x	✓	✓	✓	x	P
	Desmanthus (<i>Desmanthus virgatus</i>)	S	1	✓	✓	x	1	✓	✓	✓	x	x	P
	Gland clover (<i>Trifolium glanduliferum</i>)	W	1	x	✓	✓	1	x	x	✓	✓	✓	SA
	Subterranean clover (<i>Trifolium subterraneum</i>)	W	1	✓	✓	✓	1	✓	x	✓	✓	✓	SA
	Sulla (<i>Hedysarum coronarium</i>)	W	1	✓	✓	x	1	✓	✓	✓	✓	x	SP
550 + mm	Lucerne (<i>Medicago sativa</i>)	S/W	1	✓	✓	x	2	✓	✓	✓	✓	x	P
	Murex medic (<i>Medicago murex</i>)	W	1	x	✓	✓	1	✓	✓	✓	✓	x	SA
	Snail medic (<i>Medicago scutellata</i>)	W	1	✓	✓	x	1	✓	✓	✓	✓	x	SA
	Sphere medic (<i>Medicago sphaerocarpus</i>)	W	1	x	✓	✓	1	✓	✓	✓	✓	x	SA
	Woolly pod vetch (<i>Vicia villosa</i>)	W	2	✓	✓	✓	1	✓	✓	✓	✓	✓	SA
450 + mm	Barrel medic (<i>Medicago truncatula</i>)	W	1	✓	✓	x	1	✓	✓	✓	✓	x	SA
	Burr medic (<i>Medicago polymorpha</i>)	W	1	✓	✓	x	1	✓	✓	✓	✓	x	SA
	Hybrid disc medic (<i>Medicago tornata</i> x <i>M. littoralis</i>)	W	1	✓	✓	x	1	✓	✓	✓	✓	✓	SA
	Serradella (<i>Ornithopus spp.</i>)	W	1	x	✓	✓	1	x	x	x	x	✓	SA
Fallow	Common vetch (<i>Vicia sativa</i>)	W	1	✓	✓	✓	1	✓	✓	✓	✓	x	A
	Cowpea (<i>Vigna unguiculata</i>)	S	1	✓	✓	x	1	✓	✓	✓	✓	✓	A
	Lablab (<i>Lablab purpureus</i>)	S	1	✓	✓	x	1	✓	✓	✓	✓	x	A
	Purple Vetch (<i>Vicia benghalensis</i>)	W	2	✓	✓	x	2	✓	✓	✓	✓	x	A

Table key: See below Table 24

Table 23. Grasses for the Northern grain belt

Minimum rainfall	Grass species	Tolerances						Soil preferences					Pasture duration
		Growing season	Water logging	pH		Acid	Salt	Black earth	Heavy clay	Friable clay	Loam	Sand	
				Alkaline	Medium								
650 + mm	Cocksfoot (<i>Dactylis glomerata</i>)	W	2	x	✓	✓	1	✓	x	✓	✓	✓	P
	Phalaris (<i>Phalaris aquatica</i>)	W	4	✓	✓	x	2	✓	✓	✓	✓	x	P
600 + mm	Tall festuca (<i>Festuca arundinacea</i>)	W	3	✓	✓	✓	3	x	x	✓	✓	✓	P
	Angelton grass (<i>Dichanthium aristatum</i>)	S	4	✓	✓	x	2	✓	✓	✓	✓	x	P
	Creeping blue (<i>Bothriochloa insculpta</i>)	S	2	✓	✓	✓	2	x	x	✓	✓	x	P
550 + mm	Gatton panic (<i>Panicum maximum</i>)	S	1	✓	✓	✓	2	x	x	✓	✓	x	P
	Green panic (<i>Panicum maximum</i>)	S	1	✓	✓	x	2	x	x	✓	✓	x	P
	Queensland blue (<i>Dichanthium sericeum</i>)	S	2	✓	✓	x	1	✓	✓	✓	✓	x	P
	Rhodes grass (<i>Chloris gayana</i>)	S	1	✓	✓	✓	3	✓	x	✓	✓	✓	P
	Silk sorghum (<i>Sorghum spp.</i>)	S	2	✓	✓	x	1	✓	✓	✓	✓	x	P
	Bambatsi panic (<i>Panicum coloratum</i>)	S	4	✓	✓	x	3	✓	✓	✓	x	x	P
	Buffel grass (<i>Cenchrus ciliaris</i>)	S	1	✓	✓	✓	2	x	x	✓	✓	✓	P
450 + mm	Digit grass (<i>Digitaria eriantha</i>)	S	2	✓	✓	✓	2	x	x	✓	✓	✓	P
	Purple pigeon (<i>Setaria incrassata</i>)	S	2	✓	✓	x	1	✓	✓	✓	✓	x	P
	Tall finger grass (<i>Digitaria milanijana</i>)	S	2	✓	✓	✓	2	x	x	✓	✓	✓	P
	Forage sorghum (<i>Sorghum bicolor</i>)	S	1	✓	✓	x	1	✓	✓	✓	✓	✓	A
Fallow	Oats (<i>Avena sativa</i>)	W	2	✓	✓	✓	1	✓	✓	✓	✓	✓	A

Source: Lloyd et al (2007a) LeyGrain – Pastures in Farming Systems. Sustaining Profit and the Environment

Key:

Season S: Summer W: Winter
Tolerance 1: Poor 2: Fair 3: Good 4: Very good
Pasture duration A: Annual P: Perennial SA: Semi annual SP: Semi perennial

Soil x Vegetation: Black earth: Basaltic plains soils
 Heavy clay: Coarse brigalow, Mitchell grass downs and Coolibah soils
 Friable clay: Brigalow vine scrub, Brigalow/Belah and Basaltic uplands soils
 Loam: Belah and Poplar box/Sandalwood soils
 Sand: Cypress pine/Bullock soils

Table 24. Summary of adaptation of grass species to soil/vegetation units in south Queensland relevant to cropping areas.

Location	Soil/vegetation	Soil features	Pasture adaptation	
			Good	Average
Maranoa	Mitchell grass cracking clay	Deep alkaline friable	PD, QB	Bam, PP, R
	Fringes of above	Deep alkaline friable	CB(h), PP, PD, CB(b),	Bam, Buff
	Red poplar box	Acid, duplex loam	IB	CB(b), PD
Western Downs	Grey brigalow (no melonholes)	Deep alkaline friable	PD, R, CB(h), IB, CB(b), PP	GP, Buff
	Grey brigalow (melonholes)	Deep alkaline heavy	CB(b), PP,	R, Bam
Eastern and south-eastern Downs	Upland, heavy cracking clay, black earth	Heavy, deep alkaline	CB(b), CB(h), Bam, QB, PD	R, PP

PD – Premier digit

Bam – Bambatsi

IB – Indian blue

CB(b) – Creeping blue (Bisset)

CB(h) – Creeping blue (Hatch)

R – Rhodes (Callide)

QB- Queensland blue

PP – Purple pigeon (Inverell)

Buff – Buffel (Biloela)

GP – Petrie green panic

Source: Lloyd *et al.* (2007b).

Table 25. Nitrogen contributions from grain legumes in Queensland and NSW, expressed in terms of nitrogen fertiliser equivalents (kgN/ha) or increased cereal yield (%).

Legume	Control crop	N benefit in fertiliser equivalent (kgN/ha) #	Yield increase in subsequent cereal (%)
Marcellos (1984) ¹			
Chickpea	wheat	50+	103
Faba bean	wheat	50+	87
Strong <i>et al.</i> (1986) ¹			
Chickpea	wheat	50+	24
Faba bean	wheat	50+	17
Field pea	wheat	50+	31
Dalal (1991) ¹			
Chickpea 1987	wheat	75	50
Chickpea 1988	wheat	75+	39
Chickpea 1989	wheat	75+	61
Holford (1993) ¹			
Chickpea	wheat	35	77
Doughton, Vallis and Saffigna (unpub) ¹			
Chickpea	sorghum	100	53
Doughton and Mackenzie (1984) ¹			
Mungbean (black)	sorghum	68	79
Mungbean (green)	sorghum	68	61
Cox <i>et al.</i> (1998) ²			
Chickpea 1996	wheat	0	0
Chickpea 1997	wheat	20	7
Chickpea 1998	wheat	35	23

From: 1. Doughton J and Holford I (1997)

2. Cox *et al.* (1998)

where a + is present, N benefit exceeded that of the highest rate of nitrogen fertiliser used

Pulse (legume) crops

The major pulse crops grown in south Queensland are chickpea (*Cicer arietinum*), mungbean (*Vigna radiata*) and faba bean (*Vicia faba*). Pulse crops may contribute to available nitrogen and also provide a break in the rotation to reduce diseases of cereal crops. Pulse grains are also profitable in their own right when prices are favourable. Chickpea in particular is well suited to deep sowing because of its ability to emerge from depth. A disadvantage of pulses is low stubble cover remaining after harvest.

Nitrogen contribution from pulse crops

Trials from Queensland and northern NSW have quantified the nitrogen contribution of pulse crops. The trials results will reflect the growing conditions of that year and hence the proportion of vegetative and grain material returned or removed as a result of climatic conditions and the effectiveness of weed and disease control. Measuring the nitrogen contribution directly is difficult and subject to errors. The option used in Table 25 is the expression as 'fertiliser equivalent' and subsequent cereal yield increase.

Removal of grain from a legume (pulse) crop is a loss of nitrogen from the paddock. The net nitrogen balance following a pulse crop can vary widely. For example Schwenke *et al.* (1998) measured a range of net nitrogen balance from -47 to + 46 kgN/ha and -12 to 94 kgN/ha following chickpea and faba bean respectively.

Livestock production from pasture legumes

Detailed budgeting and planning capability is provided in the LeyGrain Manual (Lloyd *et al.* 2007a) which has a comprehensive section for estimating livestock production, days of grazing, stock and paddock area requirements and catering for feed short-falls.

Feed budgeting is complex with feed demand depending on the age, size and class of livestock constituting the herd. However, Table 26 provides the broad carrying capacities and weight gains from forages commonly used for beef production and may be used to develop a 'rough guide' to livestock production.

For each paddock, determine:

- area of each paddock
- the carrying capacity
- expected live weight gain from each pasture (paddock)
- number of grazing days
- number and liveweight of animals in and out.
- Costs will include: the purchase price of cattle (\$/kg) and other variable costs, (planting, veterinary, freight, etc)

Return = liveweight gain x beef price/kg

Table 26. Broad stocking periods, carrying capacities, and live weight gains from a number of forages.

Forage	Feeding period	Cattle carrying capacity (ha/AE*)	Daily gain kg/head
Native pasture	October – May + dry feed in winter	2.0 – 4.0	0.2 – 0.6
Sown tropical grass	October – May + dry feed in winter	1.5 – 2.5	0.4 – 0.8
Lucerne	September – May	1.5 – 2.2	0.8 – 1.2
Sown medics	May – October	1.2 – 2.0	0.6 – 1.0
Subclover	May – November	1.5 – 2.0	0.7 – 1.2
Lab lab	January – May	0.5 – 1.5	0.7 – 1.0
Butterfly pea	December – April	1.0 – 2.0	0.8 – 1.0
Oats	May – October	0.5 – 1.0	0.8 – 1.2
Barley	May – September	0.6 – 1.2	0.8 – 1.0
Forage sorghum	December – May	0.4 – 1.0	0.5 – 1.0
Hybrid millet	December – May	0.6 – 1.0	0.6 – 1.0
Barley stubble	October – Feb	2.0 – 4.0	0.1 – 0.4
Sorghum stubble	February – August	1.5 – 3.0	0.1 – 0.5

*AE = Adult Equivalent = 450 kg dry cattle; Note: 1 AE = 8 DSE (Dry Sheep Equivalents)

Source: (Lloyd *et al.* 2007a) LeyGrain Manual.

References

Cox HW, Strong WM, Lack DW and Kelly RM. (1998) Profitable double-crop rotations for central Queensland. In *Agronomy – Growing a greener Future*. Edited by D.L Michalk and J.E Pratley. *Proceedings of the 9th Australian Agronomy Conference*, 20-23 July 1998, Charles Sturt University, Wagga Wagga.

Dalal RC, Weston EJ, Strong WM, Lehane KJ, Cooper JE, Wildermuth GB, King AJ and Holmes CJ (2004) Sustaining productivity of a Vertosol at Warra, Queensland, with fertilisers, no-tillage or legumes. 7. Yield, nitrogen and disease-break benefits from lucerne in a two-year lucerne-wheat rotation. *Australian Journal of Experimental Agriculture*, **44**, 607-616.

Doughton J and Holford I (1997) Legumes. In: *Sustainable Crop Production* Ed: AL Clarke and PB Wylie. Queensland Department of Primary Industries & Fisheries.

Doughton JA and McKenzie J (1984). Comparative effects of black and green gram (mungbeans) and grain sorghum on soil mineral nitrogen and subsequent grain sorghum yields on the eastern Darling Downs. *Australian Journal of Experimental Agriculture and Animal Husbandry*. **24**, 244-9.

Lloyd DL, Johnston B, O'Brien S, Roesner, L, Boschma, S, Williams, R (2007a). Benefits of pastures to crops, soil and the environment. In: *Leygrain; Pastures in Farming Systems*, DPI&F Queensland and NSW DPI.

Lloyd DL, Johnston B and O'Brien S (2007b). Which pastures and should they be mixed? In: *Pastures for Production and Protection*, Proceedings 7th Australian Tropical Pastures Conference, Dalby.

Schwenke GD, Peoples M, Turner GL and Herridge DF. (1998). Does nitrogen fixation of commercial, dryland chickpea and faba bean crops in north-west New South Wales maintain or enhance soil nitrogen? *Australian Journal of Experimental Agriculture*, 1998, 38, 61-70.

Strong WM, Dalal RC, Weston EJ, Lehane KJ, Cooper JE, King AJ and Holmes, CJ (2006) Sustaining productivity of a Vertosol at Warra, Queensland, with fertilisers, no-tillage or legumes. 9. Production and nitrogen benefits from mixed grass and legume pastures in rotation with wheat. *Australian Journal of Experimental Agriculture*, **46**, 375-385.

Weston EJ, Dalal RC, Strong WM, Lehane KJ, Cooper JE, King AJ and Holmes CJ (2002) Sustaining productivity of a Vertosol at Warra, Queensland, with fertilisers, no-tillage or legumes. 6. Production and nitrogen benefits from annual medic in rotation with wheat. *Australian Journal of Experimental Agriculture*, **42**, 961-969.

Glossary of terms

Autotrophs	Organisms that depend on either light or on oxidation of inorganic or elemental substances for energy and CO ₂ as the sole source of carbon
Ammonification	The biochemical process of the conversion of organic compounds to ammonia compounds
Denitrification	The biochemical reduction of nitrate and nitrite to gaseous nitrogen; N ₂ O and N ₂
Heterotrophs	Organisms that require an organic source of carbon for energy and growth
Immobilisation	Conversion of an element from an inorganic form to an organic form, thus rendering the element less available
Mineralisation	Conversion of an element from an organic form to an inorganic form, thus rendering the element more available for plant uptake
N fixation	The biological conversion of elemental nitrogen (N ₂) to organic forms readily usable in the biological process
Nitrification	The biochemical oxidation of ammonium to nitrate predominantly by autotrophic bacteria
Symbiotic	two dissimilar organisms living in association for mutual benefit
Rhizobium	Bacteria living symbiotically with plants, usually in root nodules of legumes. They receive energy from the plant whilst converting atmospheric nitrogen to organic forms
¹⁵ N	A naturally occurring isotope of nitrogen
Rhizosphere	The soil surrounding the root of a plant in which the abundance and composition of microbial population is influenced by the roots

More information

For more information or to get a copy of the *WhopperCopper* software tool, contact the Department of Agriculture and Fisheries on 13 25 23 or visit www.daf.qld.gov.au

