



Celebrating the
100th Volume



A review of strawberry under protected cultivation: yields are higher under tunnels than in the open field

Christopher Michael Menzel

To cite this article: Christopher Michael Menzel (2025) A review of strawberry under protected cultivation: yields are higher under tunnels than in the open field, The Journal of Horticultural Science and Biotechnology, 100:3, 286-313, DOI: [10.1080/14620316.2024.2449026](https://doi.org/10.1080/14620316.2024.2449026)

To link to this article: <https://doi.org/10.1080/14620316.2024.2449026>



View supplementary material [↗](#)



Published online: 05 Feb 2025.



Submit your article to this journal [↗](#)



Article views: 169



View related articles [↗](#)



View Crossmark data [↗](#)

REVIEW



A review of strawberry under protected cultivation: yields are higher under tunnels than in the open field

Christopher Michael Menzel

Department of Agriculture and Fisheries, Nambour, Queensland, Australia

ABSTRACT

Strawberries are grown under tunnels to protect the plants from cold, frosts, rain, and fruit diseases. A review was conducted to determine the performance of plants under plastic tunnels. Information was collected on yields and fruit weight under tunnels and in the open field ($n = 133$ experiments) and on environmental conditions in the two areas. In a global analysis, plants under tunnels had higher relative marketable (Tunnel/Open = 1.34 ± 0.76) and total yields (Tunnel/Open = 1.30 ± 0.83) than those in the open ($p < 0.001$). In contrast, fruit weight was similar in the two growing areas (Tunnel/Open = 1.04 ± 0.22) ($p = 0.094$). Relative marketable yields (Tunnel/Open) were similar in plants in Northern and Southern Europe, and in Northern and Southern America ($p > 0.05$). Relative marketable yields were similar in areas with cool or cold winters or with spring/summer or winter/spring production seasons, and under low or high tunnels ($p > 0.05$). Lower yields under tunnels were associated with low light levels and high temperatures under the plastic and a higher incidence of powdery mildew. The use of tunnels under global warming will require attention to ventilation under the covers.

ARTICLE HISTORY

Received 20 November 2024
Accepted 29 December 2024

KEYWORDS

Fruit weight; light; profitability; temperature; yield

Introduction

Strawberries (*Fragaria × ananassa* Duch.) are the most common fruit across the berry catalogue and account for 72% of berry production worldwide (Symington & Glover, 2024; Xu et al., 2023). Major producers include China (36.2% of global production), the United States, Mexico, and several countries throughout Europe. Total production is about 10 million tonnes worth US \$22 billion from about 400,000 ha (FAOSTAT, n.d.). Average yields have remained stable over the last decade and ranged from 21.0 to 22.4 t/ha. Strawberry growing in Spain is concentrated in the southwestern region of the country in Huelva Province (de Los Santos et al., 2021). This area makes up 93% of the total Spanish production, with 6,867 ha, 377,596 tons and a market value of €392 million in 2016. Huelva is the leader in European strawberry production, producing 25% to 33% of total European production and is the world's largest exporter. Average yields under tunnels in Spain are twice those achieved in the United Kingdom (Webb et al., 2013).

Strawberry plants are adapted to a range of environments, with production in areas with a cool temperate climate (e.g. Latvia, Norway, and Canada), a warm, subtropical climate (e.g. Florida, Queensland, and Argentina), or a Mediterranean climate (e.g. California, Spain, and Türkiye) (Ahmadi et al., 2024; Amyotte & Samtani, 2023; Gullickson et al., 2024; Holmes, 2024; Kirschbaum et al., 2024; Ramos et al.,

2024; Zheng et al., 2024a; Tabatabaie & Murthy, 2016). In the tropics, the plants are grown at elevation (e.g. Arif et al., 2019; and Suminarti et al., 2023 in Indonesia). Different cultivars are grown in the various locations, with plant breeding improving yield, disease resistance, and fruit quality (Knapp et al., 2023; Porter et al., 2023; Alam et al., 2024; Fan et al., 2024; Feldmann et al., 2024; Lima et al., 2024; Prohaska et al., 2024a; Prohaska et al., 2024b).

The plants are susceptible to extreme temperatures, with frosts and heat-waves damaging the flowers, fruit, and leaves, and decreasing yields (Cordeiro & Dötterl, 2023; Hernández-Martínez et al., 2023; Johnson & Hoffmann, 2024; López et al., 2024; Luo et al., 2024; Martínez-Ferri et al., 2014; Matsui & Mochida, 2024; Montague et al., 2024; Nestby et al., 2001; Pedrozo et al., 2024; Silva et al., 2024; Ullah et al., 2024; Unnikrishnan et al., 2024). Prolonged periods of cool weather shorten the harvest period in several areas where there are low temperatures in autumn or spring. High temperatures during the peak of summer inhibit flowering and fruit set.

Production is affected by heavy or prolonged rain. The fruit are damaged directly by rain, with water-soaking and cracking of the skin. The fruit have a high water content and a thin peel or skin (Ma et al., 2024; Shanthini et al., 2025). Nearly all cultivars are susceptible when the fruit are close to harvesting (Herrington et al., 2011; Hurtado & Knoche, 2021; Hurtado et al.,

2024; Straube et al., 2024). The incidence of several fruit diseases such as grey mould (*Botrytis cinerea*), anthracnose fruit rot (*Colletotrichum acutatum* and related species), and powdery mildew (*Podosphaera aphanis*) is promoted by direct rain contact or by high humidity (Aldrighetti & Pertot, 2023; Filippi et al., 2021; Gama et al., 2023; Lynn et al., 2024).

Crops, including strawberry, are grown under protected cultivation (protected cropping or protected agriculture) to protect the plants, flowers, and fruit from extreme weather, including heavy rain and frosts (Maier et al., 2022; Verteramo Chiu et al., 2024). This article reviews the performance of strawberry under protected cultivation. Information was collected on total yield, marketable yield, and fruit weight from experiments where the plants were grown under low or high tunnels and in the open field. Additional information was collected on environmental conditions in the two systems, the incidence of the major fruit diseases affecting strawberry and on profitability of the plants. Finally, the role of protected cultivation in minimising the impacts of global warming on strawberry production is discussed.

Overview of horticultural production under protected cultivation

Many crops are grown under protected cultivation around the world, with the importance of the technology varying with the species, growing area, climate, and access to capital (Jiang & Yu, 2008 in China; Bertuglia & Calatrava, 2012 in Spain; Sabir & Singh, 2013 across the globe; Meland et al., 2017 in Norway; Asseng et al., 2020 in the United States; Gu, 2021 in the United States; Jiménez-Lao et al., 2021 in South America; Yamanaka & Kawashima, 2021 in Japan; Lubna et al., 2022 across the globe; Morel & Cartau, 2023 in France; Argento et al., 2024 in the Mediterranean area; Azad et al., 2024 in the Desert Southwest of the United States; Cusworth et al., 2024a in the United Kingdom; Cusworth et al., 2024b in China; Kaiser et al., 2024 across the globe; Kim, 2024 in Korea; Lakhiar et al., 2024 across the globe; Ranasingha et al., 2024 in the United Kingdom; Rantanen et al., 2024 in Finland; Stone et al., 2024 in Australia; and Sturiale et al., 2024 in Italy, Spain, Tunisia, and Türkiye). In Southern Spain, intensive horticulture contributes 40% of the gross domestic product in some regions (Egea & Glass, 2017; Egea et al., 2018; Talavera et al., 2024). There are 40,000 ha of plantings under plastic greenhouses or macro-tunnels. These structures produce more than 4 million tons of vegetable and berry crops that are shipped to the rest of Europe and valued at €3 billion each year.

Low-technology polytunnels contribute to 80% to 90% of the area under protected cultivation globally, with glasshouse production less important (Jayasuriya

et al., 2024). The area of the protected horticulture was 2.67 million hectares in China in 2021, accounting for more than 80% of the global protected horticulture area (Dong et al., 2024). The area under protected cultivation in the United States is about 0.3% to 1.0% of the global area (Janke et al., 2017). Cultivation under plastic is expanding in many locations within Europe, with the largest area in Spain and Italy (Scarascia-Mugnozza et al., 2011). There are 13,932 ha under protected cultivation in Australia, with 64.8% under nets and 30.0% under poly-houses and poly-tunnels (Clark et al., 2023). Glasshouses and shade houses are less important (5.2%). The total area of horticultural crops under protected cultivation in India is 30,000 to 40,000 ha (Sharma et al., 2024). Protected cultivation of fruit is at an early stage, with strawberry the only fruit crop grown commercially in greenhouses.

Rantanen et al. (2024) examined the benefits of protected cultivation for raspberry in Finland. About 70% of the crop in this area is produced under tunnels to reduce the impacts of cold and rain. Yields were up to ten times higher under tunnels than in the open field. The main environmental impacts from tunnel production came from the fertilisers and the substrates, the tunnel structures, and other materials, all explaining two-thirds of the impacts on climate change gas emissions. In contrast, the main environmental impacts from field production came from the growing activities, which explained over half of the climate change impacts. Eutrophication and land use were 60% and 70% lower with the tunnels than with field production. It was concluded that higher yields and more efficient production can reduce the environmental impacts of tunnel production.

Overview of strawberry production under tunnels in different areas

Strawberries are grown under protected cultivation using plastic covers to protect the plants from cold weather, frosts, and rain (Akpenpuun et al., 2021; Cayambe et al., 2023; Chavan et al., 2022; Clark & Mousavi-Avval, 2022; Domínguez et al., 2016; Osman et al., 2024; Santos et al., 2013; Soria et al., 2009). There are two main structures used to protect the crops from extreme weather, including low and high tunnels. Low tunnels include row covers and small portable tunnels that just cover the growing crop. High tunnels are generally permanent, although sometimes the plastic can be stored between seasons. In the United Kingdom, many of the poly-tunnels have disposable plastic covers fitted in early spring and removed in late autumn and replaced on an annual basis, although new poly-tunnel designs are moving to multi-year covers to reduce the costs of installation and removal of the plastic (Sakrabani et



Figure 1. Production of commercial strawberries under high tunnels in the United Kingdom. From top to bottom: planting without tunnels; extensive tunnel production; tunnel structures before the plastic covers are employed; close-up of tunnels; flower production on benches under tunnels; and heavy fruit production on benches under tunnels.

al., 2023; Figure 1). High tunnels allow the crop to be harvested without removing the plastic. The tunnels generally lack heating and cooling, thereby providing some of the benefits of a greenhouse at a lower cost (Argento et al., 2024; Giacomelli, 2009; Gupta et al., 2024; Nemali, 2022; Pierre et al., 2024).

There are various methods to control the temperature under high tunnels, including passive and active venting (Tian et al., 2023). The sides of the tunnels can be rolled up during warm weather to prevent excessive temperatures inhibiting plant growth and flowering. Temperatures during the day can be excessive for plant growth without venting. Temperatures during the night decrease without some method of heating (Grisey et al., 2023; Zhang et al., 2023). Most tunnels are not heated.

A range of plastics can be used to cover the plants, with different light transmission properties (Lewers et al., 2020; Ordidge et al., 2012; Pandey et al., 2023; Romero-Gómez et al., 2012). The transmission of light through the covers decreases slightly over time

as the plastic ages (Uchanski et al., 2020). A major limitation on the employment of plastic covers in tunnels is their disposal after use, which leads to pollution of soil and water environments (Divya & Sarkar, 2019; Galafton et al., 2023; Pergola et al., 2023; Gupta et al., 2024; Logan et al., 2024a; Logan et al., 2024b). Numerous photo- and bio-degradable films have been developed to reduce the impacts of plastics on the environment (Kasirajan & Ngouajio, 2012). However, these perishable films often degrade too quickly or incompletely in the air and soil. Biodegradable films for tunnels have a lifetime of 6 months, much shorter than standard films (Kapanen et al., 2008). Balocco et al. (2018) indicated that low-density polyethylene (LD-PE) degraded slowly, with new and 5-year-old plastic having similar light-transmissions.

The use of protected cultivation for strawberry varies across growing areas. Protected cultivation is important in the United Kingdom, Spain, and some other areas in Europe (Fountain et al., 2023;

Khoshnevisan et al., 2013; Kulak et al., 2013; Martínez et al., 2017; Martínez-Ferri et al., 2016; Neri et al., 2012; Nestby & Guéry, 2017; Port & Scopes, 1981; Roca et al., 1998; Romero-Gómez & Suárez-Rey, 2020; Savvas et al., 2016; Soode Schimonsky et al., 2017; Verteramo Chiu et al., 2024). In Northern America, protected cultivation is used in Canada and parts of the United States where there is the risks of frosts or snow (e.g. Arkansas, Kansas, Minnesota, New York, Texas, and Utah) (Garcia et al., 2017; Hodgdon et al., 2024; Maughan et al., 2015; Scott et al., 2021). Most of the crop in California is grown in the open. Overall, less than 10% of the crop in the United States is grown under tunnels.

Protected cultivation is well developed in Europe, particularly in the United Kingdom and Spain. Poly-tunnels account for more than 80% of the area in the United Kingdom, including Scotland (Beech & Simpson, 1989; Carter et al., 1993; Warner et al., 2010). Protected cultivation and other strategies have been associated with increases in the yields of strawberry over the past two to three decades in the United Kingdom, with yields increasing from 9.9 t/ha in 1996 to 2000 to 22.3 t/ha in 2011 to 2015 (Cusworth et al., 2022). Ninety-seven percent of the area in Spain is under protected cultivation and three percent is in the open (Evans, 2013; García-Tejero et al., 2018; Romero-Gómez & Suárez-Rey, 2020). Eighty-two percent of the area is under macro-tunnel and eighteen percent is under micro-tunnel. Ninety-six percent of the plants are planted in soil, with a mixture of fumigated and non-fumigated fields across the industry. An increase in yield under tunnels in Spain of 20 g/plant was associated with an increase in income of €1,300/ha (Guéry et al., 2018). In the United Kingdom, 55% of production takes place in substrate or soil-less cultivation with fertigation provided by drippers, while 45% is in the soil (British Summer Fruits, 2017). There are efforts to re-use peat in the soil-less systems to reduce the impacts on the environment (Gruda et al., 2024; Vandecasteele et al., 2024).

In the United Kingdom, strawberries are produced from March to December (Raffle et al., 2010). The earliest crops are harvested under glass in Southern England in early April, although additional heat and night-break lighting bring crops as early as March. The season continues under closed-fixed plastic

tunnels, followed by field-grown crops under French and Spanish tunnels. The traditionally grown, unprotected field crops are available in June and July. Hancock and Simpson (1995) provided a similar analysis for Europe and identified seven areas with different climates and production systems (Table 1).

Verteramo Chiu et al. (2024) reviewed the performance of strawberry in the open, under un-heated tunnels (plasticulture) and in heated greenhouses. Mean yields (\pm standard deviation or s.d.) were higher under the tunnels (33.5 ± 16.9 t/ha) and in the greenhouses (47.7 ± 26.7 t/ha) than in the open (18.7 ± 16.9 t/ha) ($p < 0.01$). Information was collected from forty-four tunnel experiments, eight greenhouse experiments and twenty-seven open field experiments. Some of the differences in productivity across the growing systems could be due to variations in plant densities, with plants in greenhouses grown at higher densities than in the other systems.

Performance of strawberry under tunnels around the globe

Productivity and fruit weight varied across the regions, climatic zones, production seasons, and types of tunnels (Supplementary files S1 and S2; Tables 2 to 4). Mean (\pm s.d.) marketable yield was 422 ± 351 g/plant under the tunnels and 364 ± 321 g/plant in the open. Mean total yields were 403 ± 282 g/plant and 367 ± 299 g/plant, and mean fruit weights were 13.9 ± 4.8 g and 13.7 ± 5.2 g.

In the global analysis, plants under tunnels had higher relative marketable (Tunnel/Open = 1.34 ± 0.76) and total yields (Tunnel/Open = 1.30 ± 0.83) than those in the open ($p < 0.001$). In contrast, fruit weight was similar in the two systems (Tunnel/Open = 1.04 ± 0.22) ($p = 0.094$). Variations in relative yields and fruit weight across the globe are shown in Figure 2.

Relative marketable yield (Tunnel/Open) was similar in Northern (1.13 ± 0.44) and Southern Europe (0.93 ± 0.38), and in Northern (1.42 ± 0.89) and Southern America (1.29 ± 0.29) ($p > 0.05$). Relative total yield was similar in Northern (0.99 ± 0.31) and Southern Europe (1.18 ± 0.73), and in Northern (1.39 ± 1.03) and Southern America (1.44 ± 0.39) ($p > 0.05$). Relative fruit weight was similar in Northern (1.07 ± 0.46) and Southern Europe (0.99 ± 0.20), and in

Table 1. Characteristics of strawberry-growing areas in Europe. Data are from Hancock and Simpson (1995).

Region	Climate	Production system
Spain & Southern Italy	Mild Mediterranean	Tunnels & open
Northern Italy & South-eastern France	Mild, occasionally cold winters	Tunnels & open
Germany	Continental, cold winters	Open & tunnels
The Netherlands & Belgium	Continental, cold winters	Open, greenhouse & tunnels
United Kingdom & the Republic of Ireland	Mild, maritime	Tunnels & open
Poland	Continental, severe winters	Mostly open, with some tunnels
Scandinavia	Short growing season, severe winters	Mostly open, with some tunnels

Table 2. Marketable yield of strawberries grown in the open field and under tunnels across different geographical zones, climatic zones, production seasons, and type of tunnel (low or high tunnel). Data are from Supplementary files S1 and S2. There were two studies in the Asia-Pacific region and one in Africa. s.d. = standard deviation.

	Open field		Tunnel		<i>n</i>
	Mean yield (g/plant) \pm s.d.	Median yield (g/plant)	Mean yield (g/plant) \pm s.d.	Median yield (g/plant)	
Global data set	364 \pm 321	317	422 \pm 351	362	66
<i>Geographical zone</i>					
Northern Europe	221 \pm 154	200	241 \pm 186	200	18
Southern Europe	467 \pm 230	538	401 \pm 223	397	21
Northern America	339 \pm 184	322	408 \pm 220	371	57
Southern America	752 \pm 826	500	866 \pm 837	534	11
Asia-Middle East	327 \pm 360	327	339 \pm 357	339	23
<i>Climatic zone</i>					
Cool winter	315 \pm 179	312	371 \pm 215	357	69
Cold winter	436 \pm 450	324	496 \pm 481	367	64
<i>Production season</i>					
Winter/spring	423 \pm 463	290	485 \pm 493	334	67
Spring/summer	329 \pm 190	321	384 \pm 225	367	66
<i>Type of tunnel</i>					
Low tunnel	325 \pm 169	314	395 \pm 291	391	58
High tunnel	390 \pm 390	317	440 \pm 417	346	75

Table 3. Total yield of strawberries grown in the open field and under tunnels across different geographical zones, climatic zones, production seasons, and type of tunnel (low or high tunnel). Data are from Supplementary files S1 and S2. There were two studies in the Asia-Pacific region and one in Africa. s.d. = standard deviation.

	Open field		Tunnel		<i>n</i>
	Mean yield (g/plant) \pm s.d.	Median yield (g/plant)	Mean yield (g/plant) \pm s.d.	Median yield (g/plant)	
Global data set	367 \pm 299	307	403 \pm 282	343	97
<i>Geographical zone</i>					
Northern Europe	283 \pm 192	256	263 \pm 209	175	15
Southern Europe	505 \pm 244	494	522 \pm 247	491	16
Northern America	403 \pm 239	389	472 \pm 280	429	36
Southern America	384 \pm 301	339	512 \pm 357	500	6
Asia-Middle East	278 \pm 436	159	292 \pm 272	202	23
<i>Climatic zone</i>					
Cool winter	394 \pm 354	310	428 \pm 290	357	52
Cold winter	335 \pm 219	307	375 \pm 274	301	45
<i>Production season</i>					
Winter/spring	381 \pm 344	301	412 \pm 271	370	51
Spring/summer	351 \pm 242	313	393 \pm 296	302	46
<i>Type of tunnel</i>					
Low tunnel	365 \pm 335	319	404 \pm 286	342	46
High tunnel	369 \pm 265	285	403 \pm 281	343	51

Table 4. Fruit weight of strawberries grown in the open field and under tunnels across different geographical zones, climatic zones, production seasons, and type of tunnel (low or high tunnel). Data are from Supplementary files S1 and S2. There were two studies in the Asia-Pacific region and one in Africa. s.d. = standard deviation.

	Open field		Tunnel		<i>n</i>
	Mean fruit weight (g) \pm s.d.	Median fruit weight (g)	Mean fruit weight (g) \pm s.d.	Median fruit weight (g)	
Global data set	13.7 \pm 5.2	12.8	13.9 \pm 4.8	13.7	73
<i>Geographical zone</i>					
Northern Europe	12.5 \pm 3.5	11.8	12.6 \pm 3.9	12.7	8
Southern Europe	13.7 \pm 5.5	11.4	13.2 \pm 4.7	11.7	12
Northern America	14.6 \pm 5.1	14.0	14.6 \pm 4.8	14.0	29
Southern America	13.8 \pm 5.5	11.7	15.0 \pm 5.1	13.8	7
Asia-Middle East	12.6 \pm 5.6	11.6	13.2 \pm 4.7	12.8	15
<i>Climatic zone</i>					
Cool winter	14.9 \pm 6.4	14.0	14.8 \pm 5.6	14.9	39
Cold winter	12.4 \pm 3.1	12.6	12.9 \pm 3.4	13.1	34
<i>Production season</i>					
Winter/spring	14.5 \pm 6.0	13.9	14.4 \pm 5.2	14.4	38
Spring/summer	12.9 \pm 4.1	12.6	13.4 \pm 4.3	13.2	35
<i>Type of tunnel</i>					
Low tunnel	13.0 \pm 4.1	13.6	13.4 \pm 4.0	13.2	33
High tunnel	14.3 \pm 6.0	12.2	14.4 \pm 5.3	13.9	40




Figure 2. Box plots showing the distribution of relative yield (marketable and total yield in g/plant) and relative fruit weight (g) in strawberry plants grown under tunnels and in the open field (Tunnel/Open). Data are from Supplementary files S1 and S2.

Northern (1.01 ± 0.14) and Southern America (1.12 ± 0.21) ($p > 0.05$). These results suggest that the plants responded in the same way to protected cultivation in the different areas.

Marketable yields were higher in areas with cold winters than those with cool winters and higher under tunnels than in the open in both climates (Table 2). Relative marketable and total yield and fruit weight (Tunnel/Open) were similar in areas with cool (1.32 ± 0.63 , 1.38 ± 0.89 , and 1.03 ± 0.19) or cold winters (1.34 ± 0.84 , 1.22 ± 0.76 , and 1.06 ± 0.26) ($p > 0.05$). These results suggest that the behaviour of the plants under protected cultivation was similar in the different climates.

Marketable production was affected by the cropping season and cultivation (Table 2). Yields were high in areas with harvests over winter and spring than harvests over spring and summer. Yields were also high in under tunnels than in the open. Relative marketable and total yield and fruit weight (Tunnel/Open) were similar with fruit produced in winter and spring (1.35 ± 0.65 , 1.38 ± 0.90 , and 1.03 ± 0.19) or spring and summer (1.33 ± 0.82 , 1.22 ± 0.75 , and 1.06 ± 1.15) ($p > 0.05$).

Average marketable yields were higher under high (440 ± 417 g/plant) than under low tunnels (395 ± 291 g/plant) (Table 2). Relative marketable and total yields and fruit weight (Tunnel/Open) were similar under the low (1.28 ± 0.46 , 1.35 ± 0.83 , and 1.05 ± 0.16) or high tunnels (1.37 ± 0.90 , 1.27 ± 0.84 , and 1.04 ± 0.26) ($p > 0.05$). This indicates that the relative performance of the plants under the two types of tunnels were similar.

Plants under tunnels had higher total and marketable yields than those in the open. In contrast, fruit weight was similar in the two systems. The response to protected cultivation was similar in different geographic areas, locations with different climates or production seasons, and in the low or high tunnels.

Light levels under tunnels

Light levels under tunnels vary across the globe. The information available suggests that shading under the plastic covers can reduce the yields and quality of strawberry plants.

Total light receipts are lower under tunnels than in the open, although there is a higher proportion of diffuse light under the covers (Al-Helal et al., 2020; Al-Madani et al., 2024; Jayalath et al., 2017; Maynard & O'Donnell, 2019; Retamal-Salgado et al., 2015; Rho et al., 2020; Salvadores & Bastías, 2023). There are changes in the transmission of different wavelengths, including the transmission of ultraviolet (UV) light (Fletcher et al., 2004; Gude et al., 2022; Kotilainen et al., 2018; Matamala et al., 2023; Mishra et al., 2023). In experiments in Spain, the yields of ‘Camarosa’ and ‘Ventana’ were 20% to 30% higher under UV-light blocking plastic than under UV-light transmitting plastic (Casal et al., 2009). Ripening was delayed with the UV-light blocking film, while fruit weight was increased.

Shading under the covers can reduce plant growth and yields. The impact of the plastic on productivity varies with the crop, latitude, and the season. Low light levels are more of an issue at high latitudes than at low latitudes and over winter than over summer (Robson et al., 2022). Cloud, fog, and aerosols (small particles in the atmosphere) also reduce ambient light levels. Changes in the transmission of UV light effect the development of plant diseases (Barnes et al., 2023). Onofre et al. (2022) found that UV-transmitting plastics reduced the severity of powdery mildew in strawberry under tunnels in Florida. The severity of the disease was higher under all plastics than that in the open field.

He and Li (2014) examined the performance of strawberry under tunnels in Taiwan. Light levels under the tunnels were 30% lower than those outdoors. This response was associated with a 30% decrease in net CO_2 assimilation by the canopy compared with plants in the open, a 35% decrease in leaf area, and a 60% decrease in plant dry weight. The number of flowers per plant was 55% lower under the plastic. The results of this study demonstrate that shading under tunnels can reduce the growth and yield of strawberry.

Plastic covers alter total light receipts and the levels of ultraviolet light. Anderson et al. (2019) investigated

the performance of strawberry under tunnels and in the open in Minnesota. Two types of covers were used, including a UV-transmitting and a UV-blocking film. In the first year, the UV levels were $89.1 \pm 7.8 \mu\text{mol}/\text{m}^2/\text{s}$ in the open and 59.1 ± 5.2 and $4.8 \pm 0.3 \mu\text{mol}/\text{m}^2/\text{s}$ under the two types of covers ($p < 0.001$). Total solar irradiance was $32.7 \pm 1.1 \mu\text{mol}/\text{m}^2/\text{s}$, $22.0 \pm 0.9 \mu\text{mol}/\text{m}^2/\text{s}$, and $22.7 \pm 1.0 \mu\text{mol}/\text{m}^2/\text{s}$ in the open, UV-transmitting and UV-blocking treatments ($p < 0.001$). Total yields were higher under the UV-transmitting film ($669 \pm 25 \text{ g/plant}$) and lower in the open ($520 \pm 31 \text{ g/plant}$) and under the UV-blocking film ($570 \pm 37 \text{ g/plant}$) ($p = 0.0058$). Total light levels were adequate for high yields under the tunnels, with the UV-transmitting film beneficial for production. Mean maximum temperatures varied by less than 3°C across the different areas, suggesting adequate ventilation under the tunnels.

Condori et al. (2017) studied the relationship between yield and the environment in five strawberry cultivars under tunnels and in the open in Maryland. Average yields were 40% higher under the tunnels than in the open. Higher productivity under the tunnels was attributed to warmer conditions under the plastic and higher light-use efficiency. Average maximum temperatures were 3.5°C higher under the tunnels than in the open. Changes in temperature and solar radiation accounted for 41% of the variation in yield over the season. It was estimated that the optimum temperature for yield was 26.8°C . Higher temperatures and adequate light levels under the tunnels provided acceptable fruit production in this area. Nakayama et al. (2024) grew strawberry plants under controlled-environment conditions and reported that a photosynthetic photon flux (PPF) of $359 \mu\text{mol}/\text{m}^2/\text{s}$ for 10.5 h each day was sufficient for fruit production. Maximum values of PPF at noon on sunny days in many locations are $2,500 \mu\text{mol}/\text{m}^2/\text{s}$.

Dias et al. (2015) found that temperatures under tunnels were adequate for fruit production in strawberry in Brazil. Mean daily temperatures varied by less than 2°C under the tunnel and in the open and ranged from 26.0° to 28.0°C over the season. In contrast, the average solar radiation level under the tunnel was less than half of that in the open. Total yields were 32% lower under the tunnel (7.2 t/ha) than in the open (10.7 t/ha). Productivity was lower under the tunnel due to shading by the plastic. Henschel et al. (2024) investigated the effect of different films on the performance of two strawberry cultivars under tunnels in Brazil. ‘Albion’ had higher yields under transparent, opaque, or red covers or in plots without a cover (control) compared with plots with a blue cover. In contrast, the yields of ‘Camarosa’ were not affected by the covers. Overall, ‘Albion’ produced twice the yield of ‘Camarosa’.

Solar radiation levels under tunnels vary with the type of cover and season. Shading under the tunnels can be severe enough to reduce yields compared with plants in the full sun. The covers can reduce the amount of solar radiation transmitted through the plastic and mediate extreme temperatures under the tunnels.

Temperatures under tunnels

Strawberry plants are sensitive to temperature conditions during the growing season. Plastic covers can protect the plants from cold weather and frosts. On the other hand, temperatures under the tunnels can be excessive for growth if ventilation is inadequate.

Crops can be grown under tunnels to protect the plants from cold weather and extend the production season in areas prone to cold weather or frosts (Ogden & van Iersel, 2009 in Georgia; Bruce et al., 2019 in the Mid-West of the United States; Zheng et al., 2019 in Tennessee; Arancibia et al., 2023 in Missouri; Shaik et al., 2023 in Texas; and Gaisser et al., 2024 in Massachusetts). In some situations, temperatures can be too low under the tunnels for acceptable growth, while in other situations, temperatures can be excessive (Black, 2010).

Several factors influence conditions under the tunnels. These include the location of the tunnel and the weather, the type of plastic used to cover the plants, the height of the tunnel and ventilation within the tunnel (Guttormsen, 1972; Kapanen et al., 2008; Lang et al., 2016; Lewus & Both, 2020; Lozano et al., 2016; Savage, 1980; Tan et al., 1984; Villagrán et al., 2021; Wien & Pritts, 2009; Zhang et al., 2022). Heat stress for workers under plastic tunnels or greenhouses can be problematic if temperatures are above 35°C and relative humidities are excessive (Jung & Kim, 2022 in Korea; Simane et al., 2022 in Ethiopia; Tiwari et al., 2023 in India; and Isoyama et al., 2024 in Japan). Workers can also be at greater risks of exposure to pesticides under the plastic (Cerruto et al., 2018 in Italy; and Requena et al., 2019; Ruiz-González et al., 2024 and Zheng et al., 2024b in Spain), although some reports suggest that this risk is over-stated (see Hernández et al., 2020; Swaen, 2020).

Zheng et al. (2017) modelled the energy balance and airflow within a naturally ventilated tunnel in Tennessee. The tunnel was 15 m long and 9 m wide, with sliding doors at each end. The plastic reflected 20% of the incoming solar radiation, with 1% of the solar energy stored in the air (considered negligible), 5% of the energy stored in the soil, with most of the heat from the solar radiation (74%) removed through natural ventilation. The average temperature outside was 17°C , while the temperature inside the tunnel increased from 2° to 16°C , depending on cloud cover, wind speed and direction, and how wide the

tunnel doors were opened. It was recommended that the side panels of the tunnel be raised during warmer weather to prevent over-heating.

Most of the tunnels used by commercial producers do not use heating. Heating is more important at high latitudes and cooling and heating is important at low latitudes. Yao et al. (2019) found that it was not economically feasible to use heating to avoid frost damage to apricot trees under tunnels in New Mexico. Venting within the tunnels can be passive, or active with a specific setpoint or threshold for an individual crop (Gent, 1992). There can be large spatial variations in temperature within a tunnel and differences between days and nights (Arabia et al., 2024; Hull et al., 2024; Palma et al., 2023; Yuan et al., 2004).

Wien (2009) demonstrated it was 4°C warmer at 0800 h and 1800 h under a tunnel than outdoors in Ithaca, New York. In contrast, it was 10°C warmer during the middle of the day (Figure 3). Shading under the tunnels decreases ambient temperatures during the day, whereas poor ventilation has the opposite effect. Mean temperatures from 0800 to 1800 h were $24.8^{\circ} \pm 1.9^{\circ}\text{C}$ under the tunnel and $14.8^{\circ} \pm 1.3^{\circ}\text{C}$ in the open. Lozano et al. (2016) conducted similar work in Almonte in Southern Spain. Mean solar radiation from November to June was $12.4 \pm 1.9 \text{ MJ/m}^2/\text{day}$ under the tunnels and $15.3 \pm 2.4 \text{ MJ/m}^2/\text{day}$ in the open (Figure 4). Mean maximum temperatures were $23.3^{\circ} \pm 0.9^{\circ}\text{C}$ under the tunnels and $20.4^{\circ} \pm 1.3^{\circ}\text{C}$ in the open. Mean minimum temperatures were similar in the two areas ($8.6^{\circ} \pm 0.8^{\circ}\text{C}$ and $8.5^{\circ} \pm 0.9^{\circ}\text{C}$).

Ogden et al. (2011) found that tunnels in Georgia warmed up during the day and cooled to ambient or below ambient at night. This response was more pronounced during clear days and nights. Ward and Bomford (2013) indicated that a high tunnel gave 4.3°C of frost protection for flax in Kentucky and increased temperatures by up to 7.4°C on the coldest

days. Yields of tomato in South Africa were higher in a tunnel with a fan and pad cooling system than one dependent on natural ventilation for cooling (Maboko et al., 2012). The temperature in the first tunnel ranged from 13.2° to 38.0°C and from 14.6° to 44.2°C in the second tunnel. Sharaf-Eldin et al. (2023) conducted similar work with tomato in Egypt. They examined various growing systems, including open field, a shaded tunnel with natural ventilation, a net tunnel with fogging and a plastic tunnel with evaporative cooling using a wet pad and fans. The tunnel with evaporative cooling produced the highest yield (3.6 kg/plant) compared with the open field (1.1 kg/plant). The other two treatments were intermediate.

Tunnels can protect strawberry plants from cold weather or frosts and increase fruit yields. However, sometimes temperatures under the tunnels are excessive for acceptable growth and fruit production, for at least part of the season. Leaves in strawberry were on average 2.6°C warmer than the air inside a tunnel in Spain (Peñuelas et al., 1992). This was because it was humid under the plastic and because the leaves were mostly horizontal, and mostly perpendicular to sunlight in the afternoon when they were measured.

Fernandez (2001) examined the potential of tunnels to protect strawberry plants from cold weather in North Carolina. Floating row covers were placed over plots of ‘Chandler’, ‘Camarosa’ and ‘Sweet Charlie’. Daily minimum and maximum temperatures were higher under the covers than in the open when ambient temperatures were above 10°C. The covers increased marketable and total yields by 80 to 100 g/plant compared with plants in the open. Photosynthetic photon flux (PPF) at midday was higher in the open ($1,171 \mu\text{mol/m}^2/\text{s}$) than under the covers ($689 \mu\text{mol/m}^2/\text{s}$). The covers protected the plants from the cold, with adequate light levels for growth under the plastic. Hernandez et al. (2016)




Figure 3. Changes in temperature inside and outside a plastic high tunnel (greenhouse) in Ithaca, New York. Data are from Wien (2009).




Figure 4. Changes in solar radiation and temperatures inside and outside a plastic high tunnel with strawberries in Almonte, Southern Spain. Data are from Lozano et al. (2016).

also investigated the impact of cold weather on strawberry, with the experiments conducted further south in Florida. During the study, there were eight freezing or near-freezing events with a minimum air temperature of -2.8°C . Minimum temperatures inside the row covers ranged between 0.6° and 4.4°C at the canopy level. Plants under the covers had higher marketable yields (21.8 to 23.9 t/ha) than those in the open where sprinklers were used to mitigate against the frosts (16.7 to 17.8 t/ha) ($p < 0.05$).

In another study in Florida, Salamé-Donoso et al. (2010) examined the effect of protected cultivation on the performance of 'Strawberry Festival', 'Winter Dawn', and 'Florida Elyana' over 2 years. Production in this area is affected by cold weather at the start of the season and wet weather for the whole of the season. The plants were grown under tunnels or in the open. In the first year, there was a single freeze event in early January where the minimum temperature under the tunnel was 1°C compared with -3°C in the open. In the second year, there were five freeze events in January and February. The minimum temperature

under the tunnels was 1°C compared with -3° , -6° , -5° , -3° , and -1°C in the open. Both the early and total marketable yields were higher under the tunnels than in the open (Table 5). During the first freeze, 'Strawberry Festival' under the tunnel had the highest early yield (4.8 t/ha) across all the treatments. The same response occurred with the next two freeze events. It was concluded that the other two cultivars were sensitive to freeze events in the open, whereas 'Strawberry Festival' was tolerant to low temperatures, regardless of the production system.

Tunnels can be used to extend the harvest in strawberry, especially in areas that have a cool start to the season. Liu et al. (2024) grew eight cultivars under high tunnels and in the open in Virginia. Fruit were harvested from January to June under the tunnels and from April to June in the open. Average marketable yields were lower under the tunnels (334 ± 26 g/plant) than in the open (516 ± 28 g/plant), with a wide variation in the response across the cultivars. Maximum temperatures under the tunnels in February and March were above 42°C . It was concluded that the

Table 5. Effect of growing system on early and marketable yields of three strawberry cultivars over two seasons in Florida. Means in a column within a treatment group followed by a common letter are not significantly different by the Fisher's least significant test at 5% level of significance. Early yield was from late December to early February. Data are from Salamé-Donoso et al. (2010).

Production system or cultivar	Early marketable yield (t/ha)		Total marketable yield (t/ha)	
	2007–2008	2008–2009	2007–2008	2008–2009
Open field	5.0 b	7.3 b	30.4 b	35.9 b
Tunnel	3.2 a	6.4 a	18.8 a	24.1 a
Strawberry Festival	5.9 c	8.2 c	30.9 c	46.5 b
Winter Dawn	4.1 b	6.8 b	23.4 b	22.5 a
Florida Elyana	2.5 a	5.0 a	19.3 a	20.9 a

tunnels extended the production season, but achieving acceptable yields was challenging in this environment.

Demirsoy et al. (2024) found that temperatures under tunnels were too high for acceptable yields in northern Türkiye. These authors grew nine cultivars under high tunnels and in the open. Mean marketable yield was slightly lower under the tunnels (606 g/plant) than in the open (637 g/plant) ($p > 0.05$). In contrast, fruit weight was higher under the tunnels (14.2 g) than in the open (11.7 g) ($p < 0.01$). Temperatures under the tunnels were higher than in the open. The average daily maximum temperatures in June, July, and August were 38.1°, 35.8°, and 36.4°C under the tunnels and 30.1°, 33.5°, and 31.5°C in the open. The day-neutral (DN) cultivars performed better under the tunnels than the short-day (SD) cultivars.

Tunnels can be used to increase yields, extend the production season and protect strawberry plants from frosts. Temperatures under tunnels are variable, and depend on the weather, the type of tunnel and the use of heating and cooling. Inadequate ventilation or cooling can lead to extreme temperatures and lower yields under the plastic.

Incidence of diseases under tunnels

There are mixed reports on the incidence or severity of diseases when strawberry plants are grown under tunnels. Protecting the plants from rain can eliminate the need for fungicide sprays to control some diseases under the plastic. In contrast, the development of other diseases can be promoted by the high humidity under the tunnels.

There are differences in the development of diseases when crops are grown under protected cultivation. Plastic covers alter light levels, temperatures, and relative humidities, but protect the plants from rain. Differences in the incidence or severity of diseases across growing systems vary with the crop, the pathogen, and the environment (Beacham et al., 2023; O'Connell et al., 2012; Rogers & Wszelaki, 2012; Shishido, 2011; Solís-Mera, 2021). High temperatures and high humidities promote the development of many fungal and bacterial pathogens. Overall, disease issues are less problematic under tunnels. The impact

of many diseases is expected to increase under climate change, with higher temperatures predicted to increase the level of damage to plant tissues (Gallego-Tévar et al., 2024).

Hanson et al. (2011) found that the incidence of grey mould in raspberry incited by *Botrytis cinerea* was only 1% under tunnels and 13% in the open. Leaf and cane diseases were rare under the tunnels and common in the open. Børve and Stensvand (2003) demonstrated that fungicides were not required when cherry trees were grown under a rain shelter in Norway. In contrast, high levels of disease in some crops can make production under tunnels not viable (e.g. Babadoost, 2011 with leaf mould in tomato in Illinois; Buckland et al., 2023 with conventional tomato in Oregon; and Díaz-Pérez et al., 2024 with organic tomato in Georgia). There are also heightened concerns about worker safety that limit the chemicals available for use in enclosed tunnels.

Protected cultivation alters the incidence and severity of several diseases in strawberry, including anthracnose fruit rot (AFR) incited by *Colletotrichum acutatum*, grey mould (BGM) incited by *Botrytis cinerea* and powdery mildew (PM) incited by *Podosphaera aphanis*. The impacts of AFR and BGM are more severe in the open than under tunnels, while the impact of PM is more severe under tunnels (Burlakoti et al., 2013; Evenhuis & Wanten, 2006; Nes et al., 2017; Riikonen et al., 2024). However, some authors have reported the opposite response or no difference between the two systems (Cosseboom et al., 2023; Kennedy et al., 2013). In some experiments, the difference in disease between tunnels and the open varies with the cultivar (e.g. Carisse et al., 2013 with powdery mildew in Canada). Some cultivars have a higher incidence of the disease under the tunnels than others. Applications of UV-C light reduce the severity of PM under both field and tunnel conditions (Mello et al., 2022; Onofre et al., 2021, 2022; Riikonen et al., 2024).

Differences in the development of the three main diseases affecting strawberry are due to variations in temperature and water conditions between the protected and non-protected environments. Both AFR and BGM are promoted by prolonged rain, whereas PM is promoted by high humidity but not by free

water (Aldrighetti & Pertot, 2023; Russi et al., 2024; Yousef et al., 2024). The application of fungicides reduces the differences in diseases between the two systems (e.g. Onofre et al., 2021 with powdery mildew in Florida). Alternatively, steam can be applied to the nursery plants to reduce the infection of plants in the production fields (Stensvand et al., 2024). Burlakoti et al. (2013) examined the effect of protected cultivation on diseases in strawberry over 2 years in Ontario, Canada. The incidences of AFR and BGM were low under tunnels compared with the open field. There was a five- to a twenty-fold lower incidence of AFR under the tunnels and a five-fold lower incidence of AFR. The incidence of PM was higher under the tunnels (7.0% to 12.9% of fruit affected) than in the open (0.6% to 4.9% of fruit affected). Marketable yields were two to three and half times higher under the tunnels than in the open. Tunnels protect plants from rain or free moisture and help to reduce the number of fungicide sprays needed to control AFR (Burlakoti et al., 2014; Xiao et al., 2001). For instance, Burlakoti et al. (2014) indicated that the incidence of AFR in non-sprayed plots was 0.9% to 3.1% under tunnels and 16.0% to 19.3% in the open. Marketable yields in the two plots ranged from 2.364 to 2.460 kg/1.2 m² and from 0.831 to 0.970 kg/1.2 m².

Protected cultivation alters the development of fruit diseases affecting strawberry. The incidence of AFR and BGM is lower under tunnels than in the open, whereas the incidence of PM shows the opposite trend. Tunnels that offer protection from rain can reduce or eliminate the number of fungicide sprays used to control AFR and BGM. In contrast, the incidence of PM can be problematic under plastic covers.

Pollination under tunnels

Strawberry plants require bees or other insects for effective seed and fruit set. This can be a problem under tunnels, with fewer pollinators under tunnels than in the open field.

Pollination is important for seed and fruit set in most flowering plants (Bishop et al., 2021; Kendall et al., 2021). Managed and wild bees, along with flies are the major pollinators of crop and wild species (Faure et al., 2024; Nacko et al., 2022; Feltham et al., 2015; Fenton et al., 2025; Feuerbacher et al., 2024; Mateos-Fierro et al., 2023; Thomasz et al., 2024; Warren et al., 2024).

The importance of pollination for the fertility of plants can be assessed by estimating pollination dependence (PD). This parameter can be calculated by determining fertility after self- and open-pollination (Menzel, 2023). There are various indices of fertility used in the research, including the rate of seed or fruit set, the number of seeds in a fruit, fruit weight, or yield. Species vary in their dependence on pollinators

for seed or fruit set (Ryan et al., 2023; Siopa et al., 2024). The dependence on pollinators ranges from zero (fertility not dependent on pollinators) to one (fertility completely dependent on pollinators). In some plants, the availability and transfer of pollen limit seed or fruit set (Layek et al., 2023; Makowski et al., 2024). Pollen limitation (PL) or pollination deficit varies with the plant and the methods used to assess seed or fruit set. Pollen limitation can be estimated by comparing the fertility of flowers with and without supplementary insects or by comparing the fertility of hand- and open-pollinated flowers. Low values of PL are associated with abundant pollinators and frequent visits of pollinators to the flowers.

Tunnels are a barrier to the movement of pollinating insects, and disrupt their interaction with flowers (Cao et al., 2023; Leach & Isaacs, 2018; Mateos-Fierro et al., 2024; Taniguchi et al., 2025; Wszelaki et al., 2013). Bees have a shorter life-span when placed under plastic covers, due to changes in light levels, temperatures, and relative humidity and may not be effective pollinators (Ogden & van Iersel, 2009). Hall et al. (2020) examined the success of pollination in blueberry and raspberry under tunnels with open ends in New South Wales, Australia. They found a higher abundance and a greater number of visits to the flowers by stingless bees (*Tetragonula carbonara*) and honey bees (*Apis mellifera*) at the ends of the tunnels and less frequent visits towards the middle of the tunnels. The middle of the tunnels had higher temperatures and lower wind speeds. In blueberry, yield ($p < 0.001$) and berry weight ($p < 0.05$) were positively correlated with the abundance of pollinators and were lower at the middle than at the edge of the tunnels. In raspberry, the fruit had better shape in areas when there were high populations of pollinators.

Strawberry plants are moderately dependent on pollination from bees and other insects, with fruit set limited by the availability of pollinators under natural open conditions (Gudowska et al., 2024; Hodgkiss et al., 2019; Menzel, 2023; Pioltelli et al., 2024; Tuohimetsä et al., 2014). Menzel (2023) demonstrated that the mean (\pm s.d.) pollinator dependence (PD) in strawberry (self-pollination versus open- or insect-assisted pollination) was 0.36 ± 0.26 ($p < 0.001$, $n = 52$ studies). The fertility of plants exposed to supplementary insects was higher than those exposed to pollinators under natural open conditions, with a calculated pollen limitation (PL) of 0.20 ± 0.17 ($p < 0.001$, $n = 20$ studies). Nishimoto et al. (2023) demonstrated that there were no differences in the percentage of marketable fruit in strawberry after pollination by hand or with robots under protected cultivation in Japan. There were also no differences in fruit weight or the number of achenes per fruit between the two treatments.

Piovesan et al. (2019) collected data on potential pollinators for strawberry under tunnels in Brazil. The

flowers were visited by 47 species of insects. *Apis mellifera* or honey bee (Hymenoptera: Apidae) was the most abundant, constant, dominant, and frequent species. Twelve species of native bees were identified, including *Tetragonisca fiebrigi*, *Tetrapedia* sp., *Trigona spinipes*, *Schwarziana quadripunctata*, *Plebeia emerina*, *P. remota*, *Bombus pauloensis* (Hymenoptera: Apidae), *Dialictus* sp.1, *Dialictus* sp.2, *Augochloropsis* sp.1, *Augochloropsis* sp.2, and *Augochlora* sp.1 (Hymenoptera: Halictidae). It was concluded that all these species were potential pollinators of the crop under covers. The native species, *T. fiebrigi*, *P. emerina*, and *P. remota* were the most abundant potential pollinators and were relatively easy to manage under the tunnels.

Antunes et al. (2007) found that the yields of four strawberry cultivars under protected cultivation in Brazil were increased when hives of jatai bees (*Tetragonisca angustula*) were introduced into the growing area. Average yields were 686 g/plant in the control and 968 g/plant with four hives per growing area. Wilkaniec and Radajewska (1997) reported similar results using the solitary bee (*Osmia rufa*) under tunnels in Poland. Average yields were 289 ± 77 g/plant with self-pollination and 434 ± 153 g/plant with bees. Howard et al. (2021) indicated that honey bees were better pollinators than other insects when strawberry plants were grown under tunnels in Victoria, Australia. Their work showed that the honey bees spent more time visiting the flowers than the other insects. Willden et al. (2022) investigated the pollination of strawberry under tunnels in New York. There was evidence that UV-blocking plastics resulted in poorly pollinated fruit compared with UV-transmitting plastic and open-field treatments. However, the mechanism causing this response was not clear.

Ariza et al. (2012) studied the effect of pollination on the performance of strawberry under tunnels in Spain. Plots under the tunnels were provided with bumble bees (*Bombus terrestris*) or pollinated under natural conditions (controls). From January to March, the incidence of misshapen fruit was $25.7 \pm 2.3\%$ with supplementary bees and $61.3 \pm 3.8\%$ without bees ($p < 0.05$). From April to May, the incidence of misshapen fruit was $13.4 \pm 2.8\%$ with bees and $20.3 \pm 1.9\%$ without bees ($p < 0.05$). Supplementary pollination was advisable for strawberry under covers, especially in cultivars such as 'Camarosa' that were prone to misshapen fruit. Sarıdaş et al. (2021) collected similar information in Türkiye. One tunnel was covered with a monofilament UV-stabilised white net (88% to 98% light transmission) to prevent pollinators from visiting the flowers. One commercial honey bee hive was placed near that tunnel, which allowed bees to enter the other tunnel. The plants supplied with bees had 54% higher yields (1,175 g/plant) than those not supplied with bees (763 g/plant)

($p < 0.05$). The plants with bees also had larger fruit (17.6 g versus 15.0 g) ($p < 0.05$).

Strawberry requires successful pollination for acceptable productivity and quality. Yields and fruit weight are lower under tunnels if the insects have poor access to the flowers compared with conditions in the open field. Supplementary pollination with bees or other insects can improve production under tunnels.

Profitability under tunnels

The economics of growing strawberry plants under tunnels varies around the globe. Most studies indicate that production under plastic covers is profitable, although there are exceptions.

The costs of production are higher when crops are grown under tunnels than in the open field. The economics of tunnels depends on the costs of production, yields, and the price received for the product (DiGiacomo et al., 2023; Ho et al., 2018; Sydorovych et al., 2013). In some cases, tunnels offer improved control of pests and diseases, but the technology is not economically viable (Nordey et al., 2020). Different analyses can be used to determine profitability, including net returns, net present worth, internal rate of return, benefit-to-cost ratio, and the pay-back period (Bodiroga et al., 2022; Morris et al., 2023; Nian et al., 2022; Sengar & Kothari, 2008; Subedi et al., 2023).

Galinato and Miles (2013) found that lettuce and tomato yielded three to four times more under tunnels than in the open in Washington State. Given the base crop yield and average price, it was 43% more profitable to grow lettuce in the open than under the tunnels. In contrast, tomatoes under the tunnels were three-times more profitable than those in the open. Ward et al. (2011) investigated the economics of speciality crops under tunnels in the United States. They found that even when land prices were high, tunnels were profitable, provided the produce was grown out-of-season and sold at farmers' markets. Chase (2013) investigated the production of vegetable crops under tunnels in Iowa. He found that the costs of the low-technology tunnels were paid back within a year, which is unusual in horticulture. Ernst (2020) indicated that it took 3 years to recover the additional costs of vegetables under low tunnels in Kentucky. Waterer (2003) examined the profitability of muskmelons, peppers, and tomatoes under low or high tunnels in Saskatchewan, Canada. The high tunnels were more costly to purchase and construct than low tunnels, but they were durable enough to be used for multiple seasons. Based on wholesale commodity prices, it took 2–5 years for the returns obtained with high tunnels to cover their higher capital costs.

Research on the profitability of strawberry under tunnels has given mixed results, with both positive and negative returns over costs (Clark, 2023;

Mbarushimana et al., 2022; Salamé-Donoso et al., 2010). Marin and Garcia (2004) collected economic data under tunnels in Spain. They indicated that there was range in income for different enterprises in Andalusia. The total net margin ($\text{€} \times 1000/\text{ha}$) was variable across the areas (mean \pm s.d.): Almeria (14.75 ± 57.6), Granada (12.57 ± 48.85), Huelva (23.2 ± 114.29), and Málaga (21.80 ± 38.48). This response was due to differences in production strategies, the availability of water, the type of irrigation technology, and the weather. The net margin was higher under the tunnels than in the open due to a more stable environment and more stable yields.

Zhao et al. (2015) demonstrated that overall net returns were higher under high tunnels (USD 51,755 \pm 26,317/ha) than in the open in Florida (USD 29,335 \pm 7,871/ha). The difference in returns between the two systems decreased with the increasing costs of the tunnels and the shorter life-span of the plastic. When the plastic lasted 8 years and the tunnel cost USD 56/m², tunnel production was less profitable than open production. However, when the plastic lasted 9–15 years, tunnel production was more profitable, even when the tunnels cost USD 57/m². Mbarushimana et al. (2022) investigated the performance of eight cultivars under high tunnels and in the open in Virginia. Averaged over three marketing strategies, net returns were lower under the tunnels (minus USD 35,055 \pm 19,807/ha) than in the open (positive USD 77,926 \pm 29,641/ha). Poor returns under the tunnels were due to low yields and high costs.

Salamé-Donoso et al. (2010) grew three strawberry cultivars under high tunnels and in the open over 2 years in Florida. Early marketable yields were up to 54% higher under the tunnels than in the open, and total marketable yields were up to 63% higher (Table 5). Higher yields under the tunnels were due to protection from frosts. The higher yields provided an additional USD 28,875/ha in gross returns, off-setting the costs of the tunnels of USD 15,000/ha each season. The costs of harvesting and marketing were not included in the analysis.

Lewers et al. (2017) examined the profitability of strawberry under low tunnels in Maryland. Marketable yields were 313% higher under the tunnels than in the open. The costs of the material used to construct the tunnels were calculated to determine if

they could be off-set by the extra production. An increase in the production value of USD 0.59/plant over 2 years was required to pay for the materials needed to construct a tunnel. Yields increased from 230 g/plant in the open to 720 g/plant under the tunnels, suggesting higher profitability under the plastic. The cost of labour to construct the tunnels was not included in these calculations. Orde and Grube Sideman (2021) evaluated the performance of day-neutral (DN) strawberry under low tunnels and in the open over 2 years in New Hampshire. Marketable yields were unaffected by the covers in the first year and were higher under the tunnels in the second year. Plants under the tunnels had a market value of USD 3,899 to 95,647/ha, depending on the cultivar and year. The covers cost USD 1,557/ha, assuming a life-span of 3 years. The row covers were profitable in this environment.

Maughan et al. (2015) investigated the costs and returns for strawberry under high tunnels in Utah. The tunnels provided a net return of USD 15,549/ha. The economics of open production was not recorded in the study. Sullivan et al. (2022) provided similar data in Oregon. These authors grew strawberry under high tunnels or in the open. They reported that a 15.2 m long tunnel gave a return of USD 1,970 compared with USD 1,344 for the open field, a 46.6% increase. No information was provided on the costs of the tunnel or on the life-span of the plastic.

Schmutz et al. (2011) examined the economics of organic strawberry under high tunnels in the United Kingdom. The field was planted out at 50,000 plants/ha and produced yields of 15 t/ha worth £60,000/ha. The cost of establishment was £15,624, variable costs were £40,424 and the gross margin was £19,576. Jirgena et al. (2013) studied the risks and returns of strawberry under different production systems in Latvia. The tunnels provided a better environment than the open field with fewer pests in the crop. This response contributed to a longer and more predictable shelf life for the fruit (Table 6). The tunnels extended the harvest season, with earlier harvests than in the open and higher prices for the crop. In the open area with minimum investments, aggregate expenditure exceeded total revenue and the system provided a negative return (Table 7). The open area managed

Table 6. Probabilities and severities of yield risks in strawberry production under different growing systems in Latvia. Risk severity is expressed from zero (no risk) to ten (severe risk). Data are from Jirgena et al. (2013).

Risks	Probability	Severity of risk to yield			
		Open area, with minimum investment	Open area	Plastic cover	High tunnel
Winter frost damage	0.1	10.0	10.0	10.0	10.0
Early moderate frosts	1.0	1.8	1.8	0	0
Early severe frosts	0.4	8.5	8.5	2.0	2.0
Hail	0.1	9.0	9.0	9.0	0
Heavy rain	0.3	4.5	4.5	4.5	1.0
Pests & diseases	0.5	7.5	7.5	7.5	1.0

intensively provided better returns on investment than that under the tunnels. Production in open areas under cover provided the best returns on investment.

McIntosh (2018) examined the performance of different production systems in southern Australia. Yields and gross margins were higher with substrate production under tunnels or in glasshouses than with soil production in the open field (Table 8). However, capital costs were much higher under protected cultivation than in the open. McIntosh (2018) indicated that the plants under protected cultivation had a lower incidence of some fruit diseases than those in the open.

Greco et al. (2020) and de Tommaso et al. (2021) analysed the performance of strawberry under tunnels and in the open in Europe. This market is worth € 2.4 billion at a consumer level, about 1% of the total value of agricultural production in the area. In the two studies, yields were higher under the tunnels in Spain and Belgium, and lower in the open in Italy and France (Table 9). Net returns were higher under the tunnels in Italy and lower in the open in Italy and Belgium. Where there was a direct comparison of the two systems, average returns were €13,229 ± 2,202/ha under the tunnels and €8,160 ± 1,265/ha in the open.

The economic performance of strawberry under tunnels is highly variable. Net returns vary with

yields, marketing strategies, and the costs of the tunnels. In some cases, plants under tunnels have higher yields than those in the open or they produce fruit out-of-season when prices are high. In these cases, the returns for the fruit can exceed the costs of the tunnels. In other cases, plants under tunnels have lower yields than those in the open or they produce fruit during the traditional season. In these cases, the returns for the fruit can be lower than the costs of the tunnels. The life-span of the covers has a strong effect on profitability. Plastics that last 8 or 10 years are more economical than those that last for shorter periods.

Use of tunnels to minimize the impacts of global warming on strawberry production

Several studies indicate that the yields of strawberry will be lower under global warming. Modelling in California demonstrated that productivity will decrease by 10% by 2050 and by 40% by 2099 (Deschenes & Kolstad, 2011; Lobell & Field, 2011). Higher temperatures under tunnels might exacerbate the effects of global warming on commercial production in the future.

Protected cultivation can be used to mitigate against the effects of climate change on the

Table 7. Return on investment for strawberry production under different growing systems in Latvia. Data are from Jirgena et al. (2013).

Growing system	Revenue (€/ha)	Investment (€/ha)	Expenditure (€/ha)	Gross profit (€/ha)	Return on investment
Open area, with minimum investment	12,177	11,666	15,972	-3,975	-33%
Open area	19,997	13,893	17,312	2,665	19%
Plastic cover	35,177	23,102	25,943	9,234	40%
High tunnel	104,209	81,866	97,250	6,959	9%

Table 8. The costs and returns of strawberries growing in the open field, under tunnels with substrates, and in greenhouses with substrates in Australia. Values are in AUD (Australian dollars). Data are from McIntosh (2018).

Costs, yields & gross margins	Open field	Tunnel/substrate	Greenhouse/substrate
Total capital costs (\$ per ha)	142,239	422,239	887,239
Total variable costs (\$ per ha)	183,399	300,799	396,331
Marketable yield (t/ha)	48.7	72.7	90.2
Gross returns (\$/ha)	205,434	446,855	558,249
Gross margin (\$/ha)	22,035	146,056	161,919
Gross margin (% of gross returns)	11%	33%	29%

Table 9. Yields and net returns of strawberries growing in the open field and under tunnels in Europe with soil fumigation. Mean values and standard errors (s.e.) also shown where there was a direct comparison of the two systems (values for tunnels in Spain excluded). Data are from Greco et al. (2020) and de Tommaso et al. (2021).

Country	Growing system	Yield (t/ha)	Net returns (€/ha)
Spain	Tunnel	55.1	13,611
Italy	Open field	12.7	7,892
	Tunnel	35.0	18,338
Belgium	Open field	18.4	5,624
	Tunnel	50.0	9,195
France	Open field	11.1	10,965
	Tunnel	22.6	12,154
Mean (± s.e.)	Open field	14.1 ± 1.8	8,160 ± 1,265
	Tunnel	35.9 ± 6.5	13,229 ± 2,202

productivity of some crops (Gruda et al., 2019; Jamarkattel et al., 2023; Nordey, Faye, et al., 2020; Rabbi et al., 2019; Sandison et al., 2023). Tunnels or similar structures can be modified to improve light levels, temperature, or relative humidity for better growth and yields (Fernandes de Oliveira & Nieddu, 2015 in Italy; Gruda et al., 2021 in Greece; Ahmad et al., 2023 in Pakistan; Sharaf-Eldin et al., 2023 in Egypt). Strawberry plants are grown under tunnels to protect the plants from cold weather, frosts, and rain. Temperatures under the plastic covers can be excessive for acceptable growth and yields, in the absence of adequate ventilation. Close attention to this issue is important for growing strawberry and other plants under plastic (Conner & Demchak, 2018; Yoneda & Kawashima, 2024).

Fatnassi et al. (2009) demonstrated that ventilation within a tunnel decreased with the height and density of the plants under the plastic, along with the direction of the wind. They recommended that the sides of tunnels should be rolled up to improve wind movement and ventilation. Bartzanas et al. (2004) modelled air movement and temperatures within a tunnel in France. The mean temperature in the middle of the tunnel varied from 28.2° to 29.8°C for an outside temperature of 28.0°C. However, there were regions inside the tunnels that were 6.0°C warmer than outside. Average air velocity in the tomato crop canopy varied according to the arrangement of the vents in the tunnel from 0.2 to 0.7 m/s. Ventilation can aid air movement within a tunnel, but it does not always lower the temperature. Lewus and Both (2022) modelled air-flow and temperatures under a high tunnel with different side and roof vents in Pennsylvania, United States. The roof vents increased mass-based ventilation through the tunnel by 20% to 78% compared with that in the standard tunnel with rolled-up, side vents only. However, the roof vents only lowered the temperature inside the tunnel by 0.1°C.

Rogers and Wszelaki (2012) demonstrated temperatures under tunnels growing tomatoes could be excessive in Tennessee. Despite having the side and end walls of the tunnels open, average temperatures reached 37.8°C during the middle of the day. The highest temperature recorded over a single day over the 2-year experiment under the tunnels was 51.7°C. The mean temperature range from April to August (in 2009) was 10.6° to 13.9°C in the open, a difference of 3.3°C. The mean temperature range under the tunnels during the same period was 22.2° to 27.2°C, a difference of 5.0°C. This trend was repeated in 2010. The tunnels did not retain heat into the evening, where average temperatures were similar with those in the open. Ventilation within the tunnel was inadequate for control of the temperature under the plastic.

Gázquez et al. (2008) examined different cooling strategies for sweet pepper under tunnels in Spain.

Fogging was the most efficient method for controlling maximum air temperatures and vapour pressure deficits (VPDs). In contrast, it was the least efficient method for controlling canopy temperatures. Plants that had been shaded and grown under plastic that had been white-washed had higher yields (10.23 kg/m²) than those grown under shade and fogging (8.41 kg/m²) ($p < 0.05$). Plants subjected to fogging had the highest incidence of blossom-end rot (BER). An economic evaluation showed that whitening was the most profitable cooling treatment. The researchers concluded that a combination of whitening and natural ventilation was the most efficient cooling system in terms of water and energy use. O'Connell and Tate (2017) investigated the use of protected cultivation to produce organic broccoli and cauliflower in Georgia. They recommended that the side curtains should be opened when the temperature was higher than 15.6 ± 1°C. Conversely, tall side walls (1.8 m openings) and end walls (4.9 m openings) helped to ventilate the tunnels on warm days and keep temperatures and relative humidity levels near the crop canopy similar to those in the open.

Sethi et al. (2009) examined the growth of vegetables under protected cultivation in Ludhiana, India (latitude 31°N). They reported that a combination of low tunnels in winter and shade nets in summer gave 37.6% and 11.5% higher yield in the brinjal crops compared with a conventional net house or plastic-covered greenhouse for the whole year. This was because the combined strategy provided optimum temperature and light conditions for growth. Thihe et al. (2020) collected data on the microclimate in a fan-pad evaporative cooled tunnel (FPVT) and an open-ended, naturally ventilated tunnel (NVT) in Pietermaritzburg, South Africa (latitude 29.7°S). The tunnels had floor areas of 144 m² and ridge heights of 3.5 m, and were 18 m in length, with a floor width of 8 m. The temperature in the NVT was 4° to 5°C higher than in the FPVT at midday when solar radiation levels peaked. At night, there was no difference in temperature between the two tunnels. Relative humidity in the NVT ranged from 49% to 90% and 59% to 95% in the FPVT. Kittas et al. (2005) modelled temperatures inside a plastic greenhouse in Greece and found that the difference between the inside and outside temperature was related to the ventilation rate and the incoming solar radiation. Temperatures inside the greenhouse were up to 8°C higher inside when there was poor ventilation. In contrast, temperatures were similar in the two areas when there was good ventilation.

Plastic tunnels can have the ends of the structures open and the sides of the tunnels rolled up to assist ventilation. Temperatures within single tunnels are easier to manage than those in multiple tunnels, although the costs of construction are higher. Menzel

et al. (2014) grew strawberry under tunnels in southern Queensland, Australia. The sides of the tunnels were rolled up when it was not raining. They showed that temperatures at canopy level were similar inside and outside the tunnels. In the first year of the experiment, the average daily maximum temperature was 26.0°C inside the tunnel and 26.5°C outside. Minimum temperatures were 11.9° and 10.5°C. In the second year, the average daily maximum temperature was 25.4°C inside the tunnel and 26.3°C outside. Minimum temperatures were 13.1° and 11.7°C.

Light-blocking films can be used to reduce heat-generating light from the 780 to 2,500 nm (infrared and far-infrared) region, while allowing most light in the photosynthetic active radiation (400 to 700 nm) region. However, their use in greenhouses and similar structures usually results in lower yields compared with control plots (no light-blocking films) (Chavan et al., 2023; He et al., 2024; Lin et al., 2024; Maier et al., 2023). Further research is required to determine if these films can reduce the impact of global warming for plants growing under protected cultivation.

Tunnels protect strawberry crops from cold, frosts, and rainfall. Temperatures under the covers can be too high for acceptable growth, depending on the weather, type of plastic, and ventilation. Temperatures inside the tunnels can be up to 10°C higher than those outside during the day. Temperatures inside the tunnel are similar to those outside during the night. The use of tunnels under global warming will depend on good ventilation and other strategies (light shading) to prevent excessive temperatures building up under the plastic.

Conclusions

Plants under tunnels had higher relative marketable (Tunnel/Open = 1.34 ± 0.76) and total yields (Tunnel/Open = 1.30 ± 0.83) than those in the open field. In contrast, fruit weight was similar in the two systems (Tunnel/Open = 1.04 ± 0.22). The response to protected cultivation (tunnel versus open field) were similar in Northern and Southern Europe, in Northern and Southern America, in areas with cool or cold winters or with spring/summer or winter/spring production seasons, and under low or high tunnels. Lower yields under the tunnels were associated with shading and high temperatures under the plastic and a higher incidence of powdery mildew (PM). Tunnels offer protection from rain and can help to reduce or eliminate fungicides for the control of anthracnose fruit rot (AFR) and botrytis grey mould (BGM). In contrast, the incidence of PM can be problematic under covers. Yields and fruit weight are lower under tunnels without supplementary pollination by bees or other insects. The economics of strawberry under tunnels is highly variable. Net returns vary with yields, marketing strategies, and the costs of the tunnels. Protected cultivation can be

used to mitigate against the effects of climate change on productivity, but producers need to pay close attention to ventilation under the plastic. There are concerns about the impact of the different plastics on the environment, and potential issues with the exposure of farm workers to pesticides under the tunnels.

Acknowledgements

The Queensland Government funded the research through the Department of Agriculture and Fisheries. Financial support from the Florida Strawberry Growers' Association (FSGA) is appreciated. Special thanks to Helen Macpherson, Danielle Hoffmann, Cheryl Petroeschovsky and Cass Butler from DAF for supplying much of the literature.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

The work was supported by the Florida Strawberry Research and Education Foundation [HF 11069].

Data availability statement

The author confirms that the data supporting the findings of this study are available within the supplementary materials published online with this paper or available from the author on reasonable request.

References

- Ahmad, B., Nadeem, M. U., Liu, T., Asif, M., Rizvi, F. F., Kamran, A., Virk, Z. T., Jamil, M. K., Mustafa, N., Saeed, S., & Abbas, A. (2023). Climate change impact on groundwater-based livelihood in Soan River Basin of Pakistan (South Asia) based on the perception of local farmers. *Water*, 15(7), 1287. <https://doi.org/10.3390/w15071287>.
- Ahmadi, M., Atefi, A., Ramzanpour, M., & Lin, J. (2024). A two-step deep semantic segmentation and object detection approach for runner recognition in strawberry plants. *International Journal of Fruit Science*, 24(1), 293–300. <https://doi.org/10.1080/15538362.2024.2397438>
- Akpenpuun, T. D., Na, W.-H., Ogunlowo, Q. O., Rabi, A., Adesanya, M. A., Addae, K. S., Kim, H.-T., & Lee, H.-W. (2021). Effect of greenhouse cladding materials and thermal screen configuration on heating energy and strawberry (*Fragaria ananassa* var. "Seolhyang") yield in winter. *Agronomy*, 11(12), 2498. <https://doi.org/10.3390/agronomy11122498>
- Alam, E., Moyer, C., Verma, S., Peres, N. A., & Whitaker, V. M. (2024). Exploring the genetic basis of resistance to *Neopestalotiopsis* species in strawberry. *The Plant Genome*, 17(2), e20477. <https://doi.org/10.1002/tpg2.20477>
- Aldrichetti, A., & Pertot, I. (2023). Epidemiology and control of strawberry powdery mildew: A review.

- Phytopathologia Mediterranea*, 62(3), 427–453. <https://doi.org/10.36253/phyto-14576>.
- Al-Helal, I., Alsadon, A., Shady, M., Ibrahim, A., & Abdel-Ghany, A. (2020). Diffusion characteristics of solar beams radiation transmitting through greenhouse covers in arid climates. *Energies (Basel)*, 13(2), 472. <https://doi.org/10.3390/en13020472>
- Al-Madani, A. A., Al-Helal, I. M., & Alsadon, A. A. (2024). Assessing the effectiveness of reflective and diffusive polyethylene films as greenhouse covers in arid environments. *Agronomy*, 14(5), 1082. <https://doi.org/10.3390/agronomy14051082>
- Amyotte, B., & Samtani, J. (2023). New directions for strawberry research in the 2020s. *International Journal of Fruit Science*, 23(1), 278–291. <https://doi.org/10.1080/15538362.2023.2274894>
- Anderson, H. C., Rogers, M. A., & Hoover, E. E. (2019). Low tunnel covering and microclimate, fruit yield, and quality in an organic strawberry production system. *HortTechnology*, 29(3), 590–598. <https://doi.org/10.21273/HORTTECH04319-19>.
- Antunes, O. T., Calvete, E. O., Rocha, H. C., Nienow, A. A., Cecchetti, D., Riva, E., & Maran, R. E. (2007). Produção de cultivares de morangueiro polinizadas pela abelha jataí em ambiente protegido. *Horticultura Brasileira*, 25(1), 94–99. <https://doi.org/10.1590/S0102-05362007000100018>
- Arabia, A., Pallarés, N., Munné-Bosch, S., & Muñoz, P. (2024). Variability in strawberry tunnels impacts fruit quality and limits melatonin effects. *Journal of the Science of Food and Agriculture* 105(3), 1745–1759. <https://doi.org/10.1002/jsfa.13951>
- Arancibia, R. A., Mecham, K., Harper, J., & Recker, C. (2023). Watermelon production under protected culture in Missouri, USA, to reach the local Fourth of July market. *HortTechnology*, 33(6), 520–526. <https://doi.org/10.21273/HORTTECH05283-23>
- Argento, S., García, G., & Treccarichi, S. (2024). Sustainable and low-input techniques in Mediterranean greenhouse vegetable production. *Horticulturae*, 10(9), 997. <https://doi.org/10.3390/horticulturae10090997>
- Arif, M. F., Aristya, G. R., & Kasiamdari, R. S. (2019). Genetic diversity of strawberry cultivars in Banyuroto, Magelang, Indonesia based on cleaved amplified polymorphic sequence. *Biodiversitas Journal of Biological Diversity*, 20(6), 1721–1728. <https://doi.org/10.13057/bio-div/d200631>
- Ariza, M. T., Soria, C., Medina-Mínguez, J. J., & Martínez-Ferri, E. (2012). Incidence of misshapen fruits in strawberry plants grown under tunnels is affected by cultivar, planting date, pollination, and low temperatures. *HortScience*, 47(11), 1569–1573. <https://doi.org/10.21273/HORTSCI.47.11.1569>
- Asseng, S., Guarina, J. R., Raman, M., Monje, O., Kiss, G., Despommier, D. D., Meggers, F. M., & Gauthier, P. P. G. (2020). Wheat yield potential in controlled-environment vertical farms. *Proceedings of the National Academy of Sciences of the United States of America* 117(32), 19131–19135. <https://doi.org/10.1073/pnas.2002655117>.
- Azad, M. O. K., Gruda, N. S., & Naznin, M. T. (2024). Energy efficiency of glasshouses and plant factories for sustainable urban farming in the desert southwest of the United States of America. *Horticulturae*, 10(10), 1055. <https://doi.org/10.3390/horticulturae10101055>.
- Babadoost, M. (2011). Leaf mold (*Fulvia fulva*), a serious threat to high tunnel tomato production in Illinois. *Acta Horticulturae*, 914, 93–96. <https://doi.org/10.17660/ActaHortic.2011.914.14>
- Balocco, C., Mercatelli, L., Azzali, N., Meucci, M., & Grazzini, G. (2018). Experimental transmittance of polyethylene films in the solar and infrared wavelengths. *Solar Energy*, 165, 199–205. <https://doi.org/10.1016/j.solener.2018.03.011>
- Barnes, P. W., Robson, T. M., Zepp, R. G., Bornman, J. F., Jansen, M. A. K., Ossola, R., Wang, Q.-W., Robinson, S. A., Foereid, B., Klekociuk, A. R., Martinez-Abaigar, J., Hou, W.-C., Mackenzie, R., & Paul, N. D. (2023). Interactive effects of changes in UV radiation and climate on terrestrial ecosystems, biogeochemical cycles, and feedbacks to the climate system. *Photochemical & Photobiological Sciences*, 22(5), 1049–1091. <https://doi.org/10.1007/s43630-023-00376-7>
- Bartzanas, T., Boulard, T., & Kittas, C. (2004). Effect of vent arrangement on windward ventilation of a tunnel greenhouse. *Biosystems Engineering*, 88(4), 479–490. <https://doi.org/10.1016/j.biosystemseng.2003.10.006>.
- Beacham, A. M., James, K. L., Randall, N. P., & Monaghan, J. M. (2023). Challenges and recommendations for the development of cultural control of aerial oomycete-associated diseases in protected horticulture. *European Journal of Plant Pathology*, 167(2), 207–219. <https://doi.org/10.1007/s10658-023-02695-y>
- Beech, M. G., & Simpson, D. W. (1989). Strawberry production in the United Kingdom. *Acta Horticulturae*, 265, 693–696. <https://doi.org/10.17660/ActaHortic.1989.265.107>
- Bertuglia, A., & Calatrava, J. (2012). Identifying indirect factors influencing productivity in protected horticulture: The case of greenhouses in the Spanish Mediterranean coastline. *Acta Horticulturae*, 927, 855–861. <https://doi.org/10.17660/ActaHortic.2012.927.106>
- Bishop, J., Nakagawa, S., & Requier, F. (2021). Quantifying crop pollinator dependence and its heterogeneity using multi-level meta-analysis. *Journal of Applied Ecology*, 58(5), 1030–1042. <https://doi.org/10.1111/1365-2664.13830>
- Black, B. (2010). *Temperature management in high tunnels*. Co-operative Extension. Utah State University.
- Bodiroga, R., Sredojević, Z., Vico, G., & Lučić-Govedarica, A. (2022). Influence of different production methods on economic results of greenhouse vegetable production. *Custos e Agronegocio*, 18(2), 44–62.
- Børve, J., & Stensvand, A. (2003). Use of a plastic rain shield reduces fruit decay and need for fungicides in sweet cherry. *Plant Disease*, 87(5), 523–528. <https://doi.org/10.1094/PDIS.2003.87.5.523>
- British Summer Fruits. (2017). The impact of Brexit on the UK soft fruit industry. *British Summer Fruits Seasonal Labour Report*, 25, 280–286.
- Bruce, A. B., Farmer, J. R., Maynard, E. T., & Valliant, J. C. D. (2019). Using high tunnels to extend the growing season and improve crop quality and yield: Assessing outcomes for organic and conventional growers in the U.S. Midwest. *International Journal of Agricultural Management*, 8(2), 45–55. <https://doi.org/10.5836/ijam/2019-08-45>.
- Buckland, K. R., Ocamb, C. M., Rasmussen, A. L., & Nackley, L. L. (2023). Reducing powdery mildew in high-tunnel tomato production in Oregon with ultra violet-C lighting. *HortTechnology*, 33(2), 149–151. <https://doi.org/10.21273/HORTTECH05139-22>
- Burlakoti, R. R., Zandstra, J., & Jackson, K. (2013). Comparison of epidemiology of gray mold, anthracnose fruit rot, and powdery mildew in day-neutral strawberries in field and high-tunnel conditions in Ontario.

- International Journal of Fruit Science*, 13(1–2), 19–27. <https://doi.org/10.1080/15538362.2012.696956>
- Burlakoti, R. R., Zandstra, J., & Jackson, K. (2014). Evaluation of epidemics and weather-based fungicide application programmes in controlling anthracnose fruit rot of day-neutral strawberry in outdoor field and protected cultivation systems. *Canadian Journal of Plant Pathology*, 36(1), 64–72. <https://doi.org/10.1080/07060661.2014.895422>
- Cao, Z., He, G., Mu, S., & Qu, H. (2023). Effects of bee density and hive distribution on pollination efficiency for greenhouse strawberries: A simulation study. *Agronomy*, 13(3), 731. <https://doi.org/10.3390/agronomy13030731>
- Carisse, O., Lefebvre, A., Van der Heyden, H., Roberge, L., & Brodeur, L. (2013). Analysis of incidence–severity relationships for strawberry powdery mildew as influenced by cultivar, cultivar type, and production systems. *Plant Disease*, 97(3), 354–362. <https://doi.org/10.1094/PDIS-05-12-0508-RE>
- Carter, S., Shaw, S. A., & Harris, N. (1993). Opportunities and change in the British strawberry industry. *British Food Journal*, 95(10), 18–22. <https://doi.org/10.1108/00070709310048498>
- Casal, C., Vílchez, C., Forján, E., & de la Morena, B. A. (2009). The absence of UV-radiation delays the strawberry ripening but increases the final productivity, not altering the main fruit nutritional properties. *Acta Horticulturae*, 842, 159–162. <https://doi.org/10.17660/ActaHortic.2009.842.19>
- Cayambe, J., Heredia-R, M., Torres, E., Puhl, L., Torres, B., Barreto, D., Heredia, B. N., Vaca-Lucero, A., & Diaz-Ambrona, C. G. H. (2023). Evaluation of sustainability in strawberry crops production under greenhouse and open-field systems in the Andes. *International Journal of Agricultural Sustainability*, 21(1), 2255449. <https://doi.org/10.1080/14735903.2023.2255449>
- Cerruto, E., Manetto, G., Santoro, F., & Pascuzzi, S. (2018). Operator dermal exposure to pesticides in tomato and strawberry greenhouses from hand-held sprayers. *Sustainability*, 10(7), 2273. <https://doi.org/10.3390/su10072273>
- Chase, C. (2013). *Vegetable production budgets for a high tunnel*. Extension and Outreach, Iowa State University.
- Chavan, S. G., Chen, Z.-H., Ghannoum, O., Cazzonelli, C. I., & Tissue, D. T. (2022). Current technologies and target crops: A review on Australian protected cropping. *Crops*, 2(2), 172–185. <https://doi.org/10.3390/crops2020013>
- Chavan, S. G., He, X., Maier, C., Alagoz, Y., Anwar, S., Chen, S. H., Ghannoum, O., Cazzonelli, C. I., & Tissue, D. T. (2023). An energy-saving glasshouse film reduces seasonal, and cultivar dependent capsicum yield due to light limited photosynthesis. *Annals of Agricultural Sciences*, 68(1), 21–35. <https://doi.org/10.1016/j.aos.2023.04.001>
- Clark, A., Shephard, C., Robson, A., McKechnie, J., Morrison, R. B., & Rankin, A. (2023). A multifaceted approach to developing an Australian national map of protected cropping structures. *Land*, 12(12), 2168. <https://doi.org/10.3390/land12122168>
- Clark, S. (2023). *Budget analysis of organic high-tunnel strawberry production on the Berea College farm*. Berea College.
- Clark, S., & Mousavi-Avval, S. H. (2022). Global warming potential of organic strawberry production under unheated high tunnels in Kentucky, USA. *Sustainability*, 14(3), 1778. <https://doi.org/10.3390/su14031778>
- Condori, B., Fleisher, D. H., & Lewers, K. (2017). Relationship of strawberry yield with microclimate factors in open and covered raised-bed production. *Transactions of the American Society of Agricultural and Biological Engineers*, 60(5), 1511–1525. <https://doi.org/10.13031/trans.12371>
- Conner, D. S., & Demchak, K. (2018). Farmer perceptions of tunnels for berry production: Management and marketing implications. *HortTechnology*, 28(6), 706–710. <https://doi.org/10.21273/HORTTECH04147-18>
- Cordeiro, G. D., & Dötterl, S. (2023). Global warming impairs the olfactory floral signaling in strawberry. *BMC Plant Biology*, 23(1), 549. <https://doi.org/10.1186/s12870-023-04564-6>
- Cosseboom, S. D., Schoeneberg, A., Lea-Cox, J. D., Samtani, J., Johnson, C. S., & Hu, M. (2023). Impact of floating row cover and sensor placement on strawberry anthracnose and botrytis fruit rot risk assessment. *Plant Pathology*, 72(4), 819–828. <https://doi.org/10.1111/ppa.13695>
- Cusworth, S. J., Davies, W. J., Mcainsh, M. R., & Stevens, C. J. (2022). Sustainable production of healthy, affordable food in the UK: The pros and cons of plasticulture. *Food and Energy Security*, 11(4), e404. <https://doi.org/10.1002/fes3.404>
- Cusworth, S. J., Davies, W. J., Mcainsh, M. R., Stevens, C. J., & Wang, W. (2024a). A nationwide assessment of microplastic abundance in agricultural soils: The influence of plastic crop covers within the United Kingdom. *Plants, People, Planet*, 6(2), 304–314. <https://doi.org/10.1002/ppp3.10430>
- Cusworth, S. J., Davies, W. J., Mcainsh, M. R., Stevens, C. J., & Wang, W. (2024b). Sustainable plasticulture in Chinese agriculture: A review of challenges and routes to achieving long-term food and ecosecurity. *Frontiers of Agricultural Science and Engineering*, 11(1), 155–168. <https://doi.org/10.15302/J-FASE-2023508>
- de Los Santos, B., Medina, J. J., Miranda, L., Gómez, J. A., & Talavera, M. (2021). Soil disinfestation efficacy against soil fungal pathogens in strawberry crops in Spain: An overview. *Agronomy*, 11(3), 526. <https://doi.org/10.3390/agronomy11030526>
- Demirsoy, L., Misir, D., Bektaş, A., Soysal, D., Lizalo, A., & Demirsoy, H. (2024). Influence of high tunnel conditions and cultivars on the growth, yield, and fruit quality of strawberry in northern Türkiye. *Turkish Journal of Agriculture and Forestry*, 48(1), 57–70. <https://doi.org/10.55730/1300-011X.3162>
- Deschenes, O., & Kolstad, C. (2011). Economic impacts of climate change on California agriculture. *Climatic Change*, 109(S1), 365–386. <https://doi.org/10.1007/s10584-011-0322-3>
- de Tommaso, N., López Aranda, J. M., Greco, N., Saporiti, M., Maccarini, C., & Myrta, A. (2021). Sustainability of strawberry nurseries and fruit production in relation to fumigation practices in Europe. *Acta Horticulturae*, 1309, 693–700. <https://doi.org/10.17660/ActaHortic.2021.1309.100>
- Dias, C. N., Marinho, A. B., Arruda, R. S., Silva, E. J. P., Pereira, E. D., & Femande, C. N. V. (2015). Productivity and quality of strawberry under different environments and biofertilizer doses. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 19(10), 961–966. <https://doi.org/10.1590/1807-1929/agriambi.v19n10p961-966>
- Díaz-Pérez, J. C., Bag, S., Coolong, T., Luo, X., Hodges, A., Bashyal, M., Milner, H., Konakalla, N. C., & Pitcher, A. (2024). Plant growth, fruit yield, and tomato leaf curl disease of high tunnel organic tomato affected by shade net and plastic mulch color.

- HortScience*, 59(3), 323–331. <https://doi.org/10.21273/HORTSCI17516-23>
- DiGiacomo, G., Gieske, M., Grossman, J., Jacobsen, K., Peterson, H., & Rivard, C. (2023). Economic trade-offs: Analysis of hairy vetch (*Vicia villosa*) cover crop use in organic tomato (*Solanum lycopersicum* L.) high tunnel systems across multiple regions. *Renewable Agriculture and Food Systems*, 38, e10. <https://doi.org/10.1017/S1742170523000029>
- Divya, V. U., & Sarkar, N. C. (2019). Plastic mulch pollution and introduction of biodegradable plastic mulches – a review. *Agricultural Reviews*, 40(4), 314–318. <https://doi.org/10.18805/ag.R-1913>
- Domínguez, P., Medina, J. J., Miranda, L., López-Aranda, J. M., Ariza, M. T., Soria, C., Santos, B. M., Torres-Quezada, E. A., & Hernández-Ochoa, I. (2016). Effect of planting and harvesting dates on strawberry fruit quality under high tunnels. *International Journal of Fruit Science*, 16(Sup.1), 228–238. <https://doi.org/10.1080/15538362.2016.1219291>
- Dong, J., Xu, Y., Quan, Z., Yin, Y., Zhao, Y., Xu, Q., Tian, K., Huang, B., Cai, Z., Ma, Y., & Duan, Z. (2024). The obstacles and countermeasures of soil sustainability in protected horticulture in China. *Acta Pedologica Sinica*, 61(6), 1467–1480. <https://doi.org/10.11766/trxb202311010449>
- Egea, F. J., & Glass, R. (2017). Almería: A model for sustainable intensive production. *Aspects of Applied Biology*, 136, 233–236. <https://doi.org/10.1016/j.nbt.2017.06.009>
- Egea, F. J., Torrente, R. G., & Aguilar, A. (2018). An efficient agro-industrial complex in Almería (Spain): Towards an integrated and sustainable bioeconomy model. *New Biotechnology*, 40, 103–112. <https://doi.org/10.1016/j.nbt.2017.06.009>
- Ernst, M. (2020). *Low tunnel economics*. College of Agriculture, Food and Environment. University of Kentucky.
- Evans, N. (2013). Strawberry fields forever? Conflict over neo-productivist Spanish polytunnel technology in British agriculture. *Land Use Policy*, 35, 61–72. <https://doi.org/10.1016/j.landusepol.2013.04.019>
- Evenhuis, A., & Wanten, P. J. (2006). Effect of polythene tunnels and cultivars on grey mould caused by *Botrytis cinerea* in organically grown strawberries. *Agriculturae Conspectus Scientificus*, 71(4), 111–114.
- Fan, Z., Verma, S., Lee, H., Jang, Y. J., Wang, Y., Lee, S., & Whitaker, V. M. (2024). Strawberry soluble solids QTL with inverse effects on yield. *Horticulture Research*, 11(2), uhad271 <https://doi.org/10.1093/hr/uhad271>
- FAOSTAT. (n.d.). <https://www.fao.org/faostat/en/#home>
- Fatnassi, H., Leyronas, C., Boulard, T., Bardin, M., & Nicot, P. (2009). Dependence of greenhouse tunnel ventilation on wind direction and crop height. *Biosystems Engineering*, 103(3), 338–343. <https://doi.org/10.1016/j.biosystemseng.2009.03.005>
- Faure, J., Mouysset, L., Allier, F., Decourtye, A., & Gaba, S. (2024). How pollinator dependence may mediate farmer adoption of pollinator supporting practices and perceptions: A case study from western France. *Environmental Research Communications*, 6(9), 095010. <https://doi.org/10.1088/2515-7620/ad75ee>
- Feldmann, M. J., Pincot, D. D. A., Cole, G. S., & Knapp, S. J. (2024). Genetic gains underpinning a little-known strawberry green revolution. *Nature Communications*, 15(1), 2468. <https://doi.org/10.1038/s41467-024-46421-6>
- Feltham, H., Park, K., Minderman, J., & Goulson, D. (2015). Experimental evidence that wildflower strips increase pollinator visits to crops. *Ecology and Evolution*, 5(16), 3523–3530. <https://doi.org/10.1002/ece3.1444>
- Fenton, M., Goodrich, B. K., & Penn, J. (2025). Measuring beekeepers' economic value of contract enhancements in almond pollination agreements. *Ecological Economics*, 227, 108351. <https://doi.org/10.1016/j.ecolecon.2024.108351>
- Fernandes de Oliveira, A., & Nieddu, G. (2015). Vine growth and physiological performance of two red grape cultivars under natural and reduced UV solar radiation. *Australian Journal of Grape and Wine Research*, 22(1), 105–114. <https://doi.org/10.1111/ajgw.12179>
- Fernandez, G. E. (2001). Fall-applied row covers enhance yield in plasticulture strawberries. *HortTechnology*, 11(3), 440–444. <https://doi.org/10.21273/HORTTECH.11.3.440>
- Feuerbacher, A., Herbold, T., & Krumbe, F. (2024). The economic value of pollination services for seed production: A blind spot deserving attention. *Environmental and Resource Economics*, 87(4), 881–905. <https://doi.org/10.1007/s10640-024-00840-7>
- Filippi, D., Nienow, A. A., Chiomento, J. L. T., dos Santos Trentin, T., Dornelles, A. G., Calvete, E. O., & Huzar-Novakowski, J. (2021). Development and validation of a set of standard area diagrams to assess severity of gray mold in strawberry fruit. *European Journal of Plant Pathology*, 160(2), 277–286. <https://doi.org/10.1007/s10658-021-02238-3>
- Fletcher, J. M., Tatsiopoulou, A., Hadley, P., Davis, F. J., & Henbest, R. G. C. (2004). Growth, yield and development of strawberry cv. 'Elsanta' under novel photosensitive film clad greenhouses. *Acta Horticulturae*, 633, 99–106. <https://doi.org/10.17660/ActaHortic.2004.633.11>
- Fountain, M. T., Delgado, A., Deakin, G., Davis, F., & Hemer, S. (2023). Light spectra blocking films reduce numbers of western flower thrips, *Frankliniella occidentalis* (Thysanoptera: Thripidae) in strawberry, *Fragaria x ananassa*. *Agricultural and Forest Entomology*, 25(1), 1–8. <https://doi.org/10.1111/afe.12526>
- Gaisser, R., Kuehn, K., & Pritts, M. (2024). Novel low tunnel coverings and plant type affect productivity of day-neutral strawberries. *HortTechnology*, 34(3), 381–387. <https://doi.org/10.21273/HORTTECH05409-24>
- Galafton, C., Maga, D., Sonnemann, G., & Thonemann, N. (2023). Life cycle assessment of different strawberry production methods in Germany with a particular focus on plastic emissions. *International Journal of Life Cycle Assessment*, 28(6), 611–625. <https://doi.org/10.1007/s11367-023-02167-9>
- Galinato, S. P., & Miles, C. A. (2013). Economic profitability of growing lettuce and tomato in Western Washington under high tunnel and open-field production system. *HortTechnology*, 23(4), 453–461. <https://doi.org/10.21273/HORTTECH.23.4.453>
- Gallego-Tévar, B., Gil-Martínez, M., Perea, A., Pérez-Ramos, I., & Gómez-Aparicio, L. (2024). Interactive effects of climate change and pathogens on plant performance: A global meta-analysis. *Global Change Biology*, 30(10), e17535. <https://doi.org/10.1111/gcb.17535>
- Gama, A. B., Cordova, L. G., Baggio, J. S., Mertely, J. C., & Peres, N. A. (2023). Old but gold: Captan is a valuable tool for managing anthracnose and botrytis fruit rots and improving strawberry yields based on a meta-analysis. *Plant Disease*, 107(10), 3071–3078. <https://doi.org/10.1094/PDIS-12-22-2781-RE>
- Garcia, M. E., Ernst, T., Johnson, D. T., & Dickey, D. A. (2017). Strawberry cultivar performance in high tunnels under sustainable and organic production practices in

- three climatic regions of Arkansas. *Acta Horticulturae*, 1156, 549–554. <https://doi.org/10.17660/ActaHortic.2017.1156.81>
- García-Tejero, I. F., López-Borrillo, D., Miranda, L., Medina, J. J., Arriaga, J., Muriel-Fernández, J. L., & Martínez-Ferri, E. (2018). Estimating strawberry crop coefficients under plastic tunnels in southern Spain by using drainage lysimeters. *Scientia Horticulturae*, 231, 233–240. <https://doi.org/10.1016/j.scienta.2017.12.020>
- Gázquez, J. C., López, J. C., Pérez-Parra, J. J., Baeza, E. J., Saéz, M., & Parra, A. (2008). Greenhouse cooling strategies for Mediterranean climate areas. *Acta Horticulturae*, 801, 425–432. <https://doi.org/10.17660/ActaHortic.2008.801.45>
- Gent, M. P. N. (1992). Effect of planting date, ventilation and soil temperature on growth and nutrition of tomato in high tunnels. *Plant and Soil*, 145(1), 81–91. <https://doi.org/10.1007/BF00009544>
- Giacomelli, G. A. (2009). Engineering principles impacting high-tunnel environments. *HortTechnology*, 19(1), 30–33. <https://doi.org/10.21273/HORTTECH.19.1.30>
- Greco, N., Aranda, J. M. L., Saporiti, M., Maccarini, C., de Tommaso, N., & Myrta, A. (2020). Sustainability of European vegetable and strawberry production in relation to fumigation practices in the EU. *Acta Horticulturae*, 1270, 203–210. <https://doi.org/10.17660/ActaHortic.2020.1270.24>
- Grisey, A., Garnodier, J., Stauffer, V., & Masson, A. (2023). Increasing night temperature in a tunnel without heating system. *Acta Horticulturae*, 1377, 149–156. <https://doi.org/10.17660/ActaHortic.2023.1377.18>
- Gruda, N., Bisbis, M., Katsoulas, N., & Kittas, C. (2021). Smart greenhouse production practices to manage and mitigate the impact of climate change in protected cultivation. *Acta Horticulturae*, 1320, 189–196. <https://doi.org/10.17660/ActaHortic.2021.1320.24>
- Gruda, N., Bisbis, M., & Tanny, J. (2019). Influence of climate change on protected cultivation: Impacts and sustainable adaptation strategies - a review. *Journal of Cleaner Production*, 225, 481–495. <https://doi.org/10.1016/j.jclepro.2019.03.210>
- Gruda, N. S., Hirschler, O., & Stuart, J. (2024). Peat reduction in horticulture – an overview of Europe. *Acta Horticulturae*, 1391, 545–560. <https://doi.org/10.17660/ActaHortic.2024.1391.75>
- Gu, S. (2021). *High tunnel farming*. Cooperative Extension. North Carolina Agricultural and Technical State University.
- Gude, K. M., Pliakoni, E. D., Cunningham, B., Ayub, K., Kang, Q., Rajashekar, C. B., & Rivard, C. L. (2022). High tunnel coverings alter crop productivity and microclimate of tomato and lettuce. *HortScience*, 57(2), 265–272. <https://doi.org/10.21273/HORTSCI.16208-21>
- Gudowska, A., Cwajna, A., Marjańska, E., & Moroń, D. (2024). Pollinators enhance the production of a superior strawberry – a global review and meta-analysis. *Agriculture, Ecosystems & Environment*, 362, 108815. <https://doi.org/10.1016/j.agee.2023.108815>
- Guéry, S., Lea-Cox, J. D., Martínez Bastida, M. A., Belayneh, B. E., & Ferrer-Alegre, F. (2018). Using sensor-based control to optimize soil moisture availability and minimize leaching in commercial strawberry production in Spain. *Acta horticulturae*, 1197, 171–178. <https://doi.org/10.17660/ActaHortic.2018.1197.23>
- Gullickson, M. G., DiGiacomo, G., & Rogers, M. A. (2024). Efficacy and economic viability of organic control methods for spotted-wing drosophila in day-neutral strawberry production in the Upper Midwest. *HortTechnology*, 34(5), 618–628. <https://doi.org/10.21273/HORTTECH05461-24>
- Gupta, J., Sharma, D., & Chauhan, A. (2024). Classification and suitability of protected structures. In M. C. Singh & K. K. Sharma (Eds.), *Protected cultivation: Structural design, crop management, modeling and automation* (pp. 35–61). Apple Academic Press Incorporated.
- Guttormsen, G. (1972). The effect of perforation on temperature conditions in plastic tunnels. *Journal of Agricultural Engineering Research*, 17(2), 172–177. [https://doi.org/10.1016/S0021-8634\(72\)80005-2](https://doi.org/10.1016/S0021-8634(72)80005-2)
- Hall, M. A., Jones, J., Rocchetti, M., Wright, D., Rader, R., & Strange, J. (2020). Bee visitation and fruit quality in berries under protected cropping vary along the length of polytunnels. *Journal of Economic Entomology*, 113(3), 1337–1346. <https://doi.org/10.1093/jee/toaa037>
- Hancock, J., & Simpson, D. (1995). Methods of extending the strawberry season in Europe. *HortTechnology*, 5(4), 286–290. <https://doi.org/10.21273/HORTTECH.5.4.286>
- Hanson, E., Von Weihe, M., Schilder, A. C., Chanon, A. M., & Scheerens, J. C. (2011). High tunnel and open field production of florican- and primocane-fruited raspberry cultivars. *HortTechnology*, 21(4), 412–418. <https://doi.org/10.21273/HORTTECH.21.4.412>
- He, J., Lin, T., Liang, W., Chavan, S., Sethuvenkatraman, S., Goldsworthy, M., Tissue, D., & Chen, Z.-H. (2024). Energy and fertigation consumption and economic viability of a light blocking film in greenhouse lettuce production. *Journal of Cleaner Production*, 479, 144013. <https://doi.org/10.1016/j.jclepro.2024.144013>
- He, M.-H., & Li, K.-T. (2014). The plastic tunnel system reduced irradiance, and whole plant photosynthesis and vegetative growth in strawberry cultivation. *Journal of the Taiwan Society for Horticultural Science*, 60(3), 41–49.
- Henschel, J. M., de Resende, J. T. V., Zeist, A. R., dos Santos, R. L., de Lima, V. A., Giloni-Lima, P. C., & Batista, D. S. (2024). True colors shining through: How low tunnel cover colors affect fruit yield and photosynthesis in strawberry cultivars. *Vegetos*. <https://doi.org/10.1007/s42535-024-00985-2>
- Hernández, A. F., Requena, M., López, A., Parrón, T., Navarro, A., & Alarcón, R. (2020). Reply to Swaen's letter regarding 'Environmental exposure to pesticides and risk of thyroid diseases'. *Toxicology Letters*, 331, 254–256. <https://doi.org/10.1016/j.toxlet.2020.05.039>
- Hernandez, I. M., Santos, B. M., Stanley, C. D., & Zhao, X. (2016). Comparison of freeze protection methods for strawberry production in Florida. *International Journal of Fruit Science*, 16(1), 103–112. <https://doi.org/10.1080/15538362.2015.1079515>
- Hernández-Martínez, N. R., Blanchard, C., Wells, D., & Salazar-Gutiérrez, M. R. (2023). Current state and future perspectives of commercial strawberry production: A review. *Scientia Horticulturae*, 312, 111893. <https://doi.org/10.1016/j.scienta.2023.111893>
- Herrington, M. E., Hardner, C., Wegener, M., Woolcock, L. L., & Dieters, M. J. (2011). Rain damage to strawberries grown in southeast Queensland: Evaluation and genetic control. *HortScience*, 46(6), 832–837. <https://doi.org/10.21273/HORTSCI.46.6.832>
- Ho, S.-T., Ifft, J. E., Rickard, B. J., & Turvey, C. G. (2018). Alternative strategies to manage weather risk in perennial fruit crop production. *Agricultural and Resource Economics Review*, 47(3), 452–476. <https://doi.org/10.1017/age.2017.29>

- Hodgdon, E. A., Conner, D. S., McDermott, L. G., Pritts, M. P., Handley, D. T., Orde, K. M., & Grube Sideman, R. (2024). A current view on strawberry production practices and trends in the Northeastern United States and Canada. *HortTechnology*, 34(5), 574–584. <https://doi.org/10.21273/HORTTECH05457-24>
- Hodgkiss, D., Brown, M. J. F., & Fountain, M. T. (2019). The effect of within-crop floral resources on pollination, aphid control and fruit quality in commercial strawberry. *Agriculture, Ecosystems & Environment*, 275, 112–122. <https://doi.org/10.1016/j.agee.2019.02.006>
- Holmes, G. J. (2024). The California strawberry industry: Current trends and future prospects. *International Journal of Fruit Science*, 24(1), 115–129. <https://doi.org/10.1080/15538362.2024.2342900>
- Howard, S. R., Nisal Ratnayake, M., Dyer, A. G., Garcia, J. E., Dorin, A., & Smagghe, G. (2021). Towards precision apiculture: Traditional and technological insect monitoring methods in strawberry and raspberry crop polytunnels tell different pollination stories. *PLoS One*, 16(5), e0251572. <https://doi.org/10.1371/journal.pone.0251572>
- Hull, K., van Schalkwyk, P. D., Mabitsela, M., Phiri, E. E., & Booyesen, M. J. (2024). Modelling the temperature inside a greenhouse tunnel. *AgriEngineering*, 6(1), 285–301. <https://doi.org/10.3390/agriengineering6010017>
- Hurtado, G., & Knoche, M. (2021). Water soaking disorder in strawberries: Triggers, factors, and mechanisms. *Frontiers in Plant Science*, 12, 694123. <https://doi.org/10.3389/fpls.2021.694123>
- Hurtado, G., Olbricht, K., Mercado, J. A., Pose, S., & Knoche, M. (2024). Phenotyping 172 strawberry genotypes for water soaking reveals a close relationship with skin water permeance. *PeerJ*, 12, e17960. <https://doi.org/10.7717/peerj.17960>
- Isoyama, Y., Nakashima, M., Kaneko, M., & Kiyono, K. (2024). Evaluation of heat stress of workers using wearable biosensors in protected horticulture. *Acta Horticulturae*, 1404, 259–268. <https://doi.org/10.17660/ActaHortic.2024.1404.36>
- Jamarkattel, D., Tuladhar, F., Jamir, C., & Diwakar, K. C. (2023). Tunnel farming as an adaptation tool against climate change effect among smallholder farmers in Nepal. In S. A. Narula & S. P. Raj (Eds.), *Sustainable food value chain development* (pp. 153–174). Springer Nature.
- Janke, R. R., Altamimi, M. E., & Khan, M. (2017). The use of high tunnels to produce fruit and vegetable crops in North America. *Agricultural Sciences*, 8(7), 692–715. <https://doi.org/10.4236/as.2017.87052>
- Jayalath, T. C., Boyhan, G. E., Little, E. L., Tate, R. I., & O'Connell, S. (2017). High tunnel and field system comparison for spring organic lettuce production in Georgia. *HortScience*, 52(11), 1518–1524. <https://doi.org/10.21273/HORTSCI12284-17>
- Jayasuriya, N., Guo, Y., Hu, W., & Ghannoum, O. (2024). Image based crop monitoring technologies in protected horticulture: A review (unpublished) <https://doi.org/10.48550/arXiv.2401.13928>
- Jiang, W. J., & Yu, H. J. (2008). Present situation and future development for protected horticulture in mainland China. *Acta Horticulturae*, 770, 29–35. <https://doi.org/10.17660/ActaHortic.2008.770.3>
- Jiménez-Lao, C., García-Caparrós, P., Lao, M. T., Llanderal, A., & Franco-Rodríguez, J. (2021). Protected horticulture of Ecuador: Past, present and future perspectives. In G. C. Lopez & M. B. A. Calle (Eds.), *Ecuador: Perspectives of the past, present and future: A multi-criteria approach to social evolution* (pp. 179–230). Nova Science Publishers.
- Jirgena, H., Hazners, J., Kaufmane, E., Strautina, S., Feldmane, D., & Skrivele, M. (2013). Risks and returns in strawberry, raspberry and cherry production with various methods. *Economics and Rural Development*, 9(2), 16–26.
- Johnson, J., & Hoffmann, M. (2024). Evaluation of yield, fruit chemistry, and firmness of seven strawberry (*Fragaria ×ananassa*) cultivars in an eastern North Carolina greenhouse. *HortScience*, 59(11), 1634–1643. <https://doi.org/10.21273/HORTSCI18084-24>
- Jung, W., & Kim, H. (2022). Evaluation of heat stress levels inside greenhouses during summer in Korea. *International Journal of Environmental Research and Public Health*, 19(19), 12497. <https://doi.org/10.3390/ijerph191912497>
- Kaiser, E., Kusuma, P., Vialet-Chabrand, S., Folta, K., Liu, Y., Poorter, H., Woning, N., Shrestha, S., Ciarreta, A., van Brenk, J., Karpe, M., Ji, Y., David, S., Zepeda, C., Zhu, X.-G., Huntenburg, K., Verdonk, J. C., Woltering, E. ... Marcelis, L. F. M. (2024). Vertical farming goes dynamic: Optimizing resource use efficiency, product quality, and energy costs. *Frontiers in Science*, 2, 1411259. <https://doi.org/10.3389/fsci.2024.1411259>
- Kapanen, A., Schettini, E., Vox, G., & Itävaara, M. (2008). Performance and environmental impact of biodegradable films in agriculture: A field study on protected cultivation. *Journal of Polymers and the Environment*, 16(2), 109–122. <https://doi.org/10.1007/s10924-008-0091-x>
- Kasirajan, S., & Ngouajio, M. (2012). Polyethylene and biodegradable mulches for agricultural applications: A review. *Agronomy and Sustainable Development*, 32(2), 501–529. <https://doi.org/10.1007/s13593-011-0068-3>
- Kendall, L. K., Evans, L. J., Gee, M., Smith, T. J., Gagic, V., Lobaton, J. D., Hall, M. A., Jones, J., Kirkland, L., Saunders, M. E., Sonter, C., Cutting, B. T., Parks, S., Hogendoorn, K., Spurr, C., Gracie, A., Simpson, M., & Rader, R. (2021). The effect of protective covers on pollinator health and pollination service delivery. *Agriculture, Ecosystems & Environment*, 319, 107556. <https://doi.org/10.1016/j.agee.2021.107556>
- Kennedy, C., Hasing, T. N., Peres, N. A., & Whitaker, V. M. (2013). Evaluation of strawberry species and cultivars for powdery mildew resistance in open-field and high tunnel production systems. *HortScience*, 48(9), 1125–1129. <https://doi.org/10.21273/HORTSCI.48.9.1125>
- Khoshnevisan, B., Rafiee, S., & Mousazadeh, H. (2013). Environmental impact assessment of open field and greenhouse strawberry production. *European Journal of Agronomy*, 50, 29–37. <https://doi.org/10.1016/j.eja.2013.05.003>
- Kim, H.-K. (2024). Evaluating the ventilation performance of single-span plastic greenhouses with continuous screened side openings. *Agronomy*, 14(7), 1447. <https://doi.org/10.3390/agronomy14071447>
- Kirschbaum, D. S., Quiroga, R. J., Funes, C. F., & Villagra, E. L. (2024). Strawberry cultivars performance in contrasting cropping conditions in Tucumán (Argentina). *Revista Agronómica del Noroeste Argentino*, 43(1), 26–34. <https://doi.org/10.61914/ranar.4301.02>
- Kittas, C., Karamanis, M., & Katsoulas, N. (2005). Air temperature regime in a forced ventilated greenhouse with rose crop. *Energy and Buildings*, 37(8), 807–812. <https://doi.org/10.1016/j.enbuild.2004.10.009>
- Knapp, S. J., Cole, G. S., Pincot, D. D., López, C. M., Gonzalez-Benitez, O. A., & Famula, R. A. (2023). ‘UC

- Eclipse', a summer plant-adapted photoperiod-insensitive strawberry cultivar. *HortScience*, 58(12), 1568–1572. <https://doi.org/10.21273/HORTSCI17363-23>.
- Kotilainen, T., Robson, T. M., Hernández, R., & Niedz, R. P. (2018). Light quality characterization under climate screens and shade nets for controlled-environment agriculture. *PLoS One*, 13(6), e0199628. <https://doi.org/10.1371/journal.pone.0199628>
- Kulak, M., Graves, A., & Chatterton, J. (2013). Reducing greenhouse gas emissions with urban agriculture: A life cycle assessment perspective. *Landscape and Urban Planning*, 111, 68–78. <https://doi.org/10.1016/j.landurbplan.2012.11.007>
- Lakhari, I. A., Yan, H., Zhang, J., Wang, G., Deng, S., Bao, R., Zhang, C., Syed, T. N., Wang, B., Zhou, R., & Wang, X. (2024). Plastic pollution in agriculture as a threat to food security, the ecosystem, and the environment: An overview. *Agronomy*, 14(3), 548. <https://doi.org/10.3390/agronomy1403054>.
- Lang, G. A., Sage, L., & Wilkinson, T. (2016). Ten Years of studies on systems to modify sweet cherry production environments: Retractable roofs, high tunnels, and rain-shelters. *Acta Horticulturae*, 1130, 83–90. <https://doi.org/10.17660/ActaHortic.2016.1130.12>
- Layek, U., Baghira, N. K., Das, A., Kundu, A., & Karmakar, P. (2023). Dependency of crops on pollinators and pollination deficits: An approach to measurement considering the influence of various reproductive traits. *Agriculture*, 13(8), 1563. <https://doi.org/10.3390/agriculture13081563>
- Leach, H., & Isaacs, R. (2018). Seasonal occurrence of key arthropod pests and beneficial insects in Michigan high tunnel and field grown raspberries. *Environmental Entomology*, 47(3), 567–574. <https://doi.org/10.1093/ee/nyy030>.
- Leskovar, D. I., Choi, S., Harvey, J. T., Lee, C., Nagila, A., Niu, G., Masabni, J., Zahid, A., & Dash, P. K. (2024). Controlled environment agriculture trends in Texas: Challenges, opportunities, and research approaches. *Acta Horticulturae*, 1391, 197–204. <https://doi.org/10.17660/ActaHortic.2024.1391.27>
- Lewers, K. S., Fleisher, D. H., & Daughtry, C. S. T. (2017). Low tunnels as a strawberry breeding tool and season-extending production system. *International Journal of Fruit Science*, 17(3), 233–258. <https://doi.org/10.1080/15538362.2017.1305941>
- Lewers, K. S., Fleisher, D. H., Daughtry, C. S. T., & Vinyard, B. T. (2020). Low-tunnel strawberry production: Comparison of cultivars and films. *International Journal of Fruit Science*, 20(Sup. 2), S705–S732. <https://doi.org/10.1080/15538362.2020.1768616>
- Lewus, D., & Both, A. J. (2020). Using computational fluid dynamics (CFD) to improve high tunnel ventilation. *Acta Horticulturae*, 1296, 33–40. <https://doi.org/10.17660/ActaHortic.2020.1296.5>
- Lewus, D. C., & Both, A. J. (2022). Using computational fluid dynamics to evaluate high tunnel roof vent designs. *AgriEngineering*, 4(3), 719–732. <https://doi.org/10.3390/agriengineering4030046>
- Lima, J. M. D., Fagherazzi, A. F., Nerbass, F. R., Petry, D., Kretschmar, A. A., Baruzzi, G., Rufato, L., & Bogo, A. (2024). Epidemiology of mycosphaerella leaf spot and powdery mildew and agronomic parameters of strawberry cultivars and genotypes in the Highland Region of Southern Brazil. *Agriculture*, 14(8), 1373. <https://doi.org/10.3390/agriculture14081373>
- Lin, T., Maier, C. R., Liang, W., Klause, N., He, J., Tissue, D. T., Lan, Y.-C., Sethuvenkatraman, S., Goldsworthy, M., & Chen, Z.-H. (2024). A light-blocking greenhouse film differentially impacts climate control energy use and capsicum production. *Frontiers in Energy Research*, 12, 1360536. <https://doi.org/10.3389/fenrg.2024.1360536>.
- Liu, D., Samtani, J. B., Taghavi, T., & Amyotte, B. (2024). Agronomic and post-harvest performance of strawberry cultivars in high tunnel and open-field environment in southeast Virginia. *International Journal of Fruit Science*, 24(1), 242–255. <https://doi.org/10.1080/15538362.2024.2384395>
- Lobell, D. B., & Field, C. B. (2011). California perennial crops in a changing climate. *Climatic Change*, 109(S1), 317–333. <https://doi.org/10.1007/s10584-011-0303-6>
- Logan, H., Astrup, T. F., & Damgaard, A. (2024a). Additive inclusion in plastic life cycle assessments part I: Review of mechanical recycling studies. *Journal of Industrial Ecology*, 28(6), 1582–1597. <https://doi.org/10.1111/jiec.13542>
- Logan, H., DeMeester, S., Astrup, T. F., & Damgaard, A. (2024b). Additive inclusion in plastic life cycle assessments, part II: Review of additive inventory data trends and availability. *Journal of Industrial Ecology*, 28(6), 1554–1566. <https://doi.org/10.1111/jiec.13534>
- López, M. E., Denoyes, B., & Bucher, E. (2024). Epigenomic and transcriptomic persistence of heat stress memory in strawberry (*Fragaria vesca*). *BMC Plant Biology*, 24, 405. <https://doi.org/10.1186/s12870-024-05093-6>.
- Lozano, D., Ruiz, N., & Gavilán, P. (2016). Consumptive water use and irrigation performance of strawberries. *Agricultural Water Management*, 169, 44–51. <https://doi.org/10.1016/j.agwat.2016.02.011>
- Lubna, F. A., Lewus, D. C., Shelford, T. J., & Both, A.-J. (2022). What you may not realize about vertical farming. *Horticulturae*, 8(4), 322. <https://doi.org/10.3390/horticulturae8040322>
- Luo, H., Guan, Y., Zhang, Z., Zhang, Z. I. H., Zhang, Z. H., & Li, H. (2024). *FveDREB1B* improves cold tolerance of woodland strawberry by positively regulating *FveSCL23* and *FveCHS*. *Plant, Cell & Environment*, 47(12), 4630–4650. <https://doi.org/10.1111/pce.15052>
- Lynn, S. C., Dunwell, J. M., Whitehouse, A. B., & Cockerton, H. M. (2024). Genetic loci associated with tissue-specific resistance to powdery mildew in octoploid strawberry (*Fragaria × ananassa*). *Frontiers in Plant Science*, 15, 1376061. <https://doi.org/10.3389/fpls.2024.1376061>
- Ma, Y., Yang, M., Zhao, Q., Li, D., & Liu, D. (2024). Assessing the effects of mechanical damage on optical properties of strawberries in the 950–1650 nm range. *Postharvest Biology and Technology*, 218, 113145. <https://doi.org/10.1016/j.postharvbio.2024.113145>.
- Maboko, M. M., Du Plooy, C. P., & Bertling, I. (2012). Performance of tomato cultivars in temperature and non-temperature controlled plastic tunnels. *Acta Horticulturae*, 927, 405–411. <https://doi.org/10.17660/ActaHortic.2012.927.50>
- Maier, C. R., Chavan, S. G., Klause, N., Liang, W., Cazzonelli, C. I., Ghannoum, O., Chen, Z.-H., & Tissue, D. T. (2023). Light blocking film in a glasshouse impacts *Capsicum annuum* L. yield differentially across planting season. *Frontiers in Plant Science*, 14, 1277037. <https://doi.org/10.3389/fpls.2023.1277037>
- Maier, C. R., Chen, Z.-H., Cazzonelli, C. I., Tissue, D. T., & Ghannoum, O. (2022). Precise phenotyping for improved crop quality and management in protected cropping: A review. *Crops*, 2(4), 336–350. <https://doi.org/10.3390/crops2040024>
- Makowski, H., Lamb, K., & Galloway, L. F. (2024). Support for Baker's law: Facultative self-fertilization ability

- decreases pollen limitation in experimental colonization. *American Journal of Botany*, 111(6), e16351. <https://doi.org/10.1002/ajb2.16351>
- Marin, H., & Garcia, C. R. (2004). Income distribution of protected-crop farms in Andalusia. *Acta Horticulturae*, 659, 47–52. <https://doi.org/10.17660/ActaHortic.2004.659.4>
- Martínez, F., Oliveira, J. A., Calvete, E. O., & Palencia, P. (2017). Influence of growth medium on yield, quality indexes and SPAD values in strawberry plants. *Scientia Horticulturae*, 217, 17–27. <https://doi.org/10.1016/j.scienta.2017.01.024>
- Martínez-Ferri, E., Ariza, M. T., Domínguez, P., Medina, J. J., Miranda, L., Muriel, J. L., Montesinos, P., Rodríguez-Díaz, J. A., & Soria, C. (2014). Cropping strawberry for improving productivity and environmental sustainability. In N. Malone (Ed.), *Strawberries: Cultivation, antioxidant properties and health benefits* (pp. 1–20). Nova Science Publishers.
- Martínez-Ferri, E., Soria, C., Ariza, M. T., Medina, J. J., Miranda, L., Domínguez, P., & Muriel, J. L. (2016). Water relations, growth and physiological response of seven strawberry cultivars (*Fragaria* × *ananassa* Duch.) to different water availability. *Agricultural Water Management*, 164, 73–82. <https://doi.org/10.1016/j.agwat.2015.08.014>
- Matamala, M. F., Bastías, R. M., Urrea, I., Calderón-Orellana, A., Campos, J., & Albornoz, K. (2023). Rain cover and netting materials differentially affect fruit yield and quality traits in two highbush blueberry cultivars via changes in sunlight and temperature conditions. *Plants*, 12(20), 3556. <https://doi.org/10.3390/plants12203556>
- Mateos-Fierro, Z., Garratt, M. P. D., Fountain, M. T., Ashbrook, K., & Westbury, D. B. (2023). The potential of wildflower strips to enhance pollination services in sweet cherry orchards grown under polytunnels. *Journal of Applied Ecology*, 60(6), 1044–1055. <https://doi.org/10.1111/1365-2664.14394>
- Mateos-Fierro, Z., Garratt, M. P. D., Fountain, M. T., Ashbrook, K., & Westbury, D. B. (2024). Wildflower strips in polytunnel cherry orchard alleyways support pest regulation services but do not counteract edge effects on pollination services. *Frontiers in Sustainable Food Systems*, 8, 1423511. <https://doi.org/10.3389/fsufs.2024.1423511>
- Matsui, H., & Mochida, K. (2024). Functional data analysis-based yield modeling in year-round crop cultivation. *Horticulture Research*, 11(7), uhae144. <https://doi.org/10.1093/hr/uhae144>
- Maughan, T. L., Curtis, K. R., Black, B. L., & Drost, D. T. (2015). Economic evaluation of implementing strawberry season extension production technologies in the U.S Intermountain West. *HortScience*, 50(3), 395–401. <https://doi.org/10.21273/HORTSCI.50.3.395>
- Maynard, E., & O'Donnell, M. (2019). *Managing the environment in high tunnels for cool season vegetable production*. Purdue University.
- Mbarushimana, J. C., Bosch, D. J., & Samtani, J. B. (2022). An economic comparison of high tunnel and open-field strawberry production in southeastern Virginia. *Horticulturae*, 8(12), 1139. <https://doi.org/10.3390/horticulturae8121139>
- McIntosh, K. (2018). *Transitioning to protected cropping*. www.strawberryinnovation.com.au
- Meland, M., Frøynes, O., & Kaiser, C. (2017). High tunnel production systems improve yields and fruit size of sweet cherry. *Acta Horticulturae*, 1161, 117–124. <https://doi.org/10.17660/ActaHortic.2017.1161.20>
- Mello, P. P., Onofre, R. B., Rea, M., Bierman, A., Gadoury, D. M., Ivors, K., Ganci, M., Broome, J. C., & Peres, N. A. (2022). Design, construction, and evaluation of equipment for nighttime applications of UV-C for management of strawberry powdery mildew in Florida and California. *Plant Health Progress*, 23(3), 321–327. <https://doi.org/10.1094/PHP-01-22-0002-RS>
- Menzel, C. M. (2023). Fruit set is moderately dependent on insect pollinators in strawberry and is limited by the availability of pollen under natural open conditions. *Journal of Horticultural Science and Biotechnology*, 98(6), 685–714. <https://doi.org/10.1080/14620316.2023.2212670>
- Menzel, C. M., Smith, L. A., & Moisaner, J. A. (2014). The productivity of strawberry plants growing under high plastic tunnels in a wet subtropical environment. *HortTechnology*, 24(3), 334–342. <https://doi.org/10.21273/HORTTECH.24.3.334>
- Mishra, K., Stanghellini, C., & Hemming, S. (2023). Technology and materials for passive manipulation of the solar spectrum in greenhouses. *Advanced Sustainable Systems*, 7(5), 2200503. <https://doi.org/10.1002/adsu.202200503>
- Montague, T., Villanueva-Morales, A., Khan, M. A. R., Wallace, R., & Panta, S. (2024). Comparing the accuracy and efficiency of leaf gas exchange measurements of excised and nonexcised strawberry (*Fragaria* × *ananassa* Duch. 'Camino Real') leaves. *HortTechnology*, 34(6), 728–737. <https://doi.org/10.21273/HORTTECH05516-24>
- Morel, K., & Cartau, K. (2023). Adaptation of organic vegetable farmers to climate change: An exploratory study in the Paris region. *Agricultural Systems*, 210, 103703. <https://doi.org/10.1016/j.agry.2023.103703>
- Morris, P., Gu, S., & Jefferson-Moore, K. (2023). The economic viability of tomato production using single versus double-layer high tunnels. *Journal of Food Distribution Research*, 54(1), 52–54.
- Nacko, S., Hall, M., Spooner-Hart, R., Cook, J., Bernauer, O., & Riegler, M. (2022). Cucurbit crops in temperate Australia are visited more by native solitary bees than by stingless bees. *Journal of Apicultural Research*, 61(5), 675–687. <https://doi.org/10.1080/00218839.2022.2110742>
- Nakayama, M., Fujita, S., & Nakazawa, Y. (2024). Effects of daytime LED supplemental lighting on strawberry growth and yield under subtropical climate. *Acta Horticulturae*, 1404, 233–240. <https://doi.org/10.17660/ActaHortic.2024.1404.32>
- Nemali, K. (2022). History of controlled environment horticulture: Greenhouses. *HortScience*, 57(2), 239–246. <https://doi.org/10.21273/HORTSCI16160-21>
- Neri, D., Baruzzi, G., Massetani, F., & Faedi, W. (2012). Strawberry production in forced and protected culture in Europe as a response to climate change. *Canadian Journal of Plant Science*, 92(6), 1021–1036. <https://doi.org/10.4141/cjps2011-276>
- Nes, A., Henriksen, J. K., Serikstad, G. L., & Stensvand, A. (2017). Cultivars and cultivation systems for organic strawberry production in Norway. *Acta Agriculturae Scandinavica, Section B — Soil and Plant Science*, 67(6), 485–491. <https://doi.org/10.1080/09064710.2017.1296490>
- Nestby, R., Bjørgum, R., Nes, A., Wikdahl, T., & Hageberg, B. (2001). Reactions of strawberry plants to long-term freezing and alternate freezing and thawing. *Journal of Horticultural Science and Biotechnology*, 76(3), 280–285. <https://doi.org/10.1080/14620316.2001.11511364>

- Nestby, R., & Guéry, S. (2017). Balanced fertigation and improved sustainability of June bearing strawberry cultivated three years in open polytunnel. *Journal of Berry Research*, 7(3), 203–216. <https://doi.org/10.3233/JBR-170157>.
- Nian, Y., Zhao, R., Tian, S., Zhao, X., & Gao, Z. (2022). Economic analysis of grafted organic tomato production in high tunnels. *HortTechnology*, 32(5), 459–470. <https://doi.org/10.21273/HORTTECH05101-22>.
- Nishimoto, Y., Lu, N., Ichikawa, Y., Watanabe, A., Kikuchi, M., & Takagaki, M. (2023). An evaluation of pollination methods for strawberries cultivated in plant factories: Robot vs hand. *Technology in Horticulture*, 3, 19. <https://doi.org/10.48130/THH-2023-0019>
- Nordey, T., Faye, E., Chailleux, A., Parrot, L., Simon, S., Mlowe, N., & Fernandes, P. (2020). Mitigation of climatic conditions and pest protection provided by insect-proof nets for cabbage cultivation in East Africa. *Experimental Agriculture*, 56(4), 608–619. <https://doi.org/10.1017/S0014479720000186>
- Nordey, T., Ochieng, J., Ernest, Z., Mlowe, N., Mosha, I., & Fernandes, P. (2020). Is vegetable cultivation under low tunnels a profitable alternative to pesticide use? The case of cabbage cultivation in northern Tanzania. *Crop Protection*, 134, 105169. <https://doi.org/10.1016/j.cropro.2020.105169>
- O’Connell, S., Rivard, C., Peet, M. M., Harlow, C., & Louws, F. (2012). High tunnel and field production of organic heirloom tomatoes: Yield, fruit quality, disease, and microclimate. *HortScience*, 47(9), 1283–1290. <https://doi.org/10.21273/HORTSCI.47.9.1283>
- O’Connell, S., & Tate, R. (2017). Winter broccoli and cauliflower under organic high tunnels in a humid, subtropical climate. *HortScience*, 52(11), 1511–1517. <https://doi.org/10.21273/HORTSCI.12291-17>
- Ogden, A. B., Kim, J., Radcliffe, C. A., van Iersel, M. W., Donovan, L. A., & Sugiyama, A. (2011). Leaf and bud temperatures of southern highbush blueberries (*Vaccinium corymbosum*) inside high tunnels. *Acta Horticulturae*, 893, 1319–1325. <https://doi.org/10.17660/ActaHortic.2011.893.155>
- Ogden, A. B., & van Iersel, M. W. (2009). Southern highbush blueberry production in high tunnels: Temperatures, development, yield, and fruit quality during the establishment years. *HortScience*, 44(7), 1850–1856. <https://doi.org/10.21273/HORTSCI.44.7.1850>
- Onofre, R. B., Gadoury, D. M., & Peres, N. A. (2021). High efficacy and low risk of phytotoxicity of sulfur in the suppression of strawberry powdery mildew. *Plant Health Progress*, 22, 101–107. <https://doi.org/10.1094/PHP-04-20-0034-RS>.
- Onofre, R. B., Gadoury, D. M., Stensvand, A., Bierman, A., Rea, M., & Peres, N. A. (2022). UV-transmitting plastics reduce powdery mildew in strawberry tunnel production. *Plant Disease*, 106(9), 2455–2461. <https://doi.org/10.1094/PDIS-10-21-2195-RE>.
- Orde, K. M., & Grube Sideman, R. (2021). Winter survival and second-year spring yields of day-neutral strawberry are influenced by cultivar and presence of low tunnels. *HortTechnology*, 31(1), 77–88. <https://doi.org/10.21273/HORTTECH04734-20>
- Ordidge, M., García-Macías, P., Battey, N. H., Gordon, M. H., John, P., Lovegrove, J. A., Vysini, E., Wagstaffe, A., & Hadley, P. (2012). Development of colour and firmness in strawberry crops is UV light sensitive, but colour is not a good predictor of several quality parameters. *Journal of the Science of Food and Agriculture*, 92(8), 1597–1604. <https://doi.org/10.1002/jsfa.4744>
- Osman, M., Qaryouti, M., Alharbi, S., Alghamdi, B., Al-Soqeer, A., Alharbi, A., Almutairi, K., & Abdelaziz, M. E. (2024). Impact of CO₂ enrichment on growth, yield and fruit quality of F1 hybrid strawberry grown under controlled greenhouse condition. *Horticulturae*, 10(9), 941. <https://doi.org/10.3390/horticulturae10090941>
- Palma, M., Sepúlveda, A., & Yuri, J. A. (2023). Effect of plastic roof and high tunnel on microclimate, physiology, vegetative growth and fruit characteristics of ‘Santina’ sweet cherry. *Scientia Horticulturae*, 317, 112037. <https://doi.org/10.1016/j.scienta.2023.112037>
- Pandey, G., Parks, S., & Thomas, R. G. (2023). Polymer and photo-selective covers on plant and fruit development: A review. *Agronomy Journal*, 115(6), 3074–3091. <https://doi.org/10.1002/agj2.21442>
- Pedrozo, P., Lado, B., Moltini, A. I., Vicente, E., & Lado, J. (2024). Exploration of strawberry fruit quality during harvest season under a semi-forcing culture with plants nursed without chilling. *Plants*, 13(21), 3052. <https://doi.org/10.3390/plants13213052>.
- Peñuelas, J., Savé, R., Marfà, O., & Serrano, L. (1992). Remotely measured canopy temperature of greenhouse strawberries as indicator of water status and yield under mild and very mild water stress conditions. *Agricultural and Forestry Meteorology*, 58(1–2), 63–77. [https://doi.org/10.1016/0168-1923\(92\)90111-G](https://doi.org/10.1016/0168-1923(92)90111-G)
- Pergola, M., Maffia, A., Carlucci, G., Persiani, A., Palese, A. M., Zaccardelli, M., Altieri, G., & Celano, G. (2023). An environmental and economic analysis of strawberry production in southern Italy. *Agriculture*, 13(9), 1705. <https://doi.org/10.3390/agriculture13091705>
- Pierre, J. F., Jacobsen, K. L., Wszelaki, A., Butler, D., Velandia, M., Woods, T., Sideman, R., Grossman, J., Coolong, T., Hoskins, B., da Silva, A. L. B. R., Ginakes, P., Kleinhenz, M., Zhao, X., Rivard, C., & Rudolph, R. E. (2024). Sustaining soil health in high tunnels: A paradigm shift toward soil-centered management. *HortTechnology*, 34(5), 594–603. <https://doi.org/10.21273/HORTTECH05460-24>
- Piottelli, E., Guzzetti, L., Biella, P., Sala, D., Copetta, A., Mussano, P., Galimberti, A., & Labra, M. (2024). Animal pollination shapes fruits market features, seeds functional traits and modulates their chemistry. *Scientific Reports*, 14, 22734. <https://doi.org/10.1038/s41598-024-73647-7>.
- Piovesan, B., Padilha, A. C., Botton, M., & Zotti, M. J. (2019). Entomofauna and potential pollinators of strawberry crop under semi-hydroponic conditions. *Horticultura Brasileira*, 37(3), 324–330. <https://doi.org/10.1590/s0102-053620190311>
- Port, C. M., & Scopes, N. E. A. (1981). Biological control by predatory mites (*Phytoseiulus persimilis* Athias-Henriot) of red spider mite (*Tetranychus urticae* Koch) infesting strawberries grown in walk-in’ plastic tunnels. *Plant Pathology*, 30(2), 95–99. <https://doi.org/10.1111/j.1365-3059.1981.tb01234.x>
- Porter, M., Fan, Z., Lee, S., & Whitaker, V. M. (2023). Strawberry breeding for improved flavor. *Crop Science*, 63(4), 1949–1963. <https://doi.org/10.1002/csc2.21012>
- Prohaska, A., Petit, A., Lesemann, S., Rey-Serra, P., Mazzoni, L., Masny, A., Sánchez-Sevilla, J. F., Potier, A., Gaston, A., Klamkowski, K., Rothan, C., Mezzetti, B., Amaya, I., Olbricht, K., Denoyes, B., & Costa, F. (2024a). Strawberry phenotypic plasticity in flowering time is driven by the interaction between genetic loci

- and temperature. *Journal of Experimental Botany*, 75(18), 5923–5939. <https://doi.org/10.1093/jxb/erae279>
- Prohaska, A., Rey-Serra, P., Petit, J., Petit, A., Perrotte, J., Rothan, C., & Denoyes, B. (2024b). Exploration of a European-centered strawberry diversity panel provides markers and candidate genes for the control of fruit quality traits. *Horticulture Research*, 11, uhae137. <https://doi.org/10.1093/hr/uhae137>
- Rabbi, B., Chen, Z.-H., & Sethuvenkatraman, S. (2019). Protected cropping in warm climates: A review of humidity control and cooling methods. *Energies (Basel)*, 12(14), 2737. <https://doi.org/10.3390/en12142737>
- Raffle, S., Irving, R., & Moore, G. (2010). *Extending the UK strawberry season using a range of plant types and growing systems*. Horticultural Development Company.
- Ramos, G., Goldman, P., Sharrett, J., Sacher, G. O., Pennerman, K. K., Dilla-Ermita, C. J., Jaime, J. H., Steele, M. E., Hewavitharana, S. S., Holmes, G. J., Waterhouse, H., Dundore-Arias, J. P., & Henry, P. (2024). Geospatial analysis of California strawberry fields reveals regional differences in crop rotation patterns and the potential for lengthened rotations at current levels of production. *Frontiers in Sustainable Food Systems*, 8, 1341782. <https://doi.org/10.3389/fsufs.2024.1341782>
- Ranasingha, R., Perera, A., Baghalian, K., & Geroftotis, C. (2024). Amino acid-based biostimulants and microbial biostimulants promote the growth, yield and resilience of strawberries in soilless glasshouse cultivation. *Journal of Sustainable Agriculture and Environment*, 3 (3), e12113. <https://doi.org/10.1002/sae2.12113>
- Rantanen, M., Joensuu, K., Räsänen, K., Silvenius, F., Usva, K., Rikala, K., & Karhu, S. (2024). Life cycle assessment of red raspberry production – a comparative case study of open field and tunnel production in Finland. *Acta Horticulturae*, 1388, 165–170. <https://doi.org/10.17660/ActaHortic.2024.1388.25>
- Requena, M., López-Villén, A., Hernández, A. F., Parrón, T., Navarro, A., & Alarcón, R. (2019). Environmental exposure to pesticides and risk of thyroid diseases. *Toxicology Letters*, 315, 55–63. <https://doi.org/10.1016/j.toxlet.2019.08.017>
- Retamal-Salgado, J., Bastías, R. M., Wilckens, R., & Paulino, L. (2015). Influence of microclimatic conditions under high tunnels on the physiological and productive responses in blueberry ‘O’Neal’. *Chilean Journal of Agricultural Research*, 75(3), 291–297. <https://doi.org/10.4067/S0718-58392015000400004>
- Rho, H., Colaizzi, P., Gray, J., Paetzold, L., Xue, Q., Patil, B., & Rush, C. (2020). Yields, fruit quality, and water use in a jalapeno pepper and tomatoes under open field and high-tunnel production systems in the Texas High Plains. *HortScience*, 55(10), 1632–1641. <https://doi.org/10.21273/HORTSCI15143-20>
- Riikonen, J., Ruhanen, H., Uimari, A., Poteri, M., Toljamo, A., Kokko, H., Blande, J. D., Kumpula, R., & Kivimäenpää, M. (2024). Impact of pre-harvest UVC treatment on powdery mildew infection and strawberry quality in tunnel production in Nordic conditions. *Scientia Horticulturae*, 338, 113706. <https://doi.org/10.1016/j.scienta.2024.113706>
- Robson, T. M., Pieristè, M., Durand, M., Kotilainen, T. K., & Aphalo, P. J. (2022). The benefits of informed management of sunlight in production greenhouses and poly-tunnels. *Plants, People, Planet*, 4, 314–325. <https://doi.org/10.1002/ppp3.10258>
- Roca, J. M., de San Pedro, M., Pérez, M., Marfa, O., Cots, L. L., & Martin-Closas, L. L. (1998). Effects of planting systems on yield, water use efficiency and nutrient balance of strawberry protected crop. *Acta Horticulturae*, 458, 193–200. <https://doi.org/10.17660/ActaHortic.1998.458.22>
- Rogers, M. A., & Wszelaki, A. L. (2012). Influence of high tunnel production and planting date on yield, growth, and early blight development on organically grown heirloom and hybrid tomato. *HortTechnology*, 22(4), 452–462. <https://doi.org/10.21273/HORTTECH.22.4.452>
- Romero-Gámez, M., & Suárez-Rey, E. M. (2020). Environmental footprint of cultivating strawberry in Spain. *The International Journal of Life Cycle Assessment*, 25(4), 719–732. <https://doi.org/10.1007/s11367-020-01740-w>
- Romero-Gámez, M., Suárez-Rey, E. M., Castilla, N., & Soriano, T. (2012). Evaluation of global, photosynthetically active radiation and diffuse radiation transmission of agricultural screens. *Spanish Journal of Agricultural Research*, 10(2), 306–313. <https://doi.org/10.5424/sjar/2012102-260-11>
- Ruiz-González, C., Román, P., Rueda-Ruzafa, L., Cardona, D., Requena, M., & Alarcón, R. (2024). Environmental pesticide exposure and Alzheimer’s disease in southern Spain: A cross-sectional study. *Psychiatry Research*, 337, 115932. <https://doi.org/10.1016/j.psychres.2024.115932>
- Russi, A., Eichelberger Granada, C., & Schwambach, J. (2024). Bacterial agents for controlling anthracnose and soft rot in strawberries: Present status and perspectives. *Biocontrol Science and Technology*, 34, 166–188. <https://doi.org/10.1080/09583157.2024.2307455>
- Ryan, I. C., Shutt, J. D., & Dicks, L. V. (2023). The importance of multi-year studies and commercial yield metrics in measuring pollinator dependence ratios: A case study in UK raspberries *Rubus idaeus* L. *Ecology and Evolution*, 13(5), e10044. <https://doi.org/10.1002/ece3.10044>
- Sabir, N., & Singh, N. (2013). Protected cultivation of vegetables in global arena: A review. *Indian Journal of Agricultural Sciences*, 83(2), 123–135.
- Sakrabani, R., Garnett, K., Knox, J. W., Rickson, J., Pawlett, M., Falagan, N., Girkin, N. T., Cain, M., Alamar, M. C., Burgess, P. J., Harris, J., Patchigolla, K., Sandars, D., Graves, A., Hannam, J., & Simmons, R. W. (2023). Towards net zero in agriculture: Future challenges and opportunities for arable, livestock and protected cropping systems in the UK. *Outlook on Agriculture*, 52(2), 116–125. <https://doi.org/10.1177/00307270231178889>
- Salamé-Donoso, T. P., Santos, B. M., Chandler, C. K., & Sargent, S. A. (2010). Effect of high tunnels on the growth, yields, and soluble solids of strawberry cultivars in Florida. *International Journal of Fruit Science*, 10(3), 249–263. <https://doi.org/10.1080/15538362.2010.510420>
- Salvadores, Y., & Bastías, R. M. (2023). Environmental factors and physiological responses of sweet cherry production under protective cover systems: A review. *Chilean Journal of Agricultural Research*, 83(4), 484–498. <https://doi.org/10.4067/S0718-58392023000400484>
- Sandison, F., Yeluripati, J., & Stewart, D. (2023). Does green vertical farming offer a sustainable alternative to conventional methods of production? A case study from Scotland. *Food and Energy Security*, 12(2), e438. <https://doi.org/10.1002/fes3.438>
- Santos, B. M., Vallad, G., & Torres-Quezada, E. A. (2013). Advances on protected culture of berry crops in Florida. *Journal of the American Pomological Society*, 2013(7), 11–17. <https://doi.org/10.32473/edis-hs1224-2013>
- Sarıdaş, M. A., Karabıyık, Ş., Eti, S., & Paydaş Kargı, S. (2021). Boron applications and bee pollinators increase

- strawberry yields. *International Journal of Fruit Science*, 21(1), 481–491. <https://doi.org/10.1080/15538362.2021.1907010>
- Savage, M. J. (1980). Plastic tunnels: A study of air temperature. *Agrochimophysics*, 12(1), 5–13.
- Savvas, D., Ropokis, A., Ntatsi, G., & Kittas, C. (2016). Current situation of greenhouse vegetable production in Greece. *Acta Horticulturae*, 1142, 443–448. <https://doi.org/10.17660/ActaHortic.2016.1142.67>
- Scarascia-Mugnozza, G., Sica, C., & Russo, G. (2011). Plastic materials in European agriculture: Actual use and perspectives. *Journal of Agricultural Engineering Research*, 42(3), 15–28. <https://doi.org/10.4081/jae.2011.28>
- Schmutz, U., Sumption, P., & Lennartsson, M. (2011). Economics of UK organic protected cropping. *Acta Horticulturae*, 915, 39–46. <https://doi.org/10.17660/ActaHortic.2011.915.4>
- Scott, G., Williams, C., Wallace, R. W., & Du, X. (2021). Exploring plant performance, fruit physicochemical characteristics, volatile profiles, and sensory properties of day-neutral and short-day strawberry cultivars grown in Texas. *Journal of Agricultural and Food Chemistry*, 69(45), 13299–13314. <https://doi.org/10.1021/acs.jafc.1c00915>
- Sengar, S. H., & Kothari, S. (2008). Economic evaluation of greenhouse for cultivation of rose nursery. *African Journal of Agricultural Research*, 3(6), 435–439.
- Sethi, V. P., Dubey, R. K., & Dhath, A. S. (2009). Design and evaluation of modified screen net house for off-season vegetable raising in composite climate. *Energy Conversion and Management*, 50(12), 3112–3128. <https://doi.org/10.1016/j.enconman.2009.08.001>
- Shaik, A., Singh, S., Montague, T., Siebecker, M. G., Ritchie, G., Wallace, R. W., & Stevens, R. (2023). Comparison of organic eggplant yields under open-field and high tunnel production systems in Texas. *Farming System*, 1(3), 100049. <https://doi.org/10.1016/j.farsys.2023.100049>
- Shanthini, K. S., Francis, J., George, S. N., George, S., & Devassy, B. M. (2025). Early bruise detection, classification and prediction in strawberry using Vis-NIR hyperspectral imaging. *Food Control*, 167, 10794. <https://doi.org/10.1016/j.foodcont.2024.110794>
- Sharaf-Eldin, M. A., Yaseen, Z. M., Elmetwalli, A. H., Elsayed, S., Scholz, M., Al-Khafaji, Z., & Omar, G. F. (2023). Modifying walk-in tunnels through solar energy, fogging, and evaporative cooling to mitigate heat stress on tomato. *Horticulturae*, 9(1), 77. <https://doi.org/10.3390/horticulturae9010077>
- Sharma, K. K., Chawla, S., & Singh, M. C. (2024). Protected cultivation: Technology, constraints, and its global status. In M. C. Singh & K. K. Sharma (Eds.), *Protected cultivation: Structural design, crop management, modeling and automation* (pp. 1–34). Apple Academic Press Incorporated.
- Shishido, M. (2011). Plant disease management in protected horticulture. *HortResearch*, 65, 7–18.
- Silva, L. R., Araujo, F. H. V., Ferreira, S. R., Dos Santos, J. C. B., Abreu, C. M., Silva, R. S., & Costa, M. C. (2024). Strawberries in a warming world: Examining the ecological niche of *Fragaria × ananassa* Duch. across different climate scenarios. *Journal of Berry Research*, 14(3), 193–208. <https://doi.org/10.3233/JBR-240012>
- Simane, B., Kumie, A., Berhane, K., Samet, J., Kjellstrom, T., & Patz, J. (2022). Occupational heat stress in the floriculture industry of Ethiopia: Health risks and productivity losses. *Health*, 14(2), 254–271. <https://doi.org/10.4236/health.2022.142020>
- Siopa, C., Carvalheiro, L. G., Castro, H., Loureiro, J., & Castro, S. (2024). Animal-pollinated crops and cultivars—A quantitative assessment of pollinator dependence values and evaluation of methodological approaches. *Journal of Applied Ecology*, 61(6), 1279–1288. <https://doi.org/10.1111/1365-2664.14634>
- Solis-Mera, J. A. (2021). Respuesta de tres cultivares de *Rubus* spp. en sus etapas de crecimiento vegetativo y reproductivo bajo macrotúneles. *Acta Agronómica*, 70(4), 394–406. <https://doi.org/10.15446/acag.v70n4.92460>
- Soode Schimonsky, E., Richter, K., & Weber-Blaschke, G. (2017). Product environmental footprint of strawberries: Case studies in Estonia and Germany. *Journal of Environmental Management*, 203, 564–577. <https://doi.org/10.1016/j.jenvman.2017.03.090>
- Soria, C., Lopez-Aranda, J. M., Medina, J. J., Miranda, L., & Domínguez, F. J. (2009). Evaluation of strawberry production and fruit firmness under small and large plastic tunnels in annual crop system. *Acta Horticulturae*, 842, 119–123. <https://doi.org/10.17660/ActaHortic.2009.842.10>
- Stensvand, A., Wang, N.-Y., Le, V. H., Da Silva, C. D., Jr., Asalf, B., Grieu, C., Turechek, W. W., & Peres, N. A. (2024). Aerated steam eradicates powdery mildew from strawberry transplants. *European Journal of Plant Pathology*, 168(1), 199–205. <https://doi.org/10.1007/s10658-023-02744-6>
- Stone, C. H., Sidhu, R. S., Swarts, N. D., & Close, D. C. (2024). A review of the production challenges for sweet cherries grown under protected cropping systems. *Acta Horticulturae*, 1395, 179–186. <https://doi.org/10.17660/ActaHortic.2024.1395.24>
- Straube, J., Hurtado, G., Zeisler-Diehl, V., Schreiber, L., & Knoche, M. (2024). Cuticle deposition ceases during strawberry fruit development. *BMC Plant Biology*, 24(1), 623. <https://doi.org/10.1186/s12870-024-05327-7>
- Sturiale, S., Gava, O., Gallardo, M., Buendía Guerrero, D., Buyuktas, D., Aslan, G. E., Laarif, A., Bouslama, T., Navarro, A., Incrocci, L., & Bartolini, F. (2024). Environmental and economic performance of greenhouse cropping in the Mediterranean Basin: Lessons learnt from a cross-country comparison. *Sustainability*, 16(11), 4491. <https://doi.org/10.3390/su16114491>
- Subedi, S., Tiwari, N. P., & Subedi, S. (2023). Profitability and determinants of protected vegetable farming in Nepal. *Cogent Food & Agriculture*, 9(1), 2202202. <https://doi.org/10.1080/23311932.2023.2202202>
- Sullivan, C., Black, B., Davis, A., & Sanchez, N. (2022). *Day-neutral strawberry production in central Oregon*. Extension Service, Oregon State University.
- Suminarti, N. E., Sebayang, H. T., Maghofer, M. D., & Bulan, B. (2023). Effect of para-net shade level on plant micro-environment, growth, and yield of three strawberry varieties. *Biodiversitas Journal of Biological Diversity*, 24(4), 2149–2155. <https://doi.org/10.13057/biodiv/d240426>
- Swaen, G. M. H. (2020). Letter to the Editor in response to Requena et al: Environmental exposure to pesticides and risk of thyroid diseases (Requena et al. 2019). *Toxicology Letters*, 331, 122–123. <https://doi.org/10.1016/j.toxlet.2020.05.038>
- Sydorovych, O., Rivard, C. L., O'Connell, S., Harlow, C. D., Peet, M., & Louws, F. J. (2013). Growing organic heirloom tomatoes in the field and high tunnels in North Carolina: Comparative economic analysis. *HortTechnology*, 23(2), 227–236. <https://doi.org/10.21273/HORTTECH.23.2.227>
- Symington, H. A., & Glover, B. J. (2024). Strawberry varieties differ in pollinator-relevant floral traits. *Ecology and*

- Evolution*, 14(2), e10914. <https://doi.org/10.1002/ece3.10914>
- Tabatabaie, S. M. H., & Murthy, G. S. (2016). Cradle to farm gate life cycle assessment of strawberry production in the United States. *Journal of Cleaner Production*, 127, 548–554.
- Talavera, M., Vela, M. D., & Arriaza, M. (2024). Experts' opinion on the sustainable use of nematicides in Mediterranean intensive horticulture. *Spanish Journal of Agricultural Research*, 22(1), e1001. <https://doi.org/10.5424/sjar/2024221-20568>
- Tan, C. S., Papadopoulos, A. P., & Liptay, A. (1984). Effect of various types of plastic films on the soil and air temperatures in 80-cm high tunnels. *Scientia Horticulturae*, 23(2), 105–112. [https://doi.org/10.1016/0304-4238\(84\)90013-X](https://doi.org/10.1016/0304-4238(84)90013-X)
- Taniguchi, H., Tsukuda, Y., Motoki, K., Goto, T., Yoshida, Y., & Yasuba, K. (2025). Development of an AI-based image analysis system to calculate the visit duration of a green blow fly on a strawberry flower. *The Horticulture Journal*, 94(1), 64–72. <https://doi.org/10.2503/hortj.QH-159>
- Thipe, E. L., Workneh, T. S., Odindo, A. O., & Laing, M. D. (2020). A comparison of two greenhouse structures under sub-humid conditions in terms of changes in temperature and relative humidity. *Acta Horticulturae*, 1271, 9–16. <https://doi.org/10.17660/ActaHortic.2020.1271.2>
- Thomasz, E. O., Kasanzew, A., Massot, J. M., & García-García, A. (2024). Valuing ecosystem services in agricultural production in southwest Spain. *Ecosystem Services*, 68, 101636. <https://doi.org/10.1016/j.ecoser.2024.101636>
- Tian, S., Zhao, X., Vincent, I. R., Gong, T., Ray, Z. T., Legaspi, J., Bolques, A., Coolong, T. W., & Diaz-Pérez, J. C. (2023). *Using high tunnels to enhance organic vegetable production in Florida: An overview*. Extension Publication, Institute for Food and Agricultural Science, University of Florida, Number HS1466.
- Tiwari, P., Shrivastava, A. K., & Dave, A. K. (2023). Occupational heat stress to the greenhouse workers. *The Pharma Innovation Journal*, SP-12, 3–5.
- Tuohimetsä, S., Hietaranta, T., Uosukainen, M., Kukkonen, S., & Karhu, S. (2014). Fruit development in artificially self- and cross-pollinated strawberries (*Fragaria × ananassa*) and raspberries (*Rubus idaeus*). *Acta Agriculturae Scandinavica, Section B — Soil and Plant Science*, 64(5), 408–415. <https://doi.org/10.1080/09064710.2014.919348>
- Uchanski, M. E., VanLeeuwen, D. M., Guldán, S. J., Falk, C. L., Shukla, M., & Enfield, J. (2020). Temperature and light characterization during winter production season in high tunnels in the Southwestern United States. *HortTechnology*, 30(2), 259–267. <https://doi.org/10.21273/HORTTECH04486-19>
- Ullah, I., Toor, M. D., Yerlikaya, B. A., Mohamed, H. I., Yerlikaya, S., Basit, A., & Rehman, A. (2024). High-temperature stress in strawberry: Understanding physiological, biochemical and molecular responses. *Planta*, 260, 118. <https://doi.org/10.1007/s00425-024-04544-6>
- Unnikrishnan, P., Ponnambalam, K., & Karray, F. (2024). Influence of regional temperature anomalies on strawberry yield: A study using multivariate copula analysis. *Sustainability*, 16(9), 3523. <https://doi.org/10.3390/su16093523>
- Vandecasteele, B., Claerbout, J., Denaeghel, H., & Craeye, S. (2024). The repeatability of reusing peat as horticultural substrate and the role of fertigation for optimal reuse. *Waste Management*, 190, 296–305. <https://doi.org/10.1016/j.wasman.2024.09.028>
- Verteramo Chiu, L. J., Nicholson, C. F., Gómez, M. I., & Mattson, N. S. (2024). A meta-analysis of yields and environmental performance of controlled-environment production systems for tomatoes, lettuce and strawberries. *Journal of Cleaner Production*, 469, 143142. <https://doi.org/10.1016/j.jclepro.2024.143142>
- Villagrán, E., Flores-Velazquez, J., Akrami, M., & Bojacá, C. (2021). Influence of the height in a Colombian multi-tunnel greenhouse on natural ventilation and thermal behavior: Modeling approach. *Sustainability*, 13(24), 13631. <https://doi.org/10.3390/su132413631>
- Ward, M. J., & Bomford, M. K. (2013). Row covers moderate diurnal temperature flux in high tunnels. *Acta Horticulturae*, 987, 59–66. <https://doi.org/10.17660/ActaHortic.2013.987.8>
- Ward, R., Drost, D., & Whyte, A. (2011). Assessing profitability of selected specialty crops grown in high tunnels. *Journal of Agribusiness*, 29(1), 43–58. <https://doi.org/10.22004/ag.econ.260146>
- Warner, D. J., Davies, M., Hipps, N., Osborne, N., Tzilivakis, J., & Lewis, K. A. (2010). Greenhouse gas emissions and energy use in UK-grown short-day strawberry (*Fragaria x ananassa* Duch.) crops. *The Journal of Agricultural Science*, 148(6), 667–681. <https://doi.org/10.1017/S0021859610000493>
- Warren, R. J., Colin, T., Quarrell, S. R., Barron, A. B., & Allen, G. R. (2024). Quantifying the impact of crop coverings on honey bee orientation and foraging in sweet cherry orchards using RFID. *Journal of Applied Entomology*, 148(5), 566–581. <https://doi.org/10.1111/jen.13247>
- Waterer, D. (2003). Yields and economics of high tunnels for production of warm-season vegetable crops. *HortTechnology*, 13(2), 339–343. <https://doi.org/10.21273/HORTTECH.13.2.0339>
- Webb, J., Williams, A. G., Hope, E., Evans, D., & Moorhouse, E. (2013). Do foods imported into the UK have a greater environmental impact than the same foods produced within the UK? *The International Journal of Life Cycle Assessment*, 18(7), 1325–1343. <https://doi.org/10.1007/s11367-013-0576-2>
- Wien, H. C. (2009). Microenvironmental variations within the high tunnel. *HortScience*, 44(2), 235–238. <https://doi.org/10.21273/HORTSCI.44.2.235>
- Wien, H. C., & Pritts, M. P. (2009). Use of high tunnels in the Northeastern USA: Adaptation to cold climates. *Acta Horticulturae*, 807, 55–60. <https://doi.org/10.17660/ActaHortic.2009.807.3>
- Wilkanić, Z., & Radajewska, B. (1997). Solitary bee *Osmia rufa* L. (Apoidea, Megachilidae) as pollinator of strawberry cultivated in an unheated plastic tunnel. *Acta Horticulturae*, 439, 489–494. <https://doi.org/10.17660/ActaHortic.1997.439.83>
- Willden, S. A., Pritts, M. P., & Loeb, G. M. (2022). The effect of plastic low tunnels on natural enemies and pollinators in New York strawberry. *Crop Protection*, 151, 105820. <https://doi.org/10.1016/j.cropro.2021.105820>
- Wszelaki, A., Lockwood, D., & Martin, J. (2013). *High tunnel strawberry production in Tennessee*. Institute of Agriculture, University of Tennessee.
- Xiao, C. L., Chandler, C. K., Price, J. F., Duval, J. R., Mertely, J. C., & Legard, D. E. (2001). Comparison of epidemics of botrytis fruit rot and powdery mildew of strawberry in large plastic tunnel and field production systems. *Plant Disease*, 85(8), 901–909. <https://doi.org/10.1094/PDIS.2001.85.8.901>
- Xu, S., Feng, Y., Han, L., Ran, X., Zhong, Y., Jin, Y., & Song, J. (2023). Evaluation of the wind field and deposition effect of

- a novel air-assisted strawberry sprayer. *Agriculture*, 13(2), 230. <https://doi.org/10.3390/agriculture13020230>
- Yamanaka, R., & Kawashima, H. (2021). Development of cooling techniques for small-scale protected horticulture in mountainous areas in Japan. *Japan Agricultural Research Quarterly*, 55(2), 117–125. <https://doi.org/10.6090/jarq.55.117>
- Yao, S., Guldán, S., & Heyduck, R. (2019). High tunnel apricot production in frost-prone northern New Mexico. *HortTechnology*, 29(4), 457–460. <https://doi.org/10.21273/HORTTECH04315-19>
- Yoneda, Y., & Kawashima, H. (2024). An evaluation of the ventilation performance using a new structure of plastic low tunnels. *Acta Horticulturae*, 1404, 293–300. <https://doi.org/10.17660/ActaHortic.2024.1404.40>
- Yousef, S. A. M., Ali, A. M., Elsherbiny, E. A., & Atwa, A. A. (2024). Morphological, genetic and pathogenic variability among *Botrytis cinerea* species complex causing gray mold of strawberry. *Physiological and Molecular Plant Pathology*, 134, 102395. <https://doi.org/10.1016/j.pmpp.2024.102395>
- Yuan, B. Z., Sun, J., & Nishiyama, S. (2004). Effect of drip irrigation on strawberry growth and yield inside a plastic greenhouse. *Biosystems Engineering*, 87(2), 237–245. <https://doi.org/10.1016/j.biosystemseng.2003.10.014>
- Zhang, Q., Zhang, X., Yang, Z., Huang, Q., & Qiu, R. (2022). Characteristics of plastic greenhouse high-temperature and high-humidity events and their impacts on facility tomatoes growth. *Frontiers in Earth Science*, 10, 848924. <https://doi.org/10.3389/feart.2022.848924>
- Zhang, Q., Zhu, J., Yu, X., Huang, S., Zhang, X., Zhang, S., Qiu, R., & Agathokleous, E. (2023). Assessing suitability of major meteorological factors for facility agriculture in mainland China. *Environmental Research Letters*, 18(11), 114002. <https://doi.org/10.1088/1748-9326/acffe0>
- Zhao, X., Gao, Z., & Black, Z. (2015). An economic analysis of organic strawberry production in the high tunnel system. *Proceedings of the Florida State Horticultural Society*, 128, 138–141.
- Zheng, C., Abd-Elrahman, A., Whitaker, V. M., Wang, X., Dalid, C., & Shen, K. (2024a). Strawberry canopy structural parameters estimation and growth analysis from UAV multispectral imagery using a geospatial tool. *Computers and Electronics in Agriculture*, 226, 109440. <https://doi.org/10.1016/j.compag.2024.109440>
- Zheng, M. Z., Leib, B., Butler, D. B., Wright, W., Ayers, P., Hayes, D., Haghverdi, A., Feng, L. S., Grant, T., Vanchiasong, P., & Muchoki, D. (2019). Assessing heat management practices in high tunnels to improve organic production of bell peppers. *Scientia Horticulturae*, 246, 928–941. <https://doi.org/10.1016/j.scienta.2018.10.046>
- Zheng, M. Z., Leib, B., Butler, D. M., Wright, W., Ayers, P., & Hayes, D. (2017). Modeling energy balance and airflow characteristics in a naturally ventilated high tunnel. *Transactions of the ASABE*, 60(5), 1683–1697. <https://doi.org/10.13031/trans.12080>
- Zheng, R., Romero-Del Rey, R., Garcia-Gonzalez, J., Requena-Mullor, M., Navarro-Mena, A., López-Villén, A., & Alarcon-Rodriguez, R. (2024b). Indicators of occupational pesticide exposure are associated with psychiatric symptoms. *Environmental Toxicology and Pharmacology*, 105, 104357. <https://doi.org/10.1016/j.etap.2023.104357>